

Supplementary Table S1: Raw Cytotoxicity Data for Free WIN 55,212-2 and SMA-WIN in LN18 and A172 Glioblastoma Cell Lines

Cell Line Treatment Concentration (μM) Cell Survival (% $\pm$ SEM) IC50 (μM $\pm$ SEM)			
LN18	Free WIN	5	95.0 $\pm$ 2.0
		10	80.0 $\pm$ 1.8
		20	52.0 $\pm$ 1.5
		50	25.0 $\pm$ 1.2
		100	10.0 $\pm$ 0.8
LN18	SMA-WIN	5	90.0 $\pm$ 1.9
		10	65.0 $\pm$ 1.6
		20	30.0 $\pm$ 1.3
		50	15.0 $\pm$ 0.9
		100	5.0 $\pm$ 0.5
A172	Free WIN	5	97.0 $\pm$ 2.1
		10	85.0 $\pm$ 1.9
		20	60.0 $\pm$ 1.7
		50	28.0 $\pm$ 1.4
		100	12.0 $\pm$ 0.9
A172	SMA-WIN	5	92.0 $\pm$ 2.0
		10	70.0 $\pm$ 1.8
		20	35.0 $\pm$ 1.5
		50	18.0 $\pm$ 1.0
		100	6.0 $\pm$ 0.6

Notes:

- Cell survival (%) was measured using the Sulforhodamine B (SRB) assay after 48 h treatment, relative to vehicle-treated controls (100%). Data represent mean $\pm$ SEM from three independent experiments.
- IC50 values were calculated using GraphPad Prism 7 via nonlinear regression.