

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Blood and Urinary Metal Levels among Exclusive Marijuana Users in NHANES (2005–2018)

Katlyn E. McGraw, Anne E. Nigra, Joshua Klett, Marisa Sobel, Elizabeth C. Oelsner, Ana Navas-Acien, Xin Hu, and Tiffany R. Sanchez

Table of Contents

Figure S1. Flow chart of NHANES 2005-2018 participants and exclusion criteria.

Table S1. Blood (LBX prefix) and urinary (URX prefix) metal data availability by NHANES cycle year. Seven cycles were included in our data analysis. Grey boxes indicate that metal biomarker was not collected for that cycle year and x indicates that metal biomarker was collected.

Table S2. Lower limit of detection (LLOD) values and % below limit of detection (LOD) for metal biomarkers in blood (LBX prefix) and urine (URX prefix) by NHANES cycle year. Metal biomarkers were reported in µg/L in urine and whole blood, except for lead in whole blood which is reported in µg/dL. Grayed out lines indicate metal biomarkers not measured in that cycle.

Table S3. Self-reported cigarette smoking and serum cotinine values (ng/mL) in a 2x2 table.

Table S4. Marijuana and tobacco use categories and recent use categories in a 2x2 table.

Table S5. Participant characteristics across categories of recent marijuana use, restricted to participants with available drug use questionnaire data and urinary metals (n=5,024). Self-reported cigarette smokers and individuals with serum cotinine >10ng/mL were excluded. Continuous variables are reported as mean (SD) and categorical variables as number (%).

Table S6. Urinary metal concentrations ($\mu\text{g/g}$ creatinine) by recent use, reported as median [IQR]. Metal levels measured in urine ($\mu\text{g/g}$) and in blood ($\mu\text{g/L}$ or $\mu\text{g/dL}$ for Pb), N= 5,024 NHANES 2005-2018 participants. The reference group of non-use includes individuals who neither use marijuana or tobacco nor have not used marijuana in more than one year (N=4,455). Groups of individuals who had recently used marijuana include those who had used in the last 7 days (N=226), those who had used in the last 7-30 days (N=132), or those who had used in the 31-365 days (N=211). Individuals who self-reported cigarette smoking or had serum cotinine levels $>10\text{ng/mL}$ were excluded. Values of Hg only include women of childbearing age.

Table S7. Arithmetic mean differences and 95% CIs in blood metal concentrations ($\mu\text{g/L}$ or $\mu\text{g/dL}$ for Pb) across different marijuana use categories as compared to non-marijuana/non-tobacco use (reference), N= 7,254 NHANES 2005-2018 participants, corresponding to Figure 1 in the manuscript.

Table S8. Arithmetic mean differences and 95% CIs in urinary metal levels ($\mu\text{g/g}$) across different marijuana use as compared to non-marijuana/non-tobacco use (reference), N= 7,254 NHANES 2005-2018 participants, corresponding to Figure 2 in the manuscript.

Table S9. Arithmetic mean differences and 95% CIs in blood metal concentrations ($\mu\text{g/L}$ or $\mu\text{g/dL}$ for Pb) in individuals who recently used marijuana as compared to individuals who do not use or had not used in more than one year (reference), N= 5,024 NHANES 2005-2018 participants, corresponding values for Figure 3 in the manuscript.

Table S10. Arithmetic mean differences and 95% CIs in urinary metal levels ($\mu\text{g/g}$) in individuals who recently used marijuana as compared to individuals who do not use or had not used in more than one year (reference), N= 5,024 NHANES 2005-2018 participants, corresponding values for Figure 4 in the manuscript.

Figure S2. Mean differences and 95% CIs in blood metal concentrations ($\mu\text{g/L}$ or $\mu\text{g/dL}$ for Pb) across different marijuana use groups analyzed by subgroup of age, sex, and race, N= 7,254 NHANES 2005-2018 participants.

Figure S3. Mean differences and 95% CIs in urinary metal levels ($\mu\text{g/g}$) across different marijuana use groups analyzed by subgroup of age, sex, and race, N= 7,254 NHANES 2005-2018 participants.

Figure S4. Mean differences and 95% CIs in blood metal concentrations ($\mu\text{g/L}$ or $\mu\text{g/dL}$ for Pb) in individuals who recently used marijuana analyzed by subgroup of age, sex, and race/ethnicity, N= 5,024 NHANES 2005-2018 participants.

Figure S5. Mean differences and 95% CIs in urinary metal levels ($\mu\text{g/g}$) in individuals who recently used marijuana analyzed by subgroup of age, sex, and race/ethnicity, N= 5,024 NHANES 2005-2018 participants.

Figure S6. Mean differences and 95%CIs in blood metal levels ($\mu\text{g/L}$ or $\mu\text{g/dL}$ for Pb) across different marijuana frequencies among exclusive marijuana users, N= 358 NHANES 2005-2018 participants.

Figure S7. Mean differences and 95%CIs in urinary metal levels ($\mu\text{g/g}$ of creatinine) across different marijuana frequencies among exclusive marijuana users, N= 358 NHANES 2005-2018 participants.

Figure S8. Mean differences and 95%CIs in blood metal levels ($\mu\text{g/L}$ or $\mu\text{g/dL}$ for Pb) across different marijuana use groups with and without further adjustment for race and ethnicity in the model, N= 7,254 NHANES 2005-2018 participants.

Figure S9. Mean differences and 95%CIs in urinary metal levels ($\mu\text{g/g}$ creatinine) across different marijuana use groups with and without further adjustment for race and ethnicity in the model, N= 7,254 NHANES 2005-2018 participants.

Figure S10. Mean differences and 95%CIs in blood metal levels ($\mu\text{g/L}$ or $\mu\text{g/dL}$ for Pb) in individuals who recently used marijuana with and without further regression adjustment for race and ethnicity, N= 5,024 NHANES 2005-2018 participants.

Figure S11. Mean differences and 95%CIs in urinary metal levels ($\mu\text{g/g}$ creatinine) in individuals who recently used marijuana with and without further regression adjustment for race and ethnicity, N= 5,024 NHANES 2005-2018 participants.

Figure S12. Mean differences and 95%CIs in urinary metal levels ($\mu\text{g/g}$ creatinine) across different marijuana use groups with and without further adjustment for urinary creatinine in the model, N= 7,254 NHANES 2005-2018 participants.

Figure S13. Mean differences and 95%CIs in urinary metal levels ($\mu\text{g/g}$ creatinine) in individuals who recently used marijuana with and without further adjustment for urinary creatinine in the model, N= 5,024 NHANES 2005-2018 participants.

Figure S14. Mean differences and 95%CIs in blood metal levels ($\mu\text{g/L}$ or $\mu\text{g/dL}$ for Pb) across different marijuana use with and without further regression adjustment for eGFR, N= 7,254 NHANES 2005-2018 participants.

Figure S15. Mean differences and 95%CIs in urinary metal levels ($\mu\text{g/g}$ creatinine) across different marijuana use with and without further regression adjustment for eGFR, N= 7,254 NHANES 2005-2018 participants.

Figure S16. Mean differences and 95%CIs in blood metal levels ($\mu\text{g/L}$ or $\mu\text{g/dL}$ for Pb) in individuals who recently used marijuana with and without further regression adjustment for eGFR, N= 5,024 NHANES 2005-2018 participants.

Figure S17. Mean differences and 95%CIs in urinary metal levels ($\mu\text{g/g}$ creatinine) in individuals who recently used marijuana with and without further regression adjustment for eGFR, N= 5,024 NHANES 2005-2018 participants.

Figure S18. Mean differences and 95%CIs in blood metal levels across different marijuana use with and without further regression adjustment for serum cotinine, N= 7,254 NHANES 2005-2018 participants.

Figure S19. Mean differences and 95%CIs in urinary metal levels ($\mu\text{g/g}$ creatinine) across different marijuana use with and without further regression adjustment for serum cotinine, N= 7,254 NHANES 2005-2018 participants.

Figure S20. Mean differences and 95%CIs in blood metal levels in individuals who recently used marijuana with and without further regression adjustment for serum cotinine, N= 5,024 NHANES 2005-2018 participants.

Figure S21. Mean differences and 95%CIs in urinary metal levels ($\mu\text{g/g}$ creatinine) in individuals who recently used marijuana with and without further regression adjustment for serum cotinine, N= 5,024 NHANES 2005-2018 participants.

Figure S22. Mean differences and 95%CIs in urinary and blood metal levels (UCd: urinary cadmium levels ($\mu\text{g/g}$), BCd: blood cadmium levels ($\mu\text{g/L}$), UPb: urinary lead levels ($\mu\text{g/g}$), and BPb: blood lead levels ($\mu\text{g/dL}$)) across different marijuana use groups, stratified by former smoking status, never smoking and ever smoking, in N= 7,246 NHANES 2005-2018 participants.

Figure S23. Mean differences and 95%CIs in urinary and blood metal levels (UCd: urinary cadmium levels ($\mu\text{g/g}$), BCd: blood cadmium levels ($\mu\text{g/L}$), UPb: urinary lead levels ($\mu\text{g/g}$), and BPb: blood lead levels ($\mu\text{g/dL}$)) in individuals who recently used marijuana, stratified by former smoking status, never smoking and ever smoking, in N= 5,024 NHANES 2005-2018 participants.

Figure S24. Mean differences and 95%CIs in blood metal levels across different marijuana use with and without further adjustment for different measures of seafood consumption.

Figure S25. Mean differences and 95%CIs in urinary metal levels across different marijuana use with and without further adjustment for different measures of seafood consumption.

Figure S26. Mean differences and 95%CIs in blood metal levels in individuals who recently used with and without further adjustment for different measures of seafood consumption.

Figure S27. Mean differences and 95%CIs in urinary metal levels in individuals who recently used with and without further adjustment for different measures of seafood consumption.

Additional File- Excel Document