

Assessing the Effects of a Diet of BPA Analogues-Exposed Microalgae in the Clam *Ruditapes philippinarum*

Jacopo Fabrello, Michela Dalla Fontana, Noemi Gaiani, Maria Piscato, Marco Roverso, Sara Bogiatti and Valerio Matozzo




Figure S1: haemocyte diameter, expressed as µm. Different letters indicate significant differences among all treatments. N=5.




Figure S2: haemocyte volume, expressed as pL. Different letters indicate significant differences among all treatments. N=5.




Figure S3: LDH activity expressed, as OD₄₉₀ nm. Different letters indicate significant differences among all treatments. N=5.




Figure S4: lysozyme activity in HL, expressed as µg lysozyme/mg protein. Different letters indicate significant differences among all treatments. N=5.




Figure S5: arylsulfatase activity in HL, expressed as $\mu\text{g p-nitrocatechol/h/mg protein}$. Different letters indicate significant differences among all treatments. N=5.




Figure S6: acid phosphatase activity in HL, expressed as U/mg protein . Different letters indicate significant differences among all treatments. N=5.




Figure S7: alkaline phosphatase activity in CFH, expressed as U/mg protein. Different letters indicate significant differences among all treatments. N=5.




Figure S8: CUPRAC levels in haemolymph, expressed as mM Trolox Eq/mg protein. Different letters indicate significant differences among all treatments. N=5.




Figure S9: CUPRAC levels in digestive gland, expressed as mM Trolox Eq./mg protein. Different letters indicate significant differences among all treatments. N=5.




Figure S10: SOD activity in gills, expressed as U SOD/mg protein. Different letters indicate significant differences among all treatments. N=5.




Figure S11: CAT activity in the digestive gland, expressed as U CAT/mg protein. Different letters indicate significant differences among all treatments. N=5.




Figure S12: GST activity in the digestive gland, expressed as nmol/min/mg protein. Different letters indicate significant differences among all treatments. N=5.




Figure S13: PCC levels in gills expressed as nmol/mg protein. Different letters indicate significant differences among all treatments. N=5.




Figure S14: AChE activity in gills, expressed as nmol/min/mg protein. Different letters indicate significant differences among all treatments. N=5.




Figure S15: BChE activity in gills, expressed as nmol/min/mg protein. Different letters indicate significant differences among all treatments. N=5.




Figure S16: HAT activity in gills, expressed as μmol TNB⁻/mg protein. Different letters indicate significant differences among all treatments. N=5.