

Supporting Information

Physicochemical characterization and metabolites identification of the synthetic cannabinoid MDMB-5'Br-PINACA using in silico and in vitro approaches

Alexandre B. Godoi^{†,‡}, Natalícia J. Antunes^{*§}, Júlio C. C. da Silva^{||}, Gabriel Cordeiro^{†,‡}, Tássia F. D. Castro^{†,‡}, José L. Costa^{†,§}

[†] Campinas Poison Control Center, Universidade Estadual de Campinas (UNICAMP), Campinas, SP 13083-859, Brazil

[‡] School of Medical Sciences, Universidade Estadual de Campinas (UNICAMP), Campinas, SP 13083-859, Brazil

[§] Faculty of Pharmaceutical Sciences, Universidade Estadual de Campinas (UNICAMP), Campinas, SP 13083-859, Brazil

^{||} Nova Analítica Imp. Exp. LTDA, Sao Paulo, SP 09941-202, Brazil

***Corresponding author:** Faculty of Pharmaceutical Sciences, Universidade Estadual de Campinas (UNICAMP), R. Cândido Portinari, 200, Cidade Universitária, 13083-871, Campinas, SP, Brazil.

E-mail address: natja@unicamp.br (Antunes, N.J.)

Tel.: +55-19-3521-8181

ORCID: <https://orcid.org/0000-0002-7820-552X>

FIGURES

Figure S1. MS ¹ -mass spectra of MDMB-5'Br-PINACA metabolites considering phase I metabolism in pooled human liver microsomes (pHLM) by LC-HRMS using electrospray ionization in positive mode (ESI ⁺).....	3
Figure S2. MS ² -mass spectra of MDMB-5'Br-PINACA metabolites considering phase I metabolism in pooled human liver microsomes (pHLM) by LC-HRMS using electrospray ionization in positive mode (ESI ⁺).....	10




Figure S1. MS¹-mass spectra of MDMB-5'Br-PINACA metabolites considering phase I metabolism in pooled human liver microsomes (pHLM) by LC-HRMS using electrospray ionization in positive mode (ESI⁺).






Figure S1 (Continuation). MS¹-mass spectra of MDMB-5'-Br-PINACA metabolites considering phase I metabolism in pooled human liver microsomes (pHLM) by LC-HRMS using electrospray ionization in positive mode (ESI⁺).







Figure S1 (Continuation). MS¹-mass spectra of MDMB-5'-Br-PINACA metabolites considering phase I metabolism in pooled human liver microsomes (pHLM) by LC-HRMS using electrospray ionization in positive mode (ESI⁺).




Figure S1 (Continuation). MS¹-mass spectra of MDMB-5'Br-PINACA metabolites considering phase I metabolism in pooled human liver microsomes (pHLM) by LC-HRMS using electrospray ionization in positive mode (ESI⁺).







Figure S1 (Continuation). MS¹-mass spectra of MDMB-5'-Br-PINACA metabolites considering phase I metabolism in pooled human liver microsomes (pHLM) by LC-HRMS using electrospray ionization in positive mode (ESI⁺).




Figure S1 (Continuation). MS¹-mass spectra of MDMB-5'Br-PINACA metabolites considering phase I metabolism in pooled human liver microsomes (pHLM) by LC-HRMS using electrospray ionization in positive mode (ESI⁺).




Figure S1 (Continuation). MS¹-mass spectra of MDMB-5'Br-PINACA metabolites considering phase I metabolism in pooled human liver microsomes (pHLM) by LC-HRMS using electrospray ionization in positive mode (ESI⁺).




Figure S2. MS²-mass spectra of MDMB-5'Br-PINACA metabolites considering phase I metabolism in pooled human liver microsomes (pHLM) by LC-HRMS using electrospray ionization in positive mode (ESI⁺).




Figure S2 (Continuation). MS²-mass spectra of MDMB-5'Br-PINACA metabolites considering phase I metabolism in pooled human liver microsomes (pHLM) by LC-HRMS using electrospray ionization in positive mode (ESI⁺).




Figure S2 (Continuation). MS²-mass spectra of MDMB-5'Br-PINACA metabolites considering phase I metabolism in pooled human liver microsomes (pHLM) by LC-HRMS using electrospray ionization in positive mode (ESI⁺).




Figure S2 (Continuation). MS²-mass spectra of MDMB-5'Br-PINACA metabolites considering phase I metabolism in pooled human liver microsomes (pHLM) by LC-HRMS using electrospray ionization in positive mode (ESI⁺).




Figure S2 (Continuation). MS²-mass spectra of MDMB-5'Br-PINACA metabolites considering phase I metabolism in pooled human liver microsomes (pHLM) by LC-HRMS using electrospray ionization in positive mode (ESI⁺).




Figure S2 (Continuation). MS²-mass spectra of MDMB-5'Br-PINACA metabolites considering phase I metabolism in pooled human liver microsomes (pHLM) by LC-HRMS using electrospray ionization in positive mode (ESI⁺).




Figure S2 (Continuation). MS²-mass spectra of MDMB-5'Br-PINACA metabolites considering phase I metabolism in pooled human liver microsomes (pHLM) by LC-HRMS using electrospray ionization in positive mode (ESI⁺).