

Table S1. Frequency of drug use and number of users (*n*) by trimester in +MJ and –MJ groups

	Trimester	+MJ				-MJ				<i>t</i>	<i>p</i>
		<i>n</i>	Mean	SE	Range	<i>n</i>	Mean	SE	Range		
Marijuana	1	20	13.29	5.5	0-112	0					
Joints/week	2	20	9.12	5.3	0-104	0					
	3	18	5.38	3.5	0-69	0					
	Postnatal	16	0.34	0.28	0-4.5	0					
Nicotine	1	14	10.53	1.44	3-20	15	9.77	2.26	0.02 - 40	-0.13	0.899
Cigarettes/day	2	13	6.21	1.38	0.01-17	11	10.12	3.24	1.47 - 40	-0.65	0.521
	3	8	4.90	1.45	0.65-12	11	6.04	1.38	0.50 - 16	-0.98	0.336
	Postnatal	6	4.7	2.38	0.06-15	7	8.45	2.38	0.12 - 20	-1.19	0.245
Alcohol	1	9	15.23	9.50	0.08-84	6	6.44	4.04	0.08 - 26	1.00	0.347
Drinks/week	2	3	0.41	0.18	0.07-0.7	4	4.62	2.64	0.13 - 12	-1.38	0.187
	3	2	0.43	0.07	0.36-0.5	3	5.94	3.39	0.69 - 12	-1.32	0.204
	Postnatal	1	0.50			2	2.33	1.92	0.42 – 4	-0.97	0.343
Opiates *	1	0				4	14.88	4.60	7.0 - 24.5		
Times /	2	1	2.0			3	12.83	5.83	7.0 - 24.5		
trimester	3	0				2	49.0	42.0	7.0 - 91.0		
	Postnatal	0				2	19.0	12.0	7.0 - 31.0		

Postnatal: postnatal period before MRI visit; SE: Standard Error; *n*: number of subjects in a group using the drug during that trimester. *The small number of opiate users in each group prevented valid statistical comparison.