

A 20th-century *Cannabis* pollen increase in a Pyrenean varved record and its potential causes

Valentí Rull^{1,2*}, Teresa Vegas-Vilarrúbia³

¹Spanish National Research Council, 08038 Barcelona Spain

²Catalan Institute of Paleontology, 08193 Cerdanyola del Vallès, Barcelona, Spain

³Department of Evolutionary Biology, Ecology and Environmental Sciences, Universitat de Barcelona, 08028 Barcelona, Spain

*Corresponding author: vrull@csic.es

Abstract

The continuous, varved and absolutely dated sedimentary record of Lake Montcortès (Iberian Pyrenees) has provided evidence for a distinct and characteristic 20th century (1980s) increase in *Cannabis* pollen (20C) that persists today. This event was coeval with the geographical shift of the hemp production center in the Iberian Peninsula from east to northeast (where Lake Montcortès lies), which was accompanied by a significant production increase. This increasing trend was fostered by the renewed interest of the paper industry in hemp and was promoted by the onset of European Union subsidies to hemp cultivation. Illegal cannabis crops could have also contributed to the *Cannabis* pollen increase, but sound evidence is still lacking. These preliminary conclusions should be reinforced by increasing the resolution of the current palynological record and modeling the dispersal of *Cannabis* pollen around the Montcortès region. More similar high-resolution records are needed to verify the geographical extent of the 20C event. Additionally, Lake Montcortès varved sediments are proposed as a suitable candidate to characterize the onset of the “Anthropocene” epoch (mid-20th century), as currently defined by the Anthropocene Working Group.

Keywords: *Cannabis*, hemp, pollen, cultivation, retting, 20th century, Pyrenees, Iberian Peninsula

1. Introduction

Cannabis is among the earliest human domesticates and has been an integral part of human life for millennia (Rull, 2022). Almost every part of this plant is able to provide a variety of products and services, such as textiles, cordage, fuel, paper, medicine, oil, food, drugs or biofuel, along with recreational, ritual, aesthetic and educational materials, among others (Clarke & Merlin, 2013, 2016; Gray et al., 2016). A database on the traditional uses of *Cannabis* (CANNUSE) is available at <http://cannusedb.csic.es/> (last visited 24 March 2023) (Balant et al., 2021). Taxonomically, the genus *Cannabis* has a single species, *Cannabis sativa*, subdivided into two subspecies, *C. sativa* subsp. *sativa* and *C. sativa* subsp. *indica*. However, as usual in plants intensively used by humans, a number of new varieties have been created by artificial selection to adapt the plant to a diversity of biogeographical features and cultural needs (Barcaccia et al., 2020; Kovalchuk et al., 2020). Due to the illegal condition of this plant in most Western countries, the scientific study of *Cannabis* underwent a significant decline in the mid-20th century (Duvall, 2014; Warf, 2014). However, the last decade has witnessed a revival of interest in *Cannabis*, especially in relation to the genetic and phytochemical characteristics of its different varieties, as well as its evolutionary origin, domestication and further diffusion. This resurgence is linked to a renewed interest in crop improvement, especially for medical purposes but also for other applications (Gray et al., 2016).

The Iberian Peninsula (IP) has been considered a keystone area for the natural and anthropogenic diffusion of *Cannabis* due to its peculiar biogeographical position and its diverse and dynamic cultural history (Rull, 2022). According to the recent meta-analysis of a database (CHIP, for *Cannabis-Humulus* Iberian Peninsula) with nearly 60 pollen records widespread across the IP (Rull et al., 2023), *Cannabis* would have entered the IP in its wild form by the Late Pleistocene (150-12 kyr BP). Domesticated forms would have arrived later in two main dispersal waves, the first in the Neolithic (7-5 kyr BP) and the second in the Middle Ages (500 CE onward), via continental (Europe) and maritime (Mediterranean) pathways. Maximum cultivation and hemp retting activities were recorded during the Modern Age (16th-19th centuries). A unique *Cannabis* record was obtained in the annually-laminated (varved) sediments of Lake Montcortès, situated in the northeastern sector (Fig. 1), which contains the longest (last 3000 years), continuous (gap-free) and high-resolution (bidecadal) absolutely dated pollen sequence available for the Mediterranean region (Rull et al., 2021). In this record, *Cannabis* appeared in the early Middle Ages (~600 CE) and has continued until today, showing significant variations, which have been correlated with historical sociocultural developments.




Figure 1. Left. Map of the Iberian Peninsula and the Balearic Islands (left), indicating the localities with published pollen records (white dots) (Carrión et al., 2022) and those containing *Cannabis* pollen (red dots) (Rull et al., 2023). Yellow dots indicate the sites with *Cannabis* records in the 20th century (AD, Alcúdia; CE, Cueto Espina; EN, Estanyons; ET, Estanya) (Rull et al., 2023). The location of Lake Montcortès (LM) is indicated by a yellow star. Blue boxes are the cities with aerobiological studies on *Cannabis* pollen mentioned in the text (CO, Córdoba; MA, Málaga; MU, Murcia). Blue circles are the main areas of hemp production, as explained in the text (AL, Alicante; CT, Catalonia). Right. *Cannabis* pollen curve (blue) and RNA from retting bacteria (bRNA) (blue) in percentage with respect to the total bacterial flora (Rull et al., 2021, 2022). CDA, Columbian discovery of America; DSN, dismantling of the Spanish navy; 20C, mid-20th century *Cannabis* increase.

In the Middle Ages, *Cannabis* was below 20% of the pollen assemblages, which is considered the threshold for hemp retting (Rull et al., 2022). Therefore, the *Cannabis* pollen deposited in the lake would have been wind-transported from small-scale domestic hemp crops (Rull et al., 2021). The situation changed in the Modern Age, when hemp retting for fiber extraction seems to have been the main activity around the lake. By those times, hemp fiber extraction was accomplished by immersion of plants in lakes and ponds, where autochthonous pectinolytic bacteria naturally degraded the cementing compounds to separate the fiber from the stalk (Akin, 2013). The immersion of male hemp plants in lakes caused the pollen to be released and

incorporated into sediments in larger amounts than expected if the only source of this pollen was wind transport from local and/or regional crops. The dramatic Modern-Age hemp pollen increase in Montcortès has been linked to the general increase in the hemp industry to meet the needs (mainly sails and ropes) of the Spanish navy, between the Columbian discovery of America (1492 CE) and the decadence of the Spanish empire, with the ensuing dismantling of its royal navy (1834 CE) (Rull & Vegas-Vilarrúbia, 2014). In Lake Montcortès, hemp-retting activity during this ~350-y period was demonstrated not only by *Cannabis* pollen values over 20% (Fig. 1) but also by the identification of sedimentary RNA from bacteria actually responsible for the retting process (Rull et al., 2022).

During the first half of the 20th century, hemp retting ceased in Lake Montcortès (Rull et al., 2022), and *Cannabis* pollen dropped to minimum historical values, coinciding with a major depopulation (60% reduction) of the Pallars region, where the lake lies, caused by the crisis of the subsistence economy and the massive emigration to industrial cities (Farràs, 2005). The *Cannabis* pollen suddenly increased again to values close to the retting threshold during the second half of the 20th century (Fig. 1). However, bacterial RNA (bRNA) analyses show that local hemp retting never returned (Rull et al., 2022), and therefore, the source for this pollen rise – called here the 20th century *Cannabis* increase (20C) – should be sought in other activities. Although some potential causes for the 20C event have been suggested (Trapote et al., 2018; Rull et al., 2022), the issue has not yet been addressed in depth. This paper analyzes the trends of the last century and the patterns of pollen deposition in recent years to discuss the potential causes of the 20C in relation to the main developments of the hemp industry in the IP.

2. Material and methods

The raw data used in this study are from Rull et al. (2017, 2021, 2022) and Trapote et al. (2018) and are available at <https://data.mendeley.com/datasets/mr4h3x7x35> (pollen) and the Supplementary Material of Rull et al. (2022) (bRNA). In this analysis, *Cannabis* has been excluded from the pollen sum because the parent plant is not part of the local/regional vegetation (Carreras et al., 2005-2006; Mercadé et al., 2013). Pollen percentages have been recalculated and the diagrams have been redrawn, with a focus on the 20th century (1896-2013) and a 2-year modern-analog study (2014-2015) using sediment traps submerged in the lake. Details on the study site and the methods used are available in former papers. The age-depth model was based on varve counting (Corella et al., 2014, 2016), the palynological methods are described in detail in Trapote et al. (2018) and Rull et al. (2017, 2021), and the methods for bRNA analysis are available at Rull et al. (2022).

3. Results

The pollen trends of roughly the last century (1896-2013) are shown in Fig. 2, where it can be seen that the regional vegetation, consisting mainly of a mosaic of *Pinus* and *Quercus* forests intermingled with grasslands and crops, has experienced little variation. The main vegetation shift occurred in the second part of the 17th century, when a conspicuous expansion of conifer forests coincided with a significant decline in olive groves, paralleled by a meaningful increase in total plant cover, as manifested in pollen accumulation rates (PAR). Regarding *Cannabis*, the main shifts occurred in the 1920s and 1980s. The first was a decline that signified the end of the hemp retting decline initiated in the late 19th century, as manifested in the bRNA curve. This *Cannabis* decline is only evident in percentage units, whereas influx values show that this pollen was virtually absent from the beginning of the century. The second increase is the 20C event, when *Cannabis* increased and attained values over 30% in the 1990s, which is beyond the threshold for hemp retting (Rull et al., 2022). Influx values confirm that this increment was

actually an increase in the amount of *Cannabis* pollen reaching the lake, rather than a percentage artifact. The 20C persists today, as indicated by a study of modern pollen sedimentation using sediment traps submerged in the lake (Rull et al., 2017). Indeed, the average amount of *Cannabis* pollen settled during 2014 and 2015 was slightly above 11%, with maxima over the retting threshold in winter and fall (Fig. 3). However, retting practices are no longer developed in the lake.

Pollen (Core MON-0713-G05)




Figure 2. Percentage pollen diagram corresponding to the 20th century at subdecadal (~5-year) resolution. Influx values (grains/particles cm⁻² y⁻¹) for *Cannabis* and pollen accumulation rates (PAR) are also provided. RNA from retting bacteria (bRNA) is represented in percentage with respect to the total bacterial flora (Rull et al., 2022). Solid lines are x10 exaggeration. Red arrows represent the major shifts in *Cannabis* pollen percentage. ΣP, pollen sum.




Figure 3. Seasonal distribution of sedimentary *Cannabis* pollen in Lake Montcortès during 2014 and 2015 (Rull et al., 2017). The hemp retting threshold (20%) is indicated by a red dashed line. Wt, winter; Sp, spring; Sm, summer; FI, fall.

4. Discussion

As mentioned above, wild or cultivated *Cannabis* plants have not been reported in Montcortès local and regional vegetation surveys (Carreras et al., 2005-2006; Mercadé et al., 2013), and hemp retting has not been practiced in the lake since the late 19th century, as indicated by historical records and supported by the absence of bRNA (Rull et al., 2022). Therefore, a reasonable explanation for the 20C event seems to be long-distance pollen dispersal mediated

167 by wind (anemophily). However, the possibility of local or regional illegal crops cannot be
168 disregarded (Munuera et al., 2002). In both cases, a conspicuous increase in these activities
169 would be needed to explain the 20C. Therefore, pollen dispersal patterns and historical trends
170 of *Cannabis* cultivation are critical points to be addressed.

171

172 The high productivity and dispersal power of anemophilous *Cannabis* pollen are well known
173 (e.g., Small & Antle, 2003; Li et al., 2017). On the IP, several studies have documented
174 relatively high amounts of airborne *Cannabis* pollen in areas far from known plant sources. For
175 example, a northern African origin for the airborne *Cannabis* pollen found in some localities
176 from the southeastern IP has been proposed by several authors after detailed aerobiological
177 studies including the use of long-distance dispersal modeling tools. These localities include the
178 cities of Murcia (Munuera et al., 2002; Aznar et al., 2022), Málaga (Cabezudo et al., 1997;
179 Aboulaich et al., 2013) and Córdoba (Cariñanos et al., 2004) (Fig. 1), which are situated 150-
180 300 km north of the African coasts. Pollen from northern Africa comes primarily from illegal
181 crops (Aboulaich et al., 2013). The possibility of nearest sources, such as illegal local/regional
182 crops, was suggested by Munuera et al. (2002). Recently, Aznar et al. (2022) found a significant
183 increase in airborne *Cannabis* pollen between 2017 and 2020 in the region of Murcia (Fig. 1),
184 which coincided with the predominance of local/regional wind patterns, and attributed this
185 increase to the spread of illegal crops in the region rather than to long-distance dispersal from
186 Africa. The occurrence of high *Cannabis* pollen percentages in modern sediments near crop
187 fields of this plant has recently been documented (Boutahar et al., 2023). A typical feature of
188 these crops is their high mobility and spatial turnover, which is a strategy to avoid detection
189 and makes their identification difficult (Aboulaich et al., 2013). These studies come from the
190 southernmost part of the IP, and no comparable information exists for other regions.
191 However, it is expected that the dispersal of *Cannabis* pollen follows similar spatial patterns.
192 Similar surveys should be conducted in the Montcortès region to evaluate the potential
193 contribution of long-distance vs. local/regional pollen sources. These studies could also
194 contribute to the detection of illegal cannabis crops in the area.

195

196 Regarding production, hemp cultivation almost disappeared from Spain in the early 1970s, but
197 a renewed interest in this crop began in 1972. Until the 1960s, hemp was cultivated mainly on
198 irrigated semiarid areas from the eastern IP (Alicante) to provide thread, ropes and fabric (Fig.
199 1). In 1972, however, the paper pulp industry started using hemp as a raw material instead of
200 worn textiles, and in the early 1980s, the main production center shifted from the east to the
201 northeast (Catalonia) in lowland areas (400-900 m elevation) of higher precipitation (600-700
202 mm per year), thus avoiding the need for continuous irrigation (Gorchs & Lloveras, 2003;
203 Gorchs et al., 2017). This geographical shift coincided with a considerable production increase
204 (<1000 tons in 1980 to >8000 tons in 1998) favored by the onset of EU subsidies to hemp
205 cultivation, which started in 1985 (40 € ha⁻¹) and significantly increased (~780 € ha⁻¹) in 1996-
206 97 (Fig. 4). This production increase was centered on the southern Pyrenees, where Lake
207 Montcortès (slightly above 1000 m elevation) lies, under wet climates, as a spring crop in
208 rotation with winter wheat. This rotation significantly increased wheat yields, which has been
209 attributed to the improvement of soil structure and the contribution of hemp to control pests,
210 diseases and particularly weeds, as its rapid early growth outcompetes most weed species,
211 thus reducing the weed burden for the ensuing winter crops (Gorchs et al., 2017).

212




Figure 4. Hemp production in NE Spain (HP) and European Union subsidies to hemp cultivation (EU) (raw data from Gorchs & Lloberas, 2003) compared with the *Cannabis* pollen percentage curve from Lake Montcortès sediments (CP) (Fig. 2).

The 20C event, as recorded in Lake Montcortès was coeval with this 1980s hemp production acceleration, which strongly suggests a causal relationship (Fig. 4). Due to its high dispersion power, hemp pollen could have easily reached Lake Montcortès from adjacent regional pre-Pyrenean hemp crops, which could be confirmed by modeling the *Cannabis* pollen dispersal around the lake area. Hemp cultivation declined again shortly after the 1990s and was abandoned by 2006, due to the relocation of the hemp straw processing industry (Gorchs et al., 2017). This decline was also paralleled by pollen trends (Fig. 2), thus reinforcing the potential causal relationship. However, *Cannabis* pollen remained close to the retting threshold until at least 2015 (Figs. 2 and 3), which deserves an additional explanation. Additional hemp pollen sources from illegal crops cannot be disregarded, although the available evidence is less robust. In recent decades, a relevant rise in large-scale *Cannabis* plantations has been recorded in the IP, especially along the Mediterranean coasts, in a trend to increasingly replace hashish imports from northern Africa (Alvarez et al., 2016; Belackova et al., 2016). The potential manifestation of this peninsula-wide rising trend in the Montcortès region remains to be documented, and is worth to be addressed in future studies.

The geographical extent of the 20C event remains unknown, as it has not been recorded in other IP localities. This lack of evidence could suggest that this event could be a local/regional phenomenon, but methodological constraints should not be dismissed. On the one hand, only 23 of the 56 records included in the peninsular-wide CHIP database (Rull et al., 2023) contain the 20th century, and of them, *Cannabis* pollen occurs in only four (Fig. 1), excluding Lake Montcortès. Of these four sites, *Cannabis* pollen is abundant (maximum of 25%) in only one (ET), whereas in the others this pollen has maxima of only 5% (CE) or shows single scattered occurrences (AD, EN). On the other hand, the resolution of these records, usually centennial to millennial, is not enough to resolve eventual changes that occurred within the 20th century. For example, in ET, which is the most similar to Montcortès regarding pollen abundance, the record ends in 1991, and only a couple of samples are available for the whole 20th century (Riera et al., 2004). Therefore, the lack of high-resolution records including the 20th century is a major handicap to assessing the spatial patterns of the 20C across the IP.

Another interesting observation is that the major vegetation shift documented above and the 20C occurred at or shortly after the onset of the “Anthropocene”, as currently defined by the Anthropocene Working Group (AWG) (Waters et al., 2023). Indeed, the AWG proposes that this boundary should be situated in the mid-20th century, coinciding with the Great Acceleration, when the Earth System underwent a global transformation overcoming the range

of variability of the Holocene (Steffen et al., 2015). Our results, together with the very high resolution (seasonal) of the Montcortès record, suggest that the sediments of this lake could be a suitable candidate to identify and precisely date an eventual “Anthropocene” GSSP (Global Stratotype Section and Point) (Waters et al., 2018). The proxies identified by the AWG as the best suited stratigraphic markers (radioisotopes, fly ash, stable carbon and nitrogen isotopes) have not yet been analyzed in the Montcortès record, but efforts in this direction are worth doing, especially after the detection of significant Hg and Pb increases during the Industrial Revolution and the Great Acceleration in these sediments (Corella et al., 2017).

5. Conclusions and future prospects

A distinct and characteristic increase in *Cannabis* pollen has been recorded in the unique varved record of Lake Montcortès during the 1980s, which has been related to the significant increase in hemp production documented by the same time in the southern Pyrenees, fostered by the incoming EU subsidies to this crop. Additional sources for this pollen type, notably illegal cannabis plantations, cannot be disregarded but sound evidence is still lacking. The geographical extent of this palynological event, called here 20C (20th century *Cannabis* increase), remains unknown due to the lack of similar continuous and absolutely dated high-resolution records for the 20th century. A number of future studies are proposed to complement this preliminary assessment, including (i) increasing the resolution of the available pollen record from subdecadal (~5 years) to annual, (ii) modeling the *Cannabis* pollen dispersion in the region around Montcortès, (iii) surveying the occurrence of illegal plantations as potential local sources for *Cannabis* pollen, (iv) increasing the number and geographical extent of 20th century high-resolution records similar to Montcortès, and (v) evaluating the Lake Montcortès record as a potential candidate for the identification of the current AWG “Anthropocene” prospect.

Acknowledgments

This work has been funded by the Ministry of Economy and Competitiveness, projects CGL 2012-3665 and CGL 2017-85682-R.

References

- Aboulaich, N., Trigo, M.M., Bouziane, H., Cabezudo, B., Recio, M., El Kadiri, M., et al. 2013. Variations and origin of the atmospheric pollen of *Cannabis* detected in the province of tetouan (NW Morocco): 2008-2010. *Sci. Total Environ.* 443, 413-419.
- Akin, D.E. 2013. Linen most useful: perspective on structure, chemistry and anzyme for retting flax. *ISRN Biotechnol.* 2013, 1-23.
- Alvarez, A., Gamella, J.F., Parra, I. 2016. *Cannabis* cultivation in Spain: a profile of plantations, growers and production systems. *Int. J. Drug Policy* 37, 70-81.
- Aznar, F., Negral, L., Moreno-Grau, S., Elvira-Rendueles, B., Costa-Gómez, I., Moreno, J.M. 2022. *Cannabis*, an emerging aeroallergen in southeastern Spain (Region of Murcia). *Sci. Total Environ.* 833, 155156.
- Balant, M., Gras, A., Gálvez, F., Garnatje, T., Vallès, J., Vitales, D. 2021. CANNUSE, a database of traditional *Cannabis* uses – an opportunity for new research. *Database* 2021, baab024.
- Barcaccia, G., Palumbo, F., Scariolo, F., Vannozzi, A., Bona, S. 2020. Potentials and challenges of genomics for breeding cannabis cultivars. *Front. Plant Sci.* 11, 573299.
- Belackova, V., Tomkova, A., Zabransky, T. 2016. Qualitative research in Spanish cannabis social clubs: “The moment you enter the door, you are minimising te risks”. *Int. J. Drug Policy* 34, 49-57.

305 Boutahar, A., Cariñanos, P., Picone, R.M., Crisafulli, A., Molero, J., Zahrae, F., et al. 2023.
 306 Modern pollen-vegetation relationships in the Rif mountains (Northern Morocco). *Rev.*
 307 *Palaeobot. Palynol.* 310, 104828.

308 Cabezudo, B., Recio, M., Sánchez-Laulhé, J.M., Trigo, M.M., Toro, F.J., Polvorinos, F. 1997.
 309 Atmospheric transportation of marihuana pollen from north Africa to the southwest of
 310 Europe. *Atmos. Environ.* 31, 3323-3328.

311 Cariñanos, P., Galán, C., Alcázar, P., Domínguez, E. 2004. Analysis of the particles transported
 312 with dust-clouds reaching Cordoba, Sotheastern Spain. *Arch. Environ. Contam. Toxicol.* 46,
 313 141-146.

314 Carreras, J., Vigo, J., Ferré, A. 2005-2006. Manual dels Hàbitats de Catalunya, vols. I-VIII. Dep.
 315 Medi Ambient Habitatge, Generalitat de Catalunya, Barcelona.

316 Carrión, J.S., Munuera, M., Ochando, J., López-Sáez, J.A., Casas-Gallego, M., González-
 317 Sampériz, P., et al. 2022. Paleoflora y Paleovegetación Ibérica III: Holoceno. Min. Ciencia
 318 Innovación & Fund. Séneca, Murcia.

319 Clarke, R.C., Merlin, M.D. 2013. *Cannabis*: Evolution and Ethnobotany. Univ. California Press,
 320 Los Angeles.

321 Clarke, R.C., Merlin, M.D. 2016. *Cannabis* domestication, breeding history, present-day genetic
 322 diversity, and future prospects. *Crit. Rev. Plant Sci.* 35, 293-327.

323 Corella, J.P., Benito, G., Rodríguez-Lloveras, X., Brauer, A., Valero-Garcés, B. 2014. Annually-
 324 resolved lake records of extreme hydro-meteorological events since AD 1347 in NE Iberian
 325 Peninsula. *Quat. Sci. Rev.* 93, 77-90.

326 Corella, J.P., Valero-Garcés, B., Vicente-Serrano, S.M., Brauer, A., Benito, G. 2016. Three
 327 millennia of heavy rainfalls in Western Mediterranean: frequency, seasonality and
 328 atmospheric drivers. *Sci. Rep.* 6, 38206.

329 Corella, J.P., Valero-Garcés, B., Wang, F., Martínez-Cortizas, A., Cuevas, C.A., Saiz-Lopez, A.
 330 2017. 700 years reconstruction of mercury and lead atmospheric deposition in the
 331 Pyrenees (NE Spain). *Atmos. Environ.* 155, 97-107.

332 Duvall, C. 2014. *Cannabis*. Reaktion, London.

333 Farràs, F. 2005. El Pallars contemporani. In: Marugan, C.M., Rapalino, V. (eds.), *Història del*
 334 *Pallars. Dels Orígens als Nostres Dies*. Pagés, Lleida, pp. 121-144.

335 Gorchs, G., Lloveras, J. 2003. Current status of hemp production and transformation in Spain.
 336 *J. Ind. Hemp.* 8, 45-64.

337 Gorchs, G., Loveras, J., Serrano, L., Cela, S. 2017. Hemp yields and its rotation effects on wheat
 338 under rainfed Mediterranean climates. *Agron. J.* 109, 1551-1560.

339 Gray, D.J., Clarke, R.C., Trigiano, R.N. 2016. Introduction to the special issue on *Cannabis*. *Crit.*
 340 *Rev. Plant Sci.* 35, 425-426.

341 Kovalchuk, I., Pellino, M., Rigault, P., van Velzen, R., Ebersbach, J., Ashnest, J.R., et al. 2020. The
 342 genomics of *Cannabis* and its close relatives. *Ann. Rev. Plant Biol.* 71, 713-739.

343 Li, F., Gaillard, M.-J., Sugita, S., Mazier, F., Xu, Q., Zhou, Z., et al. 2017. Relative pollen
 344 productivity estimates for major plant taxa of cultural landscapes in central eastern China.
 345 *Veget. Hist. Archaeobot.* 26, 587-605.

346 Mercadé, A., Vigo, J., Rull, V., Vegas-Vilarrúbia, V., Garcés, S., Lara, A., et al. 2013. Vegetation
 347 and landscape around Lake Montcortès (Catalan pre-Pyrenees) as a tool for
 348 palaeoecological studies of lake sediments. *Collect. Bot.* 32, 87-101.

349 Munuera, M., Carrión, J.S., Navarro, C. 2002. Seasonal fluctuations of the airborne pollen
 350 spectrum in Murcia (SE Spain). *Aerobiologia* 18, 141-151.

351 Riera, S., Wansard, G., Julià, R. 2004. 2000-year environmental history of a karstic lake in the
 352 Mediterranean pre-Pyrenees: the Estanya lakes (Spain). *Catena* 55, 293-324.

353 Rull, V. 2022. Origin, early expansion, domestication and anthropogenic diffusion of *Cannabis*,
 354 with emphasis on Europe and the Iberian Peninsula. *Persp. Plant Ecol. Evol. Syst.* 55,
 355 125670.

356 Rull, V., Vegas-Vilarrúbia, T. 2014. Preliminary report on a mid-19th century *Cannabis* pollen
357 peak in NE Spain: historical context and potential chronological significance. *Holocene* 24,
358 1378-1383.

359 Rull, V., Trapote, M.C., Safont, E., Cañellas-Boltà, N., Pérez-zanón, N., Sigró, J., Buchaca, T., et
360 al. 2017. Seasonal patterns of pollen sedimentation in Lake Montcortès (Central Pyrenees)
361 and potential applications to high-resolution paleoecology: a 2-year pilot study. *J.*
362 *Paleolimnol.* 57, 95-108.

363 Rull, V., Vegas-Vilarrúbia, T., Corella, J.P., Trapote, M.C., Montoya, E., Valero-Garcés, B. 2021. A
364 unique Pyrenean varved record provides a detailed reconstruction of Mediterranean
365 vegetation and land-use dynamics over the last three millennia. *Quat. Sci. Rev.* 268,
366 107128.

367 Rull, V., Sacristán-Soriano, O., Sánchez-Melsió, A., Borrego, C.M., Vegas-Vilarrúbia, T. 2022.
368 Bacterial phylogenetic markers in lake sediments provide direct evidence for historical
369 hemp retting. *Quat. Sci. Rev.* 295, 107803.

370 Rull, V., Burjachs, F., Carrión, J.S., Ejarque, A., Fernández, S., López-Sáez, J.A., et al. 2023.
371 Historical biogeography of *Cannabis* in the Iberian Peninsula: a probabilistic approach
372 using palynological evidence. *Persp. Plant. Ecol. Evol. Syst.* 58, 125704.

373 Small, E., Antle, T. 2003. A preliminary study of pollen dispersal in *Cannabis sativa* in relation to
374 wind direction. *J. Ind. Hemp* 8, 37-50.

375 Steffen, W., Broadgate, W., Deutsch, L., Gaffney, O., Ludwig, C. 2015. The trajectory of the
376 Anthropocene: the Great Acceleration. *Anthr. Rev.* 2, 81-98.

377 Trapote, M.C., Rull, V., Giralt, S., Corella, J.P., Montoya, E., Vegas-Vilarrúbia, T. 2018. High-
378 resolution (sub-decadal) pollen analysis of varved sediments from Lake Montcortès
379 (southern Pyrenean flank): a fine-tuned record of landscape dynamics and human impact
380 during the last 500 years. *Rev. Palaeobot. Palynol.* 259, 207-222.

381 Warf, B. 2014. High points: an historical geography of *Cannabis*. *Geogr. Rev.* 104, 414-438.

382 Waters, C.N., Zalasiewicz, J., Summerhayes, C., Fairchild, I.J., Rose, N.L., Loader, N.J., et al.
383 2018. Global Boundary Stratotype Section and Point (GSSP) for the Anthropocene Series:
384 where and how to look for potential candidates. *Earth-Sci. Rev.* 178, 379-429.

385 Waters, C.N., Turner, S.D., Zalasiewicz, J., Head, M.J. 2023. Candidate sites and other reference
386 sections for the Global boundary Stratotype Section and Point of the Anthropocene series.
387 *Anthr. Rev.* doi 10.1177/20530196221136422.