

Screening Heroin Smokers Attending Community Drug Clinics for Change in Lung Function

A Cohort Study

*Rebecca Nightingale, MSc; Kevin Mortimer, MD, PhD; Emanuele Giorgi, PhD;
Paul P. Walker, MD; Marie Stolbrink, MD; Tara Byrne, BSc; Kerry Marwood, NVQ;
Sally Morrison-Griffiths, MD; Susan Renwick, MPH; Jamie Rylance, MD, PhD; and
Hassan Burhan, MD*

CHEST 2020; 157(3):558-565

Online supplements are not copyedited prior to posting and the author(s) take full responsibility for the accuracy of all data.

© 2019 AMERICAN COLLEGE OF CHEST PHYSICIANS. Reproduction of this article is prohibited without written permission from the American College of Chest Physicians. See online for more details. DOI: 10.1016/j.chest.2019.11.006

e-Table 1: Characteristics of participants completing follow up questionnaires, with and without ATS standard spirometry

Variable	Acceptable spirometry N (%) Mean (SD)	Unacceptable spirometry or declined/excluded for medical reasons N (%) Mean (SD)	Chi ² or Ttest p-value
Sex	N=106	N=55	
Female	30.2 (32)	25.5 (14)	0.53
Male	69.8 (74)	74.5 (41)	
Age in years, mean (SD)	50.9 (5.2)	51.3 (5.6)	0.67
IMD Score	50.3 (12.9)	53.9 (12.3)	0.09
Occupation			0.65
Unemployed	85.9 (91)	83.6 (46)	
Employed	14.1 (15)	16.2 (9)	
Housing			0.76
Own home (included rental),n	77.4 (82)	76.4 (42)	
Homeless, n (%)	4.72 (5)	1.82 (1)	
Other, n (%)	17.9 (19)	21.78 (12)	
Cigarette Smoking Status			0.125
Current , n (%)	78.3 (83)	50 (90.9)	
Ex, n (%)	22 (20.8)	5 (9.1)	
Never, n (%)	1 (0.9)	0 (0)	
Heroin Smoking Status			0.312
Current, n (%)	44.34 (47)	52.7 (29)	
Ex, n(%)	55.7 (59)	47.3 (26)	
Crack smoking			0.74
Current, n (%)	18.9 (20)	23.6 (13)	
Ex, n(%)	55.6 (59)	54.6 (30)	
Never, n (%)	25.5 (27)	21.8 (12)	