

SUPPLEMENTARY TABLE S6. DIFFERENCES IN WITHDRAWAL SYMPTOMS AND BELIEFS ABOUT ADDICTIVENESS OF CANNABIS AMONG YOUNG, MIDDLE-AGE, AND OLDER CANNABIS USERS

	<i>Young</i> (n = 1300), %	<i>Middle age</i> (n = 1048), %	<i>Older</i> (n = 507) %	
Withdrawal symptoms				
Not applicable	30.2 ^a	36.3 ^a	47.1 ^b	$\chi^2 = 28.38, p < 0.001$
Irritability	35.2 ^a	37.5 ^a	22.5 ^b	$\chi^2 = 30.02, p < 0.001$
Insomnia/interrupted sleep	33.5 ^a	30.4 ^a	22.7 ^b	$\chi^2 = 16.43, p < 0.001$
Anxiety	22.8 ^{ab}	25.8 ^a	16.4 ^b	$\chi^2 = 19.92, p < 0.001$
Loss of appetite	26.3 ^a	15.8 ^b	5.9 ^c	$\chi^2 = 96.72, p < 0.001$
Vivid dreams	21.9	15.6	9.7	$\chi^2 = 12.09, p = 0.002$
Loss of productivity	12.8	13.2	9.5	$\chi^2 = 7.31, p = 0.03$
Tiredness	10.1	7.3	5.5	$\chi^2 = 3.66, p = 0.16$
Nausea	8.0 ^a	7.0 ^{a,b}	4.7 ^b	$\chi^2 = 13.57, p = 0.001$
Improved productivity	7.2	3.1	2.0	$\chi^2 = 8.17, p = 0.02$
Weight loss	5.7 ^a	2.8 ^{a,b}	1.0 ^b	$\chi^2 = 23.04, p < 0.001$
Sweating	5.1	3.4	1.8	$\chi^2 = 10.68, p = 0.005$
Tremor	1.5	1.1	1.6	$\chi^2 = 1.58, p = 0.45$
Salivation	0.9	0.3	0.2	$\chi^2 = 4.88, p = 0.09$
Addictiveness				
Trouble stopping	20.3	15.2	10.4	$\chi^2 = 9.52, p = 0.009$
Cannabis is addictive	20.0 ^a	17.5 ^a	8.3 ^b	$\chi^2 = 25.62, p < 0.001$
Cannabis is not addictive	65.8	67.2	76.4	$\chi^2 = 11.87, p = 0.003$
Don't know if it's addictive	14.2	15.2	15.3	$\chi^2 = 0.75, p = 0.69$

Percentages represent overall raw percentages (without effects of covariates removed). Bolded chi-square results indicate an overall significant difference ($p \leq 0.001$) across the three groups with the effects of the covariates (bolded in Supplementary Table S2) statistically removed. Different superscripts represent specific group differences with effects of covariates removed and $p \leq 0.001$.