

Table S2 MIC

MIC	WT	<i>ΔhexSDF</i>	<i>ΔugtB</i>	<i>ΔugtA</i>
Ampicillin (μg/ml)	1.33 ± 0.58	0.8 ± 0	1 ± 0	0.75 ± 0.43
Bacitracin (μg/ml)	256 ± 0 ^	64 ± 0 ^	256 ± 0 ^	128 ± 0 ^
Cefoxitin (μg/ml)	256 ± 0 ^	ND	ND	256 ± 0 ^
Daptomycin (μg/ml)	4 ± 0	0.833 ± 0.29 ****	4 ± 0	4 ± 0
Fosfomycin (μg/ml)	16 ± 0 ^	ND	16 ± 0 ^	32 ± 0 ^
Meropenem (μg/ml)	3.33 ± 1.15	ND	ND	3.33 ± 1.15
Nisin (μg/ml)	180 ± 0	120 ± 51.96	90 ± 0 **	90 ± 0 **
Vancomycin (μg/ml)	1 ± 0 ^	ND	0.5 ± 0 ^	1 ± 0 ^
Lauric Acid (mM)	0.25 ± 0 ^	0.25 ± 0 ^	0.25 ± 0 ^	0.25 ± 0 ^
Linoleic Acid (mM)	>32	>32	>32	>32
Lysozyme (mg/ml)	8 ± 0	6.67 ± 2.31	4 ± 0 *	3.33 ± 1.15 **
Novobiocin (μg/ml)	16 ± 0	16 ± 0	16 ± 0	10.67 ± 4.62

ND-not determined

Mean ± standard deviation

* $p < 0.05$

** $p < 0.01$

**** $p < 0.0001$

^Did not calculate statistical significance using one-way analysis of variance as the standard deviation was 0.