

Supporting information, Table S1

Oligonucleotides used for Real-Time PCR reactions

Primer	5'3' nucleotide sequence
IGF-I P1 for	GCGATGGGGAAAATCAGCAG
IGF-I P1 rev	GAAGAGCATCCACCAGCTCA
IGF-I P2 for	AGTACCCACTCTGACCTGCT
IGF-I P2 rev	GAAGAGCATCCACCAGCTCA
CB1R for	TTCCTTGTAGCAGAGAGCCAG
CB1R rev	CTCAACGTGACTGAGAAAGAGG
CB2R for	TGACAAATGACACCCAGTCTTCT
CB2R rev	ACTGCTCAGGATCATGTACTCCTT
NAPE-PLD for	CAGGTTCCAAAGAGGAACTTGAC
NAPE-PLD rev	GATGAGCTCGTCCATTTCCAC
DAGLα for	GATTCCCAGTCAGATGCCTAC
DAGLα rev	GACAGGAAAGCCAAGATGTC
FAAH for	GAGATGTATCGCCAGTCCGT
FAAH rev	ACAGGCAGGCCTATACCCTT
MGL for	TCTTCCTCCTGGGCCACT
MGL rev	AAAGTAGGTTGGCCTCTCTGC
IL-6 for	ACGGCCTTCCCTACTTCACA
IL-6 rev	CATTTCACGATTTCCAGAGA
TNFα for	GCCTCTTCTCATTCTGCTTG
TNFα rev	CTGATGAGAGGGAGGCCATT
βActin for	GTGATGGTGGGAATGGGTCA
βActin rev	CTGGATGGCTACGTACATGG