

Supplementary Materials: A Sustainable Air-Assisted Liquid–Liquid Microextraction Procedure Coupled to Liquid Chromatography–Tandem Mass Spectrometry for the Determination of 11-Nor-9-carboxy- Δ^9 -tetrahydrocannabinol and Hexahydrocannabinol Metabolites in Urine

Pamela Cabarcos-Fernández, Sara Odoardi, Serena Mestria, Valeria Valentini, Giulia Biosa, Ana María Bermejo-Barrera and Sabina Strano-Rossi

Table S1. Sensitivity study performed during method validation (CV, coefficient of variation).

	Concentration, ng/mL	S/N	Bias (%)	Accuracy (%)	CV (%)	Status
HHC- COOH	10	36.6	-0.33	99.7	18.8	QC level
	5	17.9	6.27	106.2	7.8	Accepted LLOQ
	2.5	5.77	-2.7	97.3	51.3	Below LLOQ
	1	5.11	50.6	49.3	72.8	Below LLOQ
8-OH- 9(R)- HHC	10	29	0.50	100.5	5.77	QC level
	5	14.1	19.4	119.4	9.71	Accepted LLOQ
	2.5	6.1	46.8	146.8	25.0	Below LLOQ
	1	2.48	103.2	203.3	12.8	Below LLOQ
11-OH- 9(R)- HHC	10	16.4	12.3	101.9	13.8	QC level
	5	8.3	12.9	109.0	3.12	Accepted LLOQ
	2.5	4.19	131.7	231.7	11.9	Below LLOQ
	1	1.87	314	414	62.6	Below LLOQ
THC- COOH	10	42.7	1.44	101.4	24.2	QC level
	5	31.4	14.1	120.0	12.8	Accepted LLOQ
	2.5	12.6	73.2	173.2	17.9	Below LLOQ
	1	5.64	242.7	342.7	2.4	Below LLOQ

Note: Concentrations of 1 and 2.5 ng/mL were evaluated during the sensitivity assessment but were not accepted as LLOQ because the predefined criteria for quantitative analysis (accuracy and/or precision) were not consistently fulfilled.