

Phenotype–genotype discordance in antimicrobial resistance profiles of Gram-negative uropathogens recovered from catheter-associated urinary tract infections in Egypt

Mohamed Eladawy^{1, 2}, Nathan Heslop¹, David Negus¹, Jonathan C. Thomas^{1#}, Lesley Hoyles^{1#}

¹ Department of Biosciences, School of Science and Technology, Nottingham Trent University, Nottingham NG11 8NS, UK.

² Department of Microbiology and Immunology, Faculty of Pharmacy, Mansoura University, Egypt.

Mohamed Eladawy ORCID 0000-0003-0200-0752

Nathan Heslop ORCID 0009-0005-4369-2005

David Negus ORCID 0000-0001-9047-4565

Jonathan Thomas ORCID 0000-0002-1599-9123

Lesley Hoyles ORCID 0000-0002-6418-342X

#Corresponding authors: jonathan.thomas@ntu.ac.uk, lesley.hoyles@ntu.ac.uk

Keywords: uropathogenic strains, bacterial genomics, resistance phenotypes, UPEC, EUCAST.

Abbreviations: AMR, antimicrobial resistance; CARD, Comprehensive Antibiotic Resistance Database; CAUTI, catheter-associated urinary tract infection; DDT, disc diffusion test; ESBL, extended spectrum β -lactamase; HAI, healthcare-associated infection; ICU, intensive care unit; MDR, multidrug-resistant; ME, major error; MLST, multilocus sequence type; VME, very major error; UPEC, uropathogenic *Escherichia coli*; UTI, urinary tract infection; XDR, extensively drug-resistant.

Data availability: The sequence data included in the study are available under BioProject PRJNA1208035.

ABSTRACT

Catheter-associated urinary tract infections (CAUTIs) are among the most common healthcare-associated infections in low- and middle-income countries (LMICs). In this study, phenotypic (EUCAST AMR profiles) and genomic data were generated for 132 isolates (67 *Escherichia coli*, 14 *Pseudomonas aeruginosa*, 11 *Klebsiella pneumoniae*, 9 *Proteus mirabilis*, 8 *Providencia* spp., 5 *Enterobacter hormaechei*, and 18 rare uropathogens) recovered from CAUTIs in an Egyptian hospital. Among the *Escherichia coli* isolates, phylogroup B2 was most represented (53.7 %), followed by B1 (19.4 %), A (11.9 %), D (7.4 %), F (5.9 %) and C (1.4 %). Several (22/132, 16.6 %) isolates were multidrug-resistant, while almost half (62/132, 46.9 %) were extensively drug-resistant. Comparison of phenotypic data with genotypic data from three different AMR-profiling tools (ResFinder, CARD, AMRFinder) highlighted phenotype–genotype discordance as an important consideration in resistome studies in Egypt, with a total concordance of 91.1 %, 85.7 %, and 80.3 % for ResFinder, CARD and AMRFinder, respectively. *Pseudomonas*, at the species level, exhibited the greatest discordance. At the antimicrobial level, meropenem was subject to greatest discordance. In addition to the findings from our comparative analyses, new AMR variants are reported for Egypt for *Pseudomonas* (*OXA-486*, *OXA-488*, *OXA-905*, *IMP-43*, *PDC-35*, *PDC-45*, *PDC-201*) and *Escherichia coli* (*TEM-176*, *TEM-190*). In summary, we show that there is phenotype–genotype discordance in AMR profiling among CAUTI isolates, highlighting the need for comprehensive approaches in resistome studies. We also show the genomic diversity of Gram-negative uropathogens contributing to disease burden in a little-studied LMIC setting.

INTRODUCTION

Urinary tract infections (UTIs) are an increasing public health problem caused by a range of uropathogens. The frequency of healthcare-associated UTIs is 12.9 %, 19.6 % and 24 %, respectively, in the United States, Europe and developing countries ¹. In 2019, bacterial antimicrobial resistance (AMR) contributed to 48,700 deaths associated with UTIs out of 541,000 cases of infectious syndromes ². Resistance to almost all antibiotics in healthcare-associated UTIs is above 20 %, with significant geographical variation ¹. However, meta-analysis and surveillance data are scarce from low- and middle-income countries (LMICs) ^{3, 4}. Egypt initiated a national surveillance program for healthcare-associated infections (HAIs) with international collaborators and national entities to establish benchmarks to outline the microbiological profiles and reduce AMR in clinical settings. Phase one of the National Action Plan showed nearly 50 % of *Escherichia (Esc.) coli* and 75 % of *Klebsiella pneumoniae* isolates from urine were resistant to quinolones and cephalosporins, respectively ⁵. Across 27 countries, Egypt came in third place for total consumption of antibacterials (1355.4 tonnes) in 2020 ⁶.

In Egypt, catheter-associated UTIs (CAUTIs) are among the most common types of HAIs affecting patients in intensive care units (ICUs) ⁷⁻¹⁰. Between 2011 and 2012, 15 % (69/472) of HAIs were UTIs and the rate of CAUTIs was 1.2 per 1,000 device-days ⁷. Another surveillance showed that UTIs represented 15 % (406/2688) of HAIs, the rate of CAUTI development increased up to 1.9/1,000 device-days and CAUTIs represented 98.3 % of UTIs ⁸. An epidemiological study (between 2011 and 2017) of carbapenem-resistant *Enterobacteriaceae* in Egyptian ICUs showed that UTIs accounted for 14.8 % (236/1598) of HAIs, with 63.8 % of UTIs catheter-related ¹⁰. The second most common HAIs were UTIs (12.2 %) in Egyptian clinical settings ⁹.

There are few reports on the genomic diversity of uropathogens found in Egyptian clinical settings ¹¹⁻¹⁶. However, high rates of AMR are observed in Egypt beside documented transmission of carbapenem-resistant strains from Egypt to other countries ¹⁷⁻¹⁹. We collected a range of Gram-negative uropathogens (n = 132 isolates) from an Egyptian hospital over a 9-month period, and generated genotypic and phenotypic data for them. Our aim was to investigate the AMR patterns associated with UTIs in an Egyptian clinical setting and provide evidence that could inform the development of effective antimicrobial surveillance plans and targeted infection

control policies. In addition, we assessed the agreement/concordance between phenotypic and genotypic data to evaluate the usefulness of such data from a diagnostic perspective.

METHODS

Recovery of isolates

Anonymized isolates (n = 132) with no patient data were recovered from urinary catheters between December 2020 and August 2021 by staff at the Urology and Nephrology Center, Mansoura, Egypt during routine diagnostic procedures as described previously¹⁶. The study of anonymized clinical isolates beyond the diagnostic requirement was approved by the Urology and Nephrology Center, Mansoura, Egypt. No other ethical approval was required for the use of the clinical isolates.

Antimicrobial susceptibility testing

Antimicrobial susceptibility testing was performed using the disc diffusion test (DDT) on Mueller-Hinton agar (Oxoid Ltd, UK)¹⁶. Twelve different antimicrobial discs (Oxoid Ltd, UK) were used: penicillins – piperacillin/tazobactam (30/6 µg); cephalosporins – ceftazidime (10 µg), cefepime (30 µg); monobactams – aztreonam (30 µg); carbapenems – doripenem (10 µg), meropenem (10 µg); aminoglycosides – tobramycin (10 µg), amikacin (30 µg); quinolones – ciprofloxacin (5 µg), levofloxacin (5 µg); miscellaneous – nitrofurantoin (100 µg), sulfamethoxazole/trimethoprim (1.25–23.75 µg).

Results were interpreted according to breakpoints of EUCAST guidelines version 12.0, 2022 (http://www.eucast.org/clinical_breakpoints/), with EUCAST reference strains - *Esc. coli* ATCC 25922 and *Pse. aeruginosa* ATCC 27853 - used for quality control purposes.

DNA extraction and whole-genome sequencing

DNA was extracted from isolates and their genomes were sequenced and assembled as described previously^{16, 20}.

Bioinformatic analyses

Bakta v1.5.1 (database v4.0) was used for annotating genes within genomes²¹. sourmash v4.8.4²² was used to compare and identify (with Genome Taxonomy Database release 214) species for all genomic data (**Supplementary Table 1**). Genomes that could not be assigned to a species by sourmash were identified using ribosomal multilocus sequence type (rMLST) (<https://pubmlst.org/species-id>; ²³), and by comparison with relevant reference sequences (**Supplementary Table 1**) using FastANI v1.33²⁴.

Serotypes were determined using ECTyper v1.0.0 (database v1.0)²⁵. Phylogrouping of *Esc. coli* was achieved using Ezclermont v0.7.0²⁶. MLST of each isolate was determined using *mlst* v2.23.0 (<https://github.com/tseemann/mlst>) and relevant schema from <http://pubmlst.org>²⁷⁻³³.

AMR genes were identified using (accessed February 2025) the Resistance Gene Identifier v6.0.3 of the Comprehensive Antibiotic Resistance Database (CARD) v4.0.0³⁴, ResFinder v4.6.0 (ResFinder database v2.4.0, PointFinder database v4.1.1)³⁵, and AMRFinder v4.0.3 (database version 2024-12-18.1)³⁶. Genes responsible for efflux pumps were excluded from this study³⁷. Only AMR genes that showed perfect or strict matches (identity ≥ 90 %) were reported.

For the *Esc. coli* genomic data, a phylogenetic tree was constructed using core genes identified using Panaroo v1.3.4³⁸ (strict mode, identity threshold 98 %, alignment option “core”). Core genes were present in all 67 *Esc. coli*, and were aligned with prank (v170427)³⁹. Gene alignments were concatenated with FASconCAT-G (v1.02) into a single alignment⁴⁰. Phylogenetic analysis was performed using the concatenated sequences and RaxML v1.2.0 (GTR+G8+IO)⁴¹. Node support was determined using 1000 non-parametric bootstrap replicates. The phylogenetic tree was visualized using iTOL v6.9⁴². Virulence traits encoded in genomes were detected via BLASTP (≥ 90 % coverage, ≥ 70 % identity) with the core protein sequence dataset from the Virulence Factors Database (VFDB)⁴³.

Comparison of AMR genotypes and phenotypes

According to EUCAST guidelines, “susceptible, standard dosing regimen” (S) and “susceptible, increased exposure” (I) categories are grouped under the term “susceptible”. Using CARD, ResFinder, and AMRFinder, genomic data of isolates were compared with DDT results for 132 isolates against 12 antimicrobials [except when certain species lacked breakpoints; i.e. nitrofurantoin was used only for *Esc. coli* (n = 67), sulfamethoxazole/trimethoprim was used only for *Enterobacterales* (n = 114)]. For each combination, concordance was considered positive if a) WGS data were predicted to encode AMR genes and the isolate had a phenotypic resistant profile “true resistant” (WGS-R/DDT-R) or b) WGS data were not predicted to encode AMR genes and the isolate had a phenotypic susceptible profile “true susceptible” (WGS-S/DDT-S)^{44, 45}. Discordance was considered positive where major errors (MEs) or very major errors (VMEs) were detected. MEs (i.e. false positives; WGS-R/DDT-S) are defined as a resistant genotype and susceptible phenotype. VMEs (i.e. false negatives; WGS-S/DDT-R) are defined as a susceptible genotype and resistant phenotype. To detect AMR gene mutations, PointFinder tool for ResFinder and Point mutations tool for AMRFinder results were also included in concordance analyses.

RESULTS AND DISCUSSION

The main aim of this study was to characterize a collection of Gram-negative CAUTI-associated bacteria (n = 132 isolates) isolated in Egypt to investigate AMR genotype–phenotype concordance in relation to gene data available in widely used, public databases. We also sought to understand the genomic diversity of the dominant species contributing to CAUTIs in an Egyptian clinical setting.

Genome characterization

Summary statistics for the 132 high-quality genomes assembled in this study can be found in **Supplementary Table 1**. All genomes were assigned to the class *Gammaproteobacteria*, with the exception of Me47, which was found to represent a novel member of the family *Alcaligenaceae* (**Figure (1A)**). Among our isolates, UPEC dominated

(50.7 %, 67/132), followed by *Pseudomonas aeruginosa* (10.6 %, 14/132), *K. pneumoniae* (8.3 %, 11/132), other uropathogens (6.8 %, 9/132), *Proteus mirabilis* (6.8 %, n = 9/132), *Providencia* spp. (6 %, n = 8/132), *Enterobacter hormaechei* (3.7 %, n = 5/132), *Alcaligenes* spp. (3 %, n = 4/132), *Serratia* spp. (2.2 %, n = 3/132) and *Pseudomonas vlassakiae* (1.51 %, n = 2/132). UPEC was previously reported as the main cause of UTIs for adults and neonates in Egypt, followed by other *Enterobacterales* or Gram-positive bacteria ⁴⁶⁻⁴⁹. *Providencia* – once considered a rare pathogen – is increasingly recognized as an opportunistic pathogen capable of causing serious UTIs ⁵⁰; we found it among the main taxa represented in our study.

Antimicrobial susceptibility of isolates

According to Egyptian Urological guidelines, symptomatic CAUTIs are treated like complicated UTIs with second- or third-generation cephalosporins, while uncomplicated pyelonephritis is treated with a short course of fluoroquinolones as the first-line therapy ⁵¹. AMR for quinolones (ciprofloxacin, levofloxacin) and cephalosporins (cefepime, ceftazidime) was >64 % among MDR uropathogens in Egyptian clinical settings ⁵². More than half of UPEC isolates (57.8 %) showed high-level ciprofloxacin resistance and *gyrA* mutations were detected in 76.7 % of isolates in a previous study, leading to recommendations for revision of empirical antibiotic treatment of UTIs ⁵³.

Many of our isolates were resistant to ceftazidime (68.9 %, 91/132) and ciprofloxacin (62.1 %, 82/132). Few isolates were resistant to nitrofurantoin (5.9 %, 4/67) and meropenem (18.2 %, 24/132) (**Figure (1B); Supplementary Table 2**). The prevalence of susceptible, R1 and R2 categories ^{54, 55} were 19.6 %, 7.6 % and 9.1 % respectively; 16.6 % (n = 22/132) of isolates were MDR, while 46.9 % of isolates (n=62/132) were XDR (**Figure (1C)**). *Ent. hormaechei* (60 %, 3/5) and *K. pneumoniae* (100 %, n = 11) were associated with the highest prevalence of MDR and XDR, respectively.

AMR determinants

A wide range of AMR determinants for β -lactams, quinolones, and aminoglycosides have been reported in different uropathogens ⁵⁶⁻⁵⁹. The main AMR genes encoded in our genomes

were predicted using ResFinder, CARD and AMRFinder (**Supplementary Table 3**). New AMR variants are reported here for uropathogens in Egypt: *OXA-486*, *OXA-488*, *OXA-905*, *IMP-43*, *PDC-35*, *PDC-45*, and *PDC-201* for *Pse. aeruginosa*, and *TEM-176* and *TEM-190* for *Esc. coli*.

WGS-based diagnostics may soon take over phenotypic testing for surveillance purposes, particularly in situations where the low error rate has minimal consequences⁶⁰. Comparison of WGS data and DDT results (with respect to predicted AMR genes and empirical resistance phenotypes) yielded a total concordance of 91.1 %, 85.7 % and 80.3 % for ResFinder, CARD and AMRFinder, respectively. The highest concordance was recorded for *Alcaligenes* (n = 4) species: 97.7 %, AMRFinder; 97.2 %, ResFinder; 100 % CARD (**Supplementary Table 4**).

Variations in discordance were detected for different databases, where the MEs (WGS-R/DDT-S) were 3.9 %, 11.9 % and 15.7 %, while VMEs (WGS-S/DDT-R) were 4.9 %, 2.2 % and 3.8% for ResFinder, CARD, and AMRFinder, respectively (**Figure (2)**). The use of an ineffective therapeutic agent in treatment due to a VME could result in treatment failure, while an ME may restrict therapeutic choices and create complications in the treatment process. The more severe impact of VMEs is evident in FDA regulations for the approval of diagnostic tests, where the FDA mandates that VMEs be kept below 1.5 % and MEs below 3 % for the approval of a new AMR diagnostic test or device⁶¹.

Concordance for aminoglycosides, quinolones, and miscellaneous antibiotics exceeded 89.3 % in all databases. Meropenem was associated with highest discordance in AMRFinder (48.4 %) and CARD (47.7 %), but showed 10.6 % discordance in ResFinder. The range of total discordance of β -lactam antibiotics for the three databases was 10.6–20.4 % ResFinder, 9–47.7 % CARD and 15.1–48.4 % AMRFinder (**Supplementary Table 5**).

In a comparison of CARD and ResFinder for two global collections and a total of 2,587 isolates comprising five pathogens of medical importance, ME rates were higher in CARD (42 %) than ResFinder (25 %). However, CARD showed almost no VMEs (1.1 %) compared with ResFinder (4.4 %)³⁷. In comparison, another report showed an overall concordance of 91 % for ResFinder while ME and VME rates were 6.2 % and 2.1 %, respectively, for 488 different isolates, where *Pse. aeruginosa* had the highest percentage of discordant results (44.4 %)⁴⁴. In agreement with previous studies^{16, 44, 60}, *Pseudomonas* showed the highest discordance here, and

across the three databases used in our study: 31.2 % for AMRFinder, 28.9 % for ResFinder, and 21.2 % for CARD.

In this study, 34.5 % of MEs tested phenotypically “*susceptible, increased exposure*” and were therefore considered as not resistant due to the EUCAST update of susceptibility definitions 2019 “*by the pooling of S and I isolates together into one category*”⁶². Notably in another study, 22.7 % of MEs tested phenotypically as “*susceptible at higher exposure*” in 234 non-duplicate *Esc. coli* strains⁴⁵. Discordant bioinformatic predictions of AMR from WGS have previously been shown through analysis of seven clinical isolates with three different databases (ResFinder, CARD and ARG-ANNOT) and where the overall ME “false positives” and VME “false negatives” were 20 % and 14 %, respectively⁶³. In the current study, the overall MEs and VME in three databases were 10.9 % and 3.6 %, respectively. The reduction in MEs and VMEs compared to earlier study⁶³ may in part reflect improvements in curation and coverage of AMR genes through regular database updates over time. The high MEs rate may be also due to low expression levels of enzymes, insufficient understanding of gene regulation, mutations in intergenic regions and *de novo* rRNA mutations.

ResFinder/PointFinder (overall concordance, 91.1 %) showed better *in silico* AMR prediction than CARD (85.7 %) and AMRFinder with Point mutations (80.3 %) (**Figure (2)**). PointFinder tackles chromosome mutation-mediated resistance only in *Esc. coli* and *Klebsiella*. A combined AMR prediction from ResFinder and PointFinder for ciprofloxacin was shown to result in improved prediction performance and a reduced VME rate³⁷.

For accurate *in silico* AMR predictions, our understanding of phenotypic–genotypic correlations must be improved to reach FDA requirements for clinical microbiology diagnostic testing. Excluding genes associated with efflux pump mechanisms from detected AMR genes has been recommended³⁷. Identity thresholds for AMR gene detection should be ≥ 90 % for different databases⁶³. Point mutations leading to AMR should be considered alongside acquired resistance determinants, especially if they are reported from specific clinical sources such as urine, as the most common site for discordance was urine for both *Enterobacterales* and *Pse. aeruginosa*, and mutations in *gyrA* are associated with quinolone resistance of *Enterobacteriaceae* in urine⁶⁴⁻⁶⁶. Databases should replace general terminology (β -lactams, quinolones, etc.) of antimicrobial class as a prediction to confer resistance to all members of the same antimicrobial class, so *in silico*

AMR prediction should be tailored to antibiotics themselves rather than antibiotic classes to avoid MEs. For example, the aminoglycoside gene *aac(6')-1* leads to amikacin and tobramycin resistance, while *aac(3)-IIa* leads to gentamicin and tobramycin resistance⁶⁷. This was reported recently as poor correlation (85.9 %) for aminoglycosides between EUCAST and WGS data using ARG-ANNOT and CARD databases, and the authors developed a phenotype-based algorithm to reflect mechanisms responsible for the corresponding aminoglycoside resistance phenotypes⁶⁸.

Curation of *in silico* databases is required. Some AMR genes listed in databases are not truly responsible for AMR. For example, the *crpP* gene identified by ResFinder and AMRFinder was previously known as a novel ciprofloxacin-modifying enzyme⁶⁹. However, recent studies showed that *crpP* presence is not always associated with ciprofloxacin resistance⁷⁰⁻⁷². In this study *crpP* was responsible for 7/8 MEs generated by AMRFinder and all seven MEs generated by ResFinder when analysing ciprofloxacin resistance. *crpP* was detected in only one strain (Me73B) resistant to ciprofloxacin and was the only quinolone resistance determinant detected by ResFinder for that strain. However, AMRFinder also detected a point mutation in *gyrA*, T83I, which most likely is the true explanation of ciprofloxacin resistance in Me73B.

Variations between databases can also be found. For example, the AMR phenotype associated with the gene *armA* by ResFinder was amikacin and tobramycin, while AMRFinder predicted resistance to gentamicin. Moreover, the phenotypic prediction for *aac(6')-Ic* in ResFinder and CARD is resistance to amikacin and tobramycin, while AMRFinder only provides aminoglycoside as subclass.

The *smeF* gene in *Stenotrophomonas maltophilia* is responsible for quinolones resistance⁷³ and was detected by all databases. However, its classification varied: *smeF* was excluded from both CARD and ResFinder as it is considered an efflux pump gene, whereas AMRFinder included it as a resistance determinant for quinolones. This discrepancy contributed to the difference in MEs between ResFinder and AMRFinder for ciprofloxacin in our dataset, with ResFinder showing seven MEs and AMRFinder eight MEs (**Figure (2)**). Although both ResFinder and AMRFinder identified the gene from the exact same contig region with identical coverage, the predicted gene names and associated resistance prediction differed in isolate Me21.

ResFinder annotated the gene as *aac(6')-Ib3*, predicting resistance to amikacin and tobramycin, while AMRFinder identified it as *aac(6')-Ib*, predicting resistance to gentamicin.

Detailed analyses of UPEC isolates

Pangenomic approaches highlight genetic diversity, revealing adaptations, virulence factors and resistance traits among collections of genomes. Several pangenome studies of UPEC strains have been performed for both global collections⁷⁴ and individual countries, including Ireland and Bangladesh^{75, 76}, with the smallest pangenome consisting of at least 16,797 genes in total, the number of core genes ranging from 2,926 to 2,945, and the number of unique genes ranging from 2,900 to 15,266. The UPEC pangenome (n = 67 isolates) here consisted of 10,453 genes: 3,136 (30 %) core; 6,303 (60.3 %) accessory; 1014 (9.7 %) unique.

UPEC isolates from phylogroup B2 were responsible for most of the reported UTIs (n = 36/67, 53.7 %) (**Figure (3)**), in agreement with a previous Egyptian study⁷⁷. We identified 29 different serotypes (**Supplementary Table 1**), with phylogroup B2 serotypes O25:H4 (n = 12/67, 17.9 %) and O75:H5 (n = 11/67, 16.4 %) most common. Although phylogroup B2 strains are uncommon among the commensal microbiota, they are highly virulent when present; B2 strains have been shown to persist in the intestinal microbiota of infants⁷⁸, while also being associated with wildlife and livestock⁷⁹, and herbivorous and omnivorous mammals⁸⁰. A previous Egyptian study showed the most prevalent phylogroup in the country was A followed by B2⁸¹. Other phylogroups detected by us were B1 (19.4 %), A (11.9 %), D (7.4 %), F (5.9 %) and C (1.4 %).

The principal virulence genes associated with all UPEC strains (n = 67) were type 1 fimbriae (*fimH*), outer membrane protein (*ompA*) and enterobactin (*entS*). Curli fibers (*csgA* 98 %, n = 66), common pilus (*ecp* 95 %, n = 64), nutritional and metabolic factors (*fyuA* 79 %, n = 53, *chuA* 67 %, n = 45), K1 capsule type (*kpsM* 66 %, n = 44), aerobactin (*iutA* 60 %, n = 40), P fimbriae (*papC* 52 %, n = 35, *papG* 33 %, n = 22), α -hemolysin and iron siderophores (*hlyA* and *iroN* 24 %, n = 16), exotoxins (*cnf1* 13 %, n = 9), and S fimbriae and F1C fimbriae (*sfa* and *foc* 6 %, n = 4) were also represented (**Figure (3); Supplementary Table 6**). *fimH* and *sfaX* were present in all phylogroups; all other *sfa* genes were only associated with phylogroup B2. *entD*

was only in phylogroup B1. *cnf-1* was present in all phylogroup B2 isolates and only one isolate (Me87) from B1.

fim, *ecp* and *csg* gene families have previously been found to be core to the UPEC pangenome⁷⁶. In contrast, a PCR-based study from Mansoura reported a much higher prevalence of afimbrial adhesins such as *afa/dra* (14 %) and S-fimbriae (*sfaS*, 60.6 %) ⁸², which were largely absent from our dataset. Additionally, our data show a higher prevalence of P fimbriae genes (*papG*, 33 %; *papC*, 52 %) compared to ⁸², where genes were found only in only 21.3 % and 49.3 % of isolates, respectively. Similarly, our detection of iron acquisition genes *iutA* (60 %) and *chuA* (67 %) exceeds their reported values of 34.6 % and 54 % ⁸². Notably, *cnf1* (cytotoxic necrotizing factor 1) was much rarer in our dataset (13 %) compared to 42 % in ⁸², suggesting potential differences in strain pathogenicity. Overall, our results highlight a strong presence of core adhesion and iron acquisition systems in UPEC strains, with notable variations in toxin-associated genes compared to previous studies. These findings emphasize the genetic diversity of UPEC virulence factors and the importance of comparing detection methods when assessing pathogenic potential.

The β -lactam resistance genes *bla*_{OXA-1} and *bla*_{TEM-1} were present in all phylogroups, *bla*_{TEM-176} was restricted to phylogroup B1, while other *bla*_{TEM} variants were present in only phylogroup A. *dha* gene variants were restricted to phylogroup B2, while *bla*_{NDM-5} was absent from B2. *bla*_{CTX-M-15} was present in all phylogroups, while *bla*_{CTX-M-14} and *bla*_{CTX-M-16} were present in phylogroup F only. In Egypt, *bla*_{CTX-M-15} is the most predominant ESBL-producing type among uropathogenic and diarrheagenic *Esc. coli* strains ⁸³. In terms of quinolone resistance, *qnrB4* was present in phylogroup B2 only, while *qnrS1* was present in all phylogroups. Mutations of *gyrA* and *parC* genes were present in all phylogroups. Genes of aminoglycoside-modifying enzymes were present in all phylogroups.

UPEC isolates were most susceptible to nitrofurantoin (3 % resistance). Nitrofurantoin was reported as a drug of choice for treating UTIs in previous Egyptian studies ^{84, 85}, but it has therapeutic limitations due to side-effects in certain patient groups ⁸⁶. *In silico* AMR prediction using CARD showed 95.5 % concordance with phenotypic results (**Figure (2)**). In a recent Egyptian study, the resistance level to nitrofurantoin was 51 % in *Klebsiella* spp., but it remains effective for *Esc. coli* UTIs in males and females (92 % sensitivity). In comparison, *Esc. coli* and

Klebsiella spp. showed elevated resistance rates to cephalosporins: 75 % and 81 %, respectively⁸⁷.

Conclusion

Our study provides crucial insights into the genomic and AMR landscape of CAUTI-associated uropathogens in Egypt, highlighting the importance of understanding AMR phenotype–genotype discordance in this setting. These findings reinforce the need for integrated phenotypic and genomic surveillance strategies to improve AMR diagnostics, guide treatment decisions, and inform infection control policies in resource-limited settings.

ACKNOWLEDGEMENTS

We would like to Dr Essam Elsayy and staff of Urology and Nephrology Centre, Mansoura, Egypt for providing the clinical isolates used in this study.

ME – did all phenotypic work; extracted DNA for sequencing; all bioinformatics associated with clinical isolates; characterized the AMR and virulence genes encoded by the isolates and their plasmids; MLST analysis and summary; interpreted virulence and AMR data; pangenome analysis. JCT – pangenome analysis and associated supervision. NH, DN – library preparation and genome sequencing of some clinical isolates; LH – sourmash analyses; supervised the study. ME, JCT and LH wrote the original version of the manuscript, and all authors approved the final version.

FUNDING

This work was supported by The Egyptian Ministry of Higher Education & Scientific Research represented by The Egyptian Bureau for Cultural & Educational Affairs in London. NH was funded through a placement studentship provided by Department of Biosciences, Nottingham Trent University. Computing resources used in this study were funded through the Research Contingency Fund of Nottingham Trent University.

365 **TRANSPARENCY DECLARATIONS**

366 The authors declare that there are no conflicts of interest.

REFERENCES




1. Tandogdu Z, Wagenlehner FME. Global epidemiology of urinary tract infections. *Current Opinion in Infectious Diseases* 2016; **29**: 73-9.
2. European Antimicrobial Resistance C. The burden of bacterial antimicrobial resistance in the WHO European region in 2019: a cross-country systematic analysis. *Lancet Public Health* 2022; **7**: e897-e913.
3. Allegranzi B, Bagheri Nejad S, Combescure C et al. Burden of endemic health-care-associated infection in developing countries: systematic review and meta-analysis. *Lancet (London, England)* 2011; **377**: 228-41.
4. Öztürk R, Murt A. Epidemiology of urological infections: a global burden. *World Journal of Urology* 2020; **38**: 2669-79.
5. WHO. Global antimicrobial resistance and use surveillance system (GLASS) report: early implementation 2016-2017. In: Organization WH, ed: World Health Organization, 2018.
6. WHO. Global antimicrobial resistance and use surveillance system (GLASS) report: 2022. World Health Organization, 2022.
7. See I, Lessa FC, ElAta OA et al. Incidence and pathogen distribution of healthcare-associated infections in pilot hospitals in Egypt. *Infect Control Hosp Epidemiol* 2013; **34**: 1281-8.
8. Talaat M, El-Shokry M, El-Kholy J et al. National surveillance of health care-associated infections in Egypt: Developing a sustainable program in a resource-limited country. *Am J Infect Control* 2016; **44**: 1296-301.
9. El-Kholy A, El-Mahallawy HA, Elsharnouby N et al. Landscape of Multidrug-Resistant Gram-Negative Infections in Egypt: Survey and Literature Review. *Infect Drug Resist* 2021; **14**: 1905-20.
10. Kotb S, Lyman M, Ismail G et al. Epidemiology of Carbapenem-resistant Enterobacteriaceae in Egyptian intensive care units using National Healthcare-associated Infections Surveillance Data, 2011-2017. *Antimicrob Resist Infect Control* 2020; **9**: 2.
11. Zakaria AS, Edward EA, Mohamed NM. Genomic Insights into a Colistin-Resistant Uropathogenic Escherichia coli Strain of O23:H4-ST641 Lineage Harboring mcr-1.1 on a Conjugative IncHI2 Plasmid from Egypt. *Microorganisms* 2021; **9**.
12. Zakaria AS, Edward EA, Mohamed NM. Pathogenicity Islands in Uropathogenic Escherichia coli Clinical Isolate of the Globally Disseminated O25:H4-ST131 Pandemic Clonal Lineage: First Report from Egypt. *Antibiotics (Basel)* 2022; **11**.
13. Sherif M, Palmieri M, Mirande C et al. Whole-genome sequencing of Egyptian multidrug-resistant Klebsiella pneumoniae isolates: a multi-center pilot study. *European Journal of Clinical Microbiology & Infectious Diseases* 2021; **40**: 1451-60.
14. Edward EA, Mohamed NM, Zakaria AS. Whole Genome Characterization of the High-Risk Clone ST383 Klebsiella pneumoniae with a Simultaneous Carriage of bla(CTX-M-14) on IncL/M Plasmid and bla(CTX-M-15) on Convergent IncHI1B/IncFIB Plasmid from Egypt. *Microorganisms* 2022; **10**.
15. Baiomy AA, Serry FE, Kadry AA et al. Genome Analysis of Pseudomonas aeruginosa Strains from Chronically Infected Patients with High Levels of Persister Formation. *Pathogens* 2023; **12**.
16. Eladawy M, Thomas JC, Hoyles L. Phenotypic and genomic characterization of Pseudomonas aeruginosa isolates recovered from catheter-associated urinary tract infections in an Egyptian hospital. *Microb Genom* 2023; **9**.
17. Kohler P, Tijet N, Kim HC et al. Dissemination of Verona Integron-encoded Metallo-beta-lactamase among clinical and environmental Enterobacteriaceae isolates in Ontario, Canada. *Sci Rep* 2020; **10**: 18580.

18. van Duin D, Perez F, Rudin SD et al. Surveillance of carbapenem-resistant *Klebsiella pneumoniae*: tracking molecular epidemiology and outcomes through a regional network. *Antimicrob Agents Chemother* 2014; **58**: 4035-41.
19. Hassuna NA, Khairalla AS, Farahat EM et al. Molecular characterization of Extended-spectrum β lactamase- producing *E. coli* recovered from community-acquired urinary tract infections in Upper Egypt. *Sci Rep* 2020; **10**: 2772.
20. Negus D, Foster G, Hoyles L. *Lelliottia amnigena* recovered from the lung of a harbour porpoise, and comparative analyses with *Lelliottia* spp. *Access Microbiol* 2023; **5**.
21. Schwengers O, Jelonek L, Dieckmann MA et al. Bakta: rapid and standardized annotation of bacterial genomes via alignment-free sequence identification. *Microb Genom* 2021; **7**.
22. Pierce NT, Irber L, Reiter T et al. Large-scale sequence comparisons with sourmash. *F1000Res* 2019; **8**: 1006.
23. Jolley KA, Bliss CM, Bennett JS et al. Ribosomal multilocus sequence typing: universal characterization of bacteria from domain to strain. *Microbiology (Reading, England)* 2012; **158**: 1005-15.
24. Jain C, Rodriguez-R LM, Phillippy AM et al. High throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nature communications* 2018; **9**: 1-8.
25. Bessonov K, Laing C, Robertson J et al. ECTyper: in silico *Escherichia coli* serotype and species prediction from raw and assembled whole-genome sequence data. *Microbial genomics* 2021; **7**.
26. Beghain J, Bridier-Nahmias A, Le Nagard H et al. ClermonTyping: an easy-to-use and accurate in silico method for *Escherichia* genus strain phylotyping. *Microbial genomics* 2018; **4**.
27. Jolley KA, Bray JE, Maiden MCJ. Open-access bacterial population genomics: BIGSdb software, the PubMLST.org website and their applications. *Wellcome open research* 2018; **3**: 124.
28. Wirth T, Falush D, Lan R et al. Sex and virulence in *Escherichia coli*: an evolutionary perspective. *Molecular microbiology* 2006; **60**: 1136-51.
29. Bai L, Xia S, Lan R et al. Isolation and characterization of cytotoxic, aggregative *Citrobacter freundii*. *PloS one* 2012; **7**: e33054.
30. Diancourt L, Passet V, Verhoef J et al. Multilocus sequence typing of *Klebsiella pneumoniae* nosocomial isolates. *J Clin Microbiol* 2005; **43**: 4178-82.
31. Curran B, Jonas D, Grundmann H et al. Development of a multilocus sequence typing scheme for the opportunistic pathogen *Pseudomonas aeruginosa*. *J Clin Microbiol* 2004; **42**: 5644-9.
32. Diancourt L, Passet V, Nemec A et al. The population structure of *Acinetobacter baumannii*: expanding multiresistant clones from an ancestral susceptible genetic pool. *PloS one* 2010; **5**: e10034.
33. Miyoshi-Akiyama T, Hayakawa K, Ohmagari N et al. Multilocus sequence typing (MLST) for characterization of *Enterobacter cloacae*. *PloS one* 2013; **8**: e66358.
34. McArthur AG, Wagglechner N, Nizam F et al. The comprehensive antibiotic resistance database. *Antimicrob Agents Chemother* 2013; **57**: 3348-57.
35. Florensa AF, Kaas RS, Clausen P et al. ResFinder - an open online resource for identification of antimicrobial resistance genes in next-generation sequencing data and prediction of phenotypes from genotypes. *Microbial genomics* 2022; **8**.
36. Feldgarden M, Brover V, Haft DH et al. Validating the AMRFinder Tool and Resistance Gene Database by Using Antimicrobial Resistance Genotype-Phenotype Correlations in a Collection of Isolates. *Antimicrob Agents Chemother* 2019; **63**.
37. Mahfouz N, Ferreira I, Beisken S et al. Large-scale assessment of antimicrobial resistance marker databases for genetic phenotype prediction: a systematic review. *Journal of Antimicrobial Chemotherapy* 2020; **75**: 3099-108.
38. Tonkin-Hill G, MacAlasdair N, Ruis C et al. Producing polished prokaryotic pangenomes with the Panaroo pipeline. *Genome biology* 2020; **21**: 180.
39. Loytynoja A. Phylogeny-aware alignment with PRANK. *Methods Mol Biol* 2014; **1079**: 155-70.

40. Kuck P, Longo GC. FASconCAT-G: extensive functions for multiple sequence alignment preparations concerning phylogenetic studies. *Front Zool* 2014; **11**: 81.
41. Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics (Oxford, England)* 2014; **30**: 1312-3.
42. Letunic I, Bork P. Interactive Tree Of Life (iTOL) v4: recent updates and new developments. *Nucleic Acids Res* 2019; **47**: W256-W9.
43. Chen L, Yang J, Yu J et al. VFDB: a reference database for bacterial virulence factors. *Nucleic Acids Res* 2005; **33**: D325-8.
44. Rebelo AR, Bortolaia V, Leekitcharoenphon P et al. One Day in Denmark: Comparison of Phenotypic and Genotypic Antimicrobial Susceptibility Testing in Bacterial Isolates From Clinical Settings. *Front Microbiol* 2022; **13**: 804627.
45. Vanstokstraeten R, Pierard D, Crombe F et al. Genotypic resistance determined by whole genome sequencing versus phenotypic resistance in 234 Escherichia coli isolates. *Sci Rep* 2023; **13**: 449.
46. Mohamed ES, Khairy RMM, Abdelrahim SS. Prevalence and molecular characteristics of ESBL and AmpC β -lactamase producing Enterobacteriaceae strains isolated from UTIs in Egypt. *Antimicrob Resist Infect Control* 2020; **9**: 198.
47. Shrief R, El-Ashry AH, Mahmoud R et al. Effect of Colistin, Fosfomycin and Meropenem/Vaborbactam on Carbapenem-Resistant Enterobacterales in Egypt: A Cross-Sectional Study. *Infect Drug Resist* 2022; **15**: 6203-14.
48. Baz AMK, El-Agamy OAE, Ibrahim AM. Incidence of urinary tract infection in neonates with significant indirect Hyperbilirubinemia of unknown etiology: case-control study. *Ital J Pediatr* 2021; **47**: 35.
49. Hassan R, El-Gilany AH, Abd Elaal AM et al. An overview of healthcare-associated infections in a tertiary care hospital in Egypt. *Infect Prev Pract* 2020; **2**: 100059.
50. Rajni E, Jain A, Garg VK et al. Providencia Causing Urinary Tract Infections: Are We Reaching a Dead End? *Indian J Crit Care Med* 2022; **26**: 446-51.
51. Nouh K, Kasem A, Shaher H et al. *THE EGYPTIAN UROLOGICAL GUIDELINES, First Edition, Chapter XI: Urinary Tract Infections Guidelines.*, 2021.
52. Elshamy AA, Aboshanab KM, Yassien MA et al. Prevalence of plasmid-mediated resistance genes among multidrug-resistant uropathogens in Egypt. *African health sciences* 2020; **20**: 190-8.
53. Esmaeel NE, Gerges MA, Hosny TA et al. Detection of Chromosomal and Plasmid-Mediated Quinolone Resistance Among Escherichia coli Isolated from Urinary Tract Infection Cases; Zagazig University Hospitals, Egypt. *Infect Drug Resist* 2020; **13**: 413-21.
54. Magiorakos AP, Srinivasan A, Carey RB et al. Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. *Clin Microbiol Infect* 2012; **18**: 268-81.
55. Bitew A, Tsige E. High Prevalence of Multidrug-Resistant and Extended-Spectrum beta-Lactamase-Producing Enterobacteriaceae: A Cross-Sectional Study at Arsho Advanced Medical Laboratory, Addis Ababa, Ethiopia. *J Trop Med* 2020; **2020**: 6167234.
56. Jomehzadeh N, Saki M, Ahmadi K et al. The prevalence of plasmid-mediated quinolone resistance genes among Escherichia coli strains isolated from urinary tract infections in southwest Iran. *Mol Biol Rep* 2022; **49**: 3757-63.
57. Tickler IA, Kawa D, Obradovich AE et al. Characterization of Carbapenemase- and ESBL-Producing Gram-Negative Bacilli Isolated from Patients with Urinary Tract and Bloodstream Infections. *Antibiotics (Basel)* 2023; **12**.
58. de Sousa T, Hebraud M, Alves O et al. Study of Antimicrobial Resistance, Biofilm Formation, and Motility of Pseudomonas aeruginosa Derived from Urine Samples. *Microorganisms* 2023; **11**.

59. Zeng Q, Xiao S, Gu F et al. Antimicrobial Resistance and Molecular Epidemiology of Uropathogenic Escherichia coli Isolated From Female Patients in Shanghai, China. *Frontiers in cellular and infection microbiology* 2021; **11**: 653983.
60. Ellington M, Ekelund O, Aarestrup FM et al. The role of whole genome sequencing in antimicrobial susceptibility testing of bacteria: report from the EUCAST Subcommittee. *Clinical microbiology and infection* 2017; **23**: 2-22.
61. Jorgensen JH, Ferraro MJ. Antimicrobial susceptibility testing: a review of general principles and contemporary practices. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America* 2009; **49**: 1749-55.
62. Nabal Diaz SG, Algara Robles O, Garcia-Lechuz Moya JM. New definitions of susceptibility categories EUCAST 2019: clinic application. *Rev Esp Quimioter* 2022; **35 Suppl 3**: 84-8.
63. Doyle RM, O'Sullivan DM, Aller SD et al. Discordant bioinformatic predictions of antimicrobial resistance from whole-genome sequencing data of bacterial isolates: an inter-laboratory study. *Microb Genom* 2020; **6**.
64. Shenagari M, Bakhtiari M, Mojtahedi A et al. High frequency of mutations in *gyrA* gene associated with quinolones resistance in uropathogenic Escherichia coli isolates from the north of Iran. *Iran J Basic Med Sci* 2018; **21**: 1226-31.
65. Ku PM, Hobbs DA, Gilmore M et al. 1234. Can Susceptibility to One Carbapenem be Conferred to Another? Frequency of Discordance in Gram-negative Clinical Isolates. *Open Forum Infectious Diseases*, p. S706-S7, Oxford University Press US.
66. Thompson D, Xu J, Ischia J et al. Fluoroquinolone resistance in urinary tract infections: Epidemiology, mechanisms of action and management strategies. *BJUI Compass* 2024; **5**: 5-11.
67. Krause KM, Serio AW, Kane TR et al. Aminoglycosides: An Overview. *Cold Spring Harb Perspect Med* 2016; **6**.
68. Mancini S, Marchesi M, Imkamp F et al. Population-based inference of aminoglycoside resistance mechanisms in Escherichia coli. *EBioMedicine* 2019; **46**: 184-92.
69. Chávez-Jacobo VM, Hernández-Ramírez KC, Romo-Rodríguez P et al. CrpP Is a Novel Ciprofloxacin-Modifying Enzyme Encoded by the Pseudomonas aeruginosa pUM505 Plasmid. *Antimicrob Agents Chemother* 2018; **62**.
70. Hernández-García M, García-Castillo M, García-Fernández S et al. Presence of Chromosomal crpP-like Genes Is Not Always Associated with Ciprofloxacin Resistance in Pseudomonas aeruginosa Clinical Isolates Recovered in ICU Patients from Portugal and Spain. *Microorganisms* 2021; **9**.
71. López M, Rojo-Bezares B, Chichón G et al. Resistance to Fluoroquinolones in Pseudomonas aeruginosa from Human, Animal, Food and Environmental Origin: The Role of CrpP and Mobilizable ICEs. *Antibiotics* 2022; **11**: 1271.
72. Zubyk HL, Wright GD. CrpP Is Not a Fluoroquinolone-Inactivating Enzyme. *Antimicrob Agents Chemother* 2021; **65**: e0077321.
73. Jia W, Wang J, Xu H et al. Resistance of Stenotrophomonas maltophilia to Fluoroquinolones: Prevalence in a University Hospital and Possible Mechanisms. *Int J Environ Res Public Health* 2015; **12**: 5177-95.
74. Fatima S, Akbar A, Irfan M et al. Virulence Factors and Antimicrobial Resistance of Uropathogenic Escherichia coli EQ101 UPEC Isolated from UTI Patient in Quetta, Balochistan, Pakistan. *Biomed Res Int* 2023; **2023**: 7278070.
75. Whelan S, Bottacini F, Buttner C et al. Whole genome sequencing of uropathogenic E. coli from Ireland reveals diverse resistance mechanisms and strong correlation with phenotypic (EUCAST) susceptibility testing. *Infect Genet Evol* 2024; **121**: 105600.
76. Hossain M, Tabassum T, Rahman A et al. Genotype-phenotype correlation of beta-lactamase-producing uropathogenic Escherichia coli (UPEC) strains from Bangladesh. *Sci Rep* 2020; **10**: 14549.


77. Farahat EM, Hassuna NA, Hammad AM et al. Distribution of integrons and phylogenetic groups among *Escherichia coli* causing community-acquired urinary tract infection in Upper Egypt. *Canadian Journal of Microbiology* 2021; **67**: 451-63.
78. Nowrouzian FL, Adlerberth I, Wold AE. Enhanced persistence in the colonic microbiota of *Escherichia coli* strains belonging to phylogenetic group B2: role of virulence factors and adherence to colonic cells. *Microbes Infect* 2006; **8**: 834-40.
79. Aguirre-Sanchez JR, Valdez-Torres JB, Del Campo NC et al. Phylogenetic group and virulence profile classification in *Escherichia coli* from distinct isolation sources in Mexico. *Infect Genet Evol* 2022; **106**: 105380.
80. Gordon DM, Cowling A. The distribution and genetic structure of *Escherichia coli* in Australian vertebrates: host and geographic effects. *Microbiology (Reading, England)* 2003; **149**: 3575-86.
81. El-Mahdy R, Mahmoud R, Shrief R. Characterization of *E. coli* Phylogroups Causing Catheter-Associated Urinary Tract Infection. *Infect Drug Resist* 2021; **14**: 3183-93.
82. El-Baz R, Said HS, Abdelmegeed ES et al. Characterization of virulence determinants and phylogenetic background of multiple and extensively drug resistant *Escherichia coli* isolated from different clinical sources in Egypt. *Appl Microbiol Biotechnol* 2022; **106**: 1279-98.
83. Abdelrahim SS, Fouad M, Abdallah N et al. Comparative Study of CTX-M-15 Producing *Escherichia coli* ST131 Clone Isolated from Urinary Tract Infections and Acute Diarrhoea. *Infect Drug Resist* 2021; **14**: 4027-38.
84. Shash RY, Elshimy AA, Soliman MY et al. Molecular Characterization of Extended-Spectrum β -Lactamase Enterobacteriaceae Isolated from Egyptian Patients with Community- and Hospital-Acquired Urinary Tract Infection. *Am J Trop Med Hyg* 2019; **100**: 522-8.
85. Elzayat M, Barnett-Vanes A, Dabour MF et al. Prevalence of undiagnosed asymptomatic bacteriuria and associated risk factors during pregnancy: a cross-sectional study at two tertiary centres in Cairo, Egypt. *BMJ Open* 2017; **7**: e013198.
86. Recht J, Chansamouth V, White NJ, Ashley EA. Nitrofurantoin and glucose-6-phosphate dehydrogenase deficiency: a safety review. *JAC Antimicrob Resist*. 2022 May 3;4(3):dlac045. doi: 10.1093/jacamr/dlac045. PMID: 35529053; PMCID: PMC9070801.
87. Tarek A, Abdalla S, Dokmak NA et al. Bacterial Diversity and Antibiotic Resistance Patterns of Community-Acquired Urinary Tract Infections in Mega Size Clinical Samples of Egyptian Patients: A Cross-Sectional Study. *Cureus* 2024; **16**: e51838.



587

588

589 **Figure (1): AMR profiles of uropathogens (n = 132 isolates) included in this study.** A) Dendrogram generated from comparison of
 590 sourmash signatures for uropathogens showing their species affiliations and whether they are MDR or XDR. Phylogroup affiliations
 591 are given for *Esc. coli* isolates. B) Resistance summary for the 132 uropathogens using seven different antimicrobial classes according
 592 to EUCAST guidelines. (C) Data presented using the category designations suggested by ^{54, 55}. Penicillins: piperacillin/tazobactam.
 593 Monobactams: aztreonam. Cephalosporins: ceftazidime and cefepime. Carbapenems: doripenem and meropenem. Quinolones:
 594 ciprofloxacin and levofloxacin. Aminoglycosides: amikacin and tobramycin. Miscellaneous: sulfamethoxazole/trimethoprim and
 595 nitrofurantoin.



596

597

Figure (2): AMR genotype–phenotype discordance summarized for the 132 isolates included in this study. (A, B) Percentage and number of genotype–phenotype concordances and discordances are shown, when comparing WGS data and DDT results using ResFinder, CARD, and AMRFinder. The bars are coloured according to Concordance or Discordance (refer to colour legend at the top of the figure). White numbers on top of the bars correspond to the respective number of results. (A) Species-level data, (B) antimicrobial-level data. Numbers in parentheses in (A) correspond to the number of isolates per species / the number of isolate-antimicrobial combinations. (A) Others: *Paenaltcaligenes suwonensis*, *Klebsiella ornithinolytica*, *Citrobacter portucalensis*, *Leclercia adecarboxylata*, *Morganella morganii*, *Stenotrophomonas maltophilia*, *Acinetobacter baumannii*, and *Achromobacter xylosoxidans*. (B) Nitrofurantoin concordance was analyzed only via CARD. Levofloxacin and doripenem were not analyzed with ResFinder/PointFinder. Piperacillin/tazobactam (TZP), aztreonam (ATM), ceftazidime (CAZ), cefepime (FEP), doripenem (DOR), meropenem (MEM), ciprofloxacin (CIP), levofloxacin (LEV), amikacin (AK), tobramycin (TOB), sulfamethoxazole/trimethoprim (SXT), and nitrofurantoin (NIF). Major errors (ME) are WGS-R/DDT-S, while very major error (VME) are WGS-S/DDT-R.

