

Auditory perceptual history is communicated through alpha oscillations

Hao Tam Ho^{1,3,*}, Johahn Leung^{1,2}, David C. Burr^{1,5,*}, David Alais¹, Maria Concetta Morrone^{3,4}

¹ School of Psychology, University of Sydney, Brennan MacCallum Building A18, Manning Rd, Camperdown, NSW 2006, Australia

² School of Medical Sciences, University of Sydney, Anderson Stuart Building F13 Eastern Avenue, NSW 2006, Australia

³ Department of Translational Research on New Technologies in Medicine and Surgery, University of Pisa, via San Zeno 31, 56123 Pisa, Italy

⁴ Scientific Institute Stella Maris, Viale del Tirreno, 331, 56018 Calambrone, Pisa, Italy

⁵ Department of Neuroscience, Psychology, Pharmacology, and Child Health, University of Florence, Via di San Salvi 12, 50139 Florence, Italy

Abbreviated title: Prior information influences criterion oscillation

***Corresponding authors:** tam.ho@sydney.edu.au, dave@in.cnr.it

Funding: The research was supported by the Australian Research Council Discovery Project (DP150101731) and the European Research Council (FPT/2007-2013) under grant agreement 338866 Ecsplain.

Word count: 5,827

Number of figures: 8

Competing interests: The authors declare that no competing interests exist.

Abstract

Sensory expectations from the accumulation of information over time exert strong predictive biases on forthcoming perceptual decisions. These anticipatory mechanisms help to maintain a coherent percept in a noisy environment. Here we present novel behavioural evidence that past sensory experience biases perceptual decisions rhythmically through alpha oscillations. Participants identified the ear of origin of a brief sinusoidal tone masked by dichotic white noise, and response bias oscillated over time at ~9 Hz. Importantly, the oscillations occurred only for trials preceded by a target to the same ear and lasted for at least two trials. These findings suggest that each stimulus elicits an oscillating memory trace, specific to the ear of origin, which subsequently biases perceptual decisions. This trace is phase-reset by the noise onset of the next trial, and remains within the circuitry of the ear in which it was elicited, modulating the sensory representations in that ear.

Keywords: Alpha oscillation, Auditory perception, Decision criteria, Sensory expectations, Serial dependence and Signal detection theory.

Introduction

It has been long known that perception depends heavily on expectations and perceptual experience. Helmholtz (1867) introduced the concept of *unconscious inference*, suggesting that perception is at least partly *inferential*, or *generative*. Gregory (1997) described perception as a series of *hypotheses* to be verified against sensory data, using many compelling illusions to back his view. Both suggested that perception is a proactive, *predictive* process, where the brain makes best guesses about the world to test against sensory data, updating the guesses as needed.

Recent techniques lend themselves to quantitative study of predictive perception. One example is *serial dependence*: under many conditions, the appearance of images in a sequence depends strongly on the stimulus presented just prior to the current one. Judgements of orientation (Fischer and Whitney, 2014), numerosity (Cicchini et al., 2014), motion (Alais et al., 2017), facial identity or gender (Liberman et al., 2014; Taubert et al., 2016), beauty and even perceived body size (Alexi et al., 2018) are strongly biased towards the previous image. Strong serial biases are also observed in audition, so far only for pitch discrimination (Arzounian et al., 2017; Chambers and Pressnitzer, 2014). To our knowledge, no study has investigated whether other aspects of auditory processing, such as sound localisation, also exhibit serial dependence.

Sequential effects can last for seconds, even minutes (Chopin and Mamassian, 2012), suggesting that perception does not rely solely on the instantaneous stimulation, but on predictions conditioned by events over a long time-course. Counter-intuitively, the biases introduced by serial dependence can lead to more efficient perception (Cicchini et al., 2018): while the biases constitute perceptual errors, these errors are offset by a reduction in response variance and are therefore beneficial overall. As the world tends to remain constant over the

short term (Dong and Atick, 1995; Voss, 1975), past events are good predictors of the future. On the assumption that nothing has physically changed, the previous stimulus acts as a predictor, or a *prior*, to be combined with the current sensory data to enhance the signal. And on average, that predictor is useful in reducing error.

However, not all serial effects are positive. Classical *adaptation aftereffects* are negative (Thompson and Burr, 2009): prolonged inspection of downward motion causes stationary stimuli displayed to the same location to appear to move upwards (Addams, 1834). Similar negative aftereffects have been observed in almost every dimension, including orientation (Gibson and Radner, 1937), colour (Neitz et al., 2002), faces (Leopold et al., 2001), and numerosity (Burr and Ross, 2008). They also exist in audition and cover a similarly wide range of aspects, from auditory motion (Grantham and Wightman, 1979), sound localisation (Kashino and Nishida, 1998) and tone intensity (Reinhardt-Rutland and Anstis, 1982) to gender of human voices (Schweinberger et al., 2008) and even vocal affect (Bestelmeyer et al., 2010). While the classical paradigm usually involves long periods of adaptation, sizeable negative aftereffects can arise after very brief presentations in vision as in audition (Aagten-Murphy and Burr, 2016; Alais et al., 2017, 2015).

What determines whether assimilation or contrastive effects prevail, and how do the two opposing mechanisms interact? One factor is stimulus strength: strong, salient, high-contrast, long-duration stimuli tend to lead to negative aftereffects, while brief, less salient low-contrast stimuli result in positive aftereffects (Kanai and Verstraten, 2005; Pantle et al., 2000; Yoshimoto et al., 2014). Taubert et al. (2016) reported strong positive serial dependence for judgments of the gender of faces (a stable attribute) and negative aftereffects for judgments of emotion (a labile attribute, where change is important) on the same stimuli. However, negative and positive effects may also reflect individual preferences. Abrahamyan et al. (2016) found that some individuals, termed “switchers”, tend to respond opposite to

their previous choice, while others prefer to “remain” with whatever response they gave in the preceding trial. Finally, Chopin and Mamassian (2012) made a surprising observation within the same experiment: stimuli close in time were negatively correlated with the current percept, while those more remote positively. Importantly, their modelling results suggest that both negative and positive dependencies could be explained by a single predictive adaptation mechanism. However, how such a predictive adaptation mechanism could be implemented neurally is unclear.

Generally, the neuronal mechanisms underlying serial dependence are largely unknown. It is assumed that the prior is generated at mid-high levels of analysis, and fed back to early sensory areas, which in turn modify the prior (Friston, 2005; Lee and Mumford, 2003; Summerfield and de Lange, 2014; Summerfield and Koechlin, 2008; Yuille and Kersten, 2006), but we do not know how this information is propagated, nor do we have a grasp on the underlying neural mechanisms. One possibility is that recursive propagation and updating of the prior is related to low-frequency neural oscillations (Friston et al., 2015; Sherman et al., 2016; VanRullen, 2017).

Although controversial, evidence is gathering that synchronous rhythmic neural activity may serve to bind stimulus components into a unitary percept (Buzsáki and Draguhn, 2004; Engel et al., 2001; Gray et al., 1989). Besides the neural evidence in animals and humans, recent psychophysical evidence shows that neural oscillations in mammalian brain modulate perceptual performance, and that these oscillations can be synchronized either by abrupt perceptual stimuli (Fiebelkorn et al., 2013, 2011; Landau and Fries, 2012), or by motor-action (hand or eye movements) (Benedetto et al., 2018, 2016; Benedetto and Morrone, 2017; Tomassini et al., 2017, 2015). Using *signal detection theory* (Green and Swets, 1966; Macmillan and Creelman, 2004), we have recently shown within the same experimental setup that oscillations can occur in both *sensitivity* (accuracy) and *criterion*

(response bias) at different frequencies (theta for sensitivity, alpha for criterion), suggesting separate mechanisms underlying those two perceptual properties (Ho et al., 2017). Oscillations in bias could plausibly reflect the reverberation of recursive error propagation within a generative perception framework.

Alpha oscillations in criteria are consistent with EEG evidence of an association between criteria shifts and modulations of alpha power (Craddock et al., 2017; Haegens et al., 2014; Iemi et al., 2017; Iemi and Busch, 2018; Limbach and Corballis, 2016) and phase (Sherman et al., 2016). Decreases in alpha power consistently coincide with more liberal decision criteria in vision (Iemi et al., 2017; Iemi and Busch, 2018; Limbach and Corballis, 2016) as well as touch (Craddock et al., 2017; Haegens et al., 2014). Notably, the alpha modulation can be observed several hundred milliseconds before stimulus onset (Craddock et al., 2017; Limbach and Corballis, 2016), suggesting that criteria shifts are driven, at least in part, by alpha oscillations. This notion is further supported by the observation that the phase of prestimulus alpha can predict criteria changes (Sherman et al., 2016). Furthermore, the relationship between prestimulus alpha and criteria change was strongly influenced by target expectancy, reinforcing the view that alpha oscillations are involved in sensory anticipation or prediction (de Lange et al., 2013; Klimesch et al., 1998; Mayer et al., 2016; Premereur et al., 2012; Rohenkohl and Nobre, 2011; Sanders et al., 2014).

VanRullen and Macdonald (2012) have proposed an oscillatory mechanism by which past perceptual history is stored in memory. By cross-correlating the EEG response with the corresponding visual stimulus, they showed that random, non-periodic luminance changes elicited a “perceptual echo”, a reverberatory response at ~10 Hz that lasted for at least 1 s. Recent findings have confirmed that this long-lasting oscillation serves to maintain sensory representations over time (Chang et al., 2017; Huang et al., 2018), which could influence subsequent perception.

Given these findings, we hypothesised that the alpha oscillations observed in criteria (Ho et al., 2017) may underlie a predictive mechanism that biases perceptual decisions, giving rise to sequential effects. To test this idea, we asked participants to indicate the ear of origin of a brief monaural sinusoidal tone masked by uncorrelated streams of dichotic white noise in each ear (illustrated in Fig. 1A). The tone was presented at random intervals after the noise onset which served to reset the phase of ongoing neural oscillations. From our previous work (Ho et al., 2017), we expected decision criterion to oscillate at alpha frequencies (8–12 Hz). If these oscillations are instrumental in communicating perceptual expectations, the rhythmic fluctuations in criteria should be contingent on the previous stimuli.




Figure 1. (A) Schematic of a trial. On each trial, white noise was presented simultaneously to both ears for 2 s. A pure tone of 1 kHz and 10 ms duration was delivered with equiprobability to the left or right ear. The SOA was randomly selected from an interval of 0.2-1.2 s post noise onset. The ITI jittered randomly between 1.2-2.2 s. **(B)** Application of signal detection theory (SDT). To look for oscillations in sensitivity and criterion, we first pooled across all 15 participants (creating an aggregate subject) and binned the data with non-overlapping, rectangular windows of 10 ms. For each bin, we calculated d-prime (d') and decision criterion (c) using the hits (H) and false alarm (FA) from the left- and right-target conditions, respectively. (M: misses, CR: correct rejections). The calculations follow the Equations 1-2. **(C)** Schematic illustration of the linear regression analysis based on single trials. We

computed the sine and cosine of the time at which the target was presented ($t = 1, 2 \dots n$) for every tested frequency from 4-12 Hz (in 0.01-Hz steps). Based on the sine and cosine regressors, we estimated parameters β_1 and β_2 (β_0 is a constant) for all responses (coded as 0 = 'left' and 1 = 'right') in y .

Results

Stimulus history affects response bias but not sensitivity

Participants performed a two-alternative forced-choice (2AFC) task which required them to identify the ear of origin of a weak, brief tone (10 ms) embedded in white noise (2 s in duration). The target tone was delivered randomly with equal probability in either ear with a stimulus-onset asynchrony (SOA) randomly drawn from an interval of 0.2–1.2 s post noise onset. Using a staircase procedure, we kept the target's intensity around individuals' thresholds (75 % accuracy).

We first examined whether observer sensitivity and response bias, as measured respectively by d-prime (d') and decision criterion (c) (Eq. 1&2, see also Fig. 1B), were contingent on the preceding stimuli. As shown in Fig. 2A, stimulus history did not influence observer sensitivity. Whether the last target (1-back) and second-last target (2-back) occurred in the left or right ear had no effect on detecting the current target: the difference in sensitivity contingent on the preceding left (blue bars) and right stimuli (red bars) was not statistically significant after multiple comparison correction (with FDR = 5 %), $p = 0.07$ and $p = 0.56$ for 1- and 2-back, respectively. However, Fig. 2B shows that criterion was affected by stimulus history. Curiously, there was no significant 1-back effect (when FDR adjusted), $p = 0.86$, but a strong 2-back effect ($p = 0.0002$) of observer response biases towards the previously presented stimulus, either left or right. The effects remained significant for stimuli

three trials back, $p = 0.008$, and four trials back, $p = 0.009$, but was non-significant five trials back, $p = 0.85$.

We then examined the dependency of the *previous response* (Fig. 2C), which revealed a *negative* influence of one trial back, marginally significant after FDR correction ($p = 0.053$). This response-dependent negative serial dependence could result from response switching, mentioned in the Introduction. However, two trials back, which should not be affected by response switching, showed the same positive aftereffect found for the stimulus-based analysis, $p = 0.0001$ (FDR corrected). Trials further in the past (not shown) had no significant effect, all p 's > 0.05 . Even though not statistically robust, this repulsive effect could account for the stimulus-based null effect in the 1-back (Fig. 2B), discussed further in the Discussion section.

To gain a better understanding of the sequential effects in response bias, we examined the effects in individual subjects. Figure 2D plots the individual biases (differences in responses preceded by left from those preceded by right) for 1-back trials against 2-back trials. All subjects except one (subject 5) showed a positive serial effect for trials that were 2-back, while there was much greater subject variability in the 1-back condition, with 5 participants showing a positive and the remaining 9 showing a negative effect. This pattern of results is consistent with previous reports of individual differences in switching responses from one trial to the next (Abrahamyan et al., 2016).

That both negative and positive effects are observed only for criteria is particularly interesting, as we have previously shown that response biases oscillate at low alpha frequencies, ~ 8 Hz, and there is evidence that alpha oscillations are involved in predictive processes and modulation of decision criteria (de Lange et al., 2013; Sherman et al., 2016). In a preliminary analysis of the temporal dynamics of the 1-back effect, we sorted the individual response data by target SOA (from noise onset) and grouped them into 12 bins of 83 ms,

computed the difference in left and right bias contingent on the preceding trial and averaged across all participants. As can be seen from Fig. 2E, the effect of stimulus history on response bias oscillates smoothly over time, warranting further investigation.




Figure 2. Results of the serial dependence analysis. **(A)** Group mean sensitivity contingent on whether the last (1-back) and second last (2-back) target occurred the left (blue bars) or right ear (red bars). There was no significant influence of stimulus history on observer sensitivity. **(B)** Group mean response bias contingent on the ear of origin of the preceding 1- to 5-back stimuli. The difference between the contingent left and right (blue and red bars, respectively) are significant for the 2-, 3- and 4-back stimuli (FDR corrected). **(C)** Group mean response bias contingent on the response 1-2 trials back. The difference between the contingent left and right (blue and red bars, respectively) is significant for both 1- and 2-back (FDR corrected). **(D)** Individual stimulus-based bias differences in the 1-back plotted against those in the 2-back. These differences were computed by subtracting the contingent left bias (blue bars in (B)) from the contingent right bias (red bars). Except for subject 5, all other subjects show a positive bias difference in the 2-back. In contrast, there was greater subject variability in the 1-back, with only 5 participants showing a positive effect. **(E)** Temporal dynamics of the 1-back effect. The individual response data were grouped into 12 bins of 83 ms. For each bin, we computed the difference in left and right bias contingent on the preceding trial as in (C) and averaged across all participants. Not only does the difference in bias change from negative to positive but it also increases and decreases in a smooth transition depending on the target SOA (from noise onset), suggesting that the 1-back effect fluctuates rhythmically over time. All error bars indicate ± 1 MSE.

Oscillation of response bias but not sensitivity

We applied two different methods to examine response bias and sensitivity for rhythmic fluctuations. The first is the same *curve fitting* approach we used in our previous study (Ho et al., 2017). For this analysis, we pooled the individual trials across all participants, sorted the trials by target SOA and grouped them into hundred 10-ms bins, from 0.2 to 1.2 s post noise onset. For each bin, we computed d' (Eq. 1) and c (Eq. 2) and fitted sine curves of different frequencies to the temporal sequences of d' and c .

We first considered the data as a whole (without dividing the trials contingent on the preceding stimuli) and searched for significant modulations within the range of 4-12 Hz in 0.1-Hz steps. This frequency range encompasses the theta and alpha band. Figures 3 A&B show the binned aggregate data as a function of target SOA from noise onset, expressed as both sensitivity (Fig. 3A) and criterion (Fig. 3B). The shaded yellow and green areas enveloping the lines represent ± 1 standard error ($N = 14$) of the group mean (SEM), computed by bootstrapping the individual trials 2,000 times and applying the same curve fitting analysis to the surrogate data. It is evident that while sensitivity shows no clear oscillation, the criterion does oscillate around the frequency of 9.4 Hz, as indicated by the grey curve in Fig. 3B. For comparison, we fitted a sinusoidal with the same frequency, 9.4 Hz (grey curve in Fig. 3A) to the sensitivity data.

We evaluated the goodness of each harmonic fit (4-12 Hz) using R^2 combined with a permutation procedure, which involved shuffling the individual responses 2,000 times and applying the same curve fitting analysis to the surrogate data. From the surrogate data, we constructed a distribution of *maximal* R^2 (across all tested frequencies) for sensitivity (Fig. 3D) and criterion (Fig. 3E), against which we compared the R^2 of the original data. The results are summarised in Fig. 3C. For sensitivity (orange line), no frequency produced a good fit, none near the corrected 95% confidence threshold of $R^2 = 0.12$ (dotted black line).

However, the criterion data (green line) showed significant modulations between 9.2 and 9.6 Hz, with a strong peak at 9.4 Hz ($R^2 = 0.15$). The significance test (illustrated in Fig. 3D&E) confirmed that this frequency, 9.4 Hz, was significant for criterion, $p = 0.009$ (Fig. 3D), but not sensitivity, $p = 0.9$ (Fig. 3D). The phase of the 9.4 Hz oscillation at trial onset relative to the noise burst onset was $179^\circ \pm 16^\circ$ SD (by bootstrap).




Figure 3. Results of the curve fitting analysis. **(A)** The yellow line shows the time course of the detrended sensitivity (d') based on the aggregate data (by pooling all trials across the 14 participants and binned with a rectangular smoothing window of 20 ms moved every 10 ms). For display, we smoothed the data but conducted all statistical analyses on the non-smoothed data. The shaded area around the yellow line represents ± 1 SD of the bootstrapped data. The black curve depicts a 9.4 Hz oscillation fitted to the sensitivity data. **(B)** The analysis of the criterion (c) data (green line) followed the same binning and curve fitting procedure. The 9.4 Hz oscillation (black curve) fits the criterion better than the sensitivity data. **(C)** The goodness of the harmonic fit R^2 obtained by the procedure illustrated in A&B. The R^2 for the criterion (green line) is highest round 9.4 Hz, while it is low and constant for sensitivity (yellow line). The dotted black line represents the 95 percentiles of the permutation distribution depicted in E. **(D-E)** To obtain a significance evaluation of the goodness of the fit, we shuffled the aggregate data 2,000 times. After each shuffle, we fitted the data and extracted the maximal R^2 between 4-12 Hz illustrated in the yellow (sensitivity, D) and green distributions (criterion, E). The red lines indicate the R^2 yielded by the fit the original data at 9.4 Hz. The p -values reflect the proportion of permuted R^2 that is equal or greater than the original R^2 (red line) across all tested frequencies.

The results shown in Fig. 3 were based on an aggregate data analysis (i.e., by pooling all trials across subjects). In principle, the criterion oscillation in Fig. 3B could derive from a few subjects with very strong oscillatory effects. To verify that this is not the case, we examined the individual data and evaluated their coherence as a group using the *linear regression analysis* in Eq. 5 (illustrated in Fig. 1C). As detailed in the Methods, this approach does not require data binning. We applied the regression analysis to the individual *accuracy* (correct or incorrect) and *response* data ('left' or 'right').

The results, summarised in Fig. 4, corroborate those of the aggregate data analysis. Figure 4A plots the amplitude spectrum for *accuracy* (an approximation of sensitivity) and Fig. 4B that for *response bias* (an approximation of criterion), which were computed using the individual estimates of the *fixed-effect regression parameters*, β_1 and β_2 , and their vectorial mean (Eq. 6). The shaded yellow and green areas enveloping the lines represent ± 1 SEM. Response bias (Fig. 4B) shows a strong peak around the same frequency, 9.4 Hz, as the R^2 results for criterion in Fig. 3C. Using a similar permutation procedure as in the aggregate analysis, we evaluated the significance of each frequency between 4-12 Hz in 0.1-Hz steps. The corrected p -values are plotted in Fig. 4D&E for accuracy and response bias, respectively. The results for response bias shows that the oscillation at 9.4 Hz is significant ($p = 0.027$) after correction for multiple comparisons with maximal statistics (illustrated in Fig. 4F). The amplitude at 9.4 Hz is $A = 0.02 \pm 0.01$ SEM. While there are several peaks in the amplitude spectrum for accuracy (Fig. 4A), none was significant after multiple comparison correction (Fig. 4D).

Figure 4F illustrates how the p -values were computed and corrected. The green dots (2,000 in total) clustering in a ring around the origin (0, 0) represent the joined distribution of *maximal* B_1 and B_2 (group averages of the individual β_1 and β_2) obtained by permutation. To construct this distribution, we determined the maximal vector of each randomisation,

irrespective of the frequency. The red dot shows the group mean estimated from the original data at the peak frequency, 9.4 Hz. The red circle has a radius equal to the distance of the red dot (B_1, B_2) to the origin (0, 0). The p -value reflects the proportion of permuted data exceeding the original data (the green points that fall outside the circle).

Finally, Fig. 4C shows the individual phase and amplitude vectors for response bias at 9.4 Hz. The length of the vectors reflects the amplitude of the 9.4 Hz oscillation and the vector direction indicate individual phases at noise onset. The vectors are tightly clustered around a phase angle, θ , of $172^\circ \pm 16^\circ$ SEM (Eq. 7). This is consistent with the phase angle we obtained from the curve fitting analysis with the aggregate data (see above).




Figure 4. Results of the linear regression analysis based on individual data and single trials. (A) The yellow line represents the amplitude spectrum for accuracy computed from the vectorial average of the individual estimates of β_1 and β_2 (see Eq. 7). The shaded area around the line indicates ± 1 SEM. (B) Amplitude spectrum of response bias based on the same analyses as for accuracy. (C) Individual 2D vectors (β_1, β_2) at 9.4 Hz for response bias. The length of the line indicates the amplitude and the direction of the line the phase at noise onset (i.e., phase reset). (D) The results of the 2D permutation test for accuracy for the frequency range of interest, 4-12 Hz. We corrected for multiple comparisons using maximal statistics illustrated in F. (E) Corrected p -values for response bias obtained by the same 2D

permutation as for accuracy. The dotted black line indicates $\alpha = 0.05$. **(F)** Illustration of the 2D permutation test by which the p -values in D&E were computed. They reflect the proportion of maximal permutation vectors that exceed the group mean (red dot), i.e., the green points that fall outside the red circle.

Oscillation in response bias is driven by stimulus history

Having established the existence of rhythmic fluctuations in bias and criterion in both the aggregate and individual data, we investigated how the oscillations relate to serial effects driven by expectations from stimuli of previous trials using two different analysis. In the first analysis, we separated the trials based on whether the previous stimulus had been presented to the same ear (congruent) or a different ear (incongruent), and tested both sets of data for oscillations in both sensitivity and bias, using the same curve fitting and linear regression analyses as above. As predicted, the oscillation observed in criterion (Figs. 3&4) was driven by the preceding stimulus history, occurring only when the previous stimulus was *congruent* with the current stimulus.

Figure 5 shows the results of the analysis of the aggregate data. As before, we fitted the temporal sequences of sensitivity (d') and criterion (c) for congruent (Fig. 5A) and incongruent trials (Fig. 5C) with sinusoids ranging in frequency between 4-12 Hz in 0.1 Hz steps. Congruent trials (dark green line), but not incongruent trials (light green line), showed a good fit at 9.4 Hz (thick grey line). The goodness of fit at all tested frequencies are plotted in Fig. 5D. The largest R^2 was obtained around 9.4 Hz for congruent trials, with $R^2 = 0.15$, while for incongruent trials, the goodness of fit did not approach significance at any frequency. As in the previous analysis, we created a distribution of maximal R^2 (Fig. 5D) from the 2,000 surrogate datasets obtained by permutation and determined the 95 percentile for both congruent (Fig. 5D) and incongruent (Fig. 5E) trials. Any frequency with R^2 exceeding this threshold ($R^2 = 0.12$, dotted black line in Fig. 5B), was considered significant. Only the peak (green line, congruent trials) around 9.4 Hz survived the multiple comparison

correction, with $p = 0.01$. The phase of this 9.4-Hz oscillation for congruent trials was, at noise onset, $180^\circ \pm 17^\circ$ SD (by bootstrap). For incongruent trials, the phase at 9.4 Hz was $179^\circ \pm 34^\circ$ SD.




Figure 5. Results of the one-back analysis for criterion with aggregate data. (A) The dark green line shows the binned *congruent* trials. For display, we smoothed the data but conducted all statistical analyses on the non-smoothed data. The error bars indicate ± 1 SD obtained by bootstrapping the aggregate data 2,000 times. The thick grey line represents the 9.4-Hz oscillation which we fitted to the criterion data. The *incongruent* trials were submitted to the same binning, curve fitting and bootstrapping procedure; the results are summarised in panel (B). (C) The goodness of fit for congruent (dark green line) and incongruent trials (light green line) at all tested frequencies from 4-12 Hz in 0.1-Hz steps. The black dotted line indicates the 95 percentiles of the distribution of maximal R^2 obtained by permuting the individual trials. (D) Distribution of maximal R^2 for congruent trials. The vertical red line indicates the R^2 of the original data and the p -value reflects the proportion of maximal R^2 greater or equal the R^2 of the original data. (E) Distribution of maximal R^2 for incongruent trials, calculated as in panel (D).

As before, we examined the individual subject data using the linear model in Eq. 5, separately for congruent and incongruent trials. The amplitude spectra in Fig. 6A and 6D are based on the individual estimates of β_1 and β_2 averaged across the 14 participants (see Eq. 7). As for the aggregate data, the congruent trials (dark green line) yielded a large peak around

9.4 Hz with $A = 0.03 \pm 0.009$ SEM (Fig. 6A). At this frequency, the amplitude is reduced for the incongruent amplitude spectrum with $A = 0.02 \pm 0.01$ SEM (Fig. 6D). Inspection of the individual vectors shows a tight cluster around a mean phase angle (at noise onset) of $164^\circ \pm 13^\circ$ SEM for congruent trials (Fig. 6B), while incongruent trials had a slightly greater phase dispersion, with a mean phase angle (at noise onset) of $177^\circ \pm 18^\circ$ SEM (Fig. 6E).

The results of the 2D permutation test plotted in Fig. 6G corroborate the observations in Figs. 5A&D. The only frequencies to survive the strict multiple comparison correction were around 9.4 Hz (dark green line, congruent trials) with $p = 0.045$. In contrast, incongruent trials showed no significant frequencies (light green line). The 2D permutation distributions for congruent and incongruent trials at 9.4 Hz are plotted in Fig. 6C and 6F, respectively. The thick red dot shows the amplitude and phase of the group mean with respect to the permutation distribution. Only in the congruent case does the group mean exceed the distribution of maximal vectors significantly (Fig. 6C).

We also analysed the sensitivity for congruent and incongruent trials, but failed to find significant oscillations in either data set (data not shown). The lack of modulation in sensitivity in the present study is consistent with our previous findings that sensitivity oscillates in antiphase between the left and right ears (Ho et al., 2017): because they are out of phase, they should cancel each other under the conditions of this study, where results for left and right ears need to be combined for the sensitivity analysis. We checked this hypothesis by a post-hoc regression analysis of the accuracy of congruent trials separated by ear of origin. We first examined the phase coherence across subjects at all frequencies of interest (4-12 Hz) for each ear and congruence condition separately. Observers showed significant phase consistencies for *both* ears for the *congruent* trials at only one region in the frequency spectrum, from 9.1 to 9.7 Hz (consistent with the grey shaded region of significance in Fig. 6B). At no frequency was there strong phase coherence in the

incongruent trials. Using the Watson-Williams test (circular analogue to a two-sample t -test; Berens, 2009), we confirmed that the group phase distributions for left- and right-ear accuracy in the congruent trials were significantly different ($p < 0.05$, Bonferroni corrected) between 9.1 and 9.7 Hz, peaking at around 9.4 Hz. Figure 6H shows the individual vectors at 9.4 Hz for congruent trials containing a left target with a mean direction (thick red line) of $\sim 189^\circ \pm 12^\circ$ SEM. For congruent trials containing a right target, the individual vectors show a mean direction of $\sim 304^\circ \pm 14^\circ$ SEM at 9.4 Hz (Fig. 6I). Although the relationship between the left and right accuracy on congruent trials is not exactly antiphase ($\sim 115^\circ$), they do show a trend in this direction. This suggests that the absence of sensitivity oscillations (Figs. 2&3) in the present study is likely due to a cancellation of left and right ear oscillations.




Figure 6. Results of the one-back analysis for response bias with individual subject data. (A) Amplitude spectrum for the congruent trials computed based on the individual estimates of β_1 and β_2 averaged across participants. The shaded area around the dark green line indicates ± 1

SEM. By the same method, we computed the amplitude spectrum with incongruent trials; the results can be inspected in panel **(D)**. **(B)** Individual phase and amplitude vectors (at noise onset, i.e., phase reset) based on congruent trials at 9.4 Hz. The subjects are coded with the same colours as in Fig. 4C. The individual vectors for incongruent trials at 9.4 Hz are shown in **(E)**. **(C)** The joined distribution of maximal vectors for congruent trials. The p -value reflects the proportion of vectors that exceed the group mean (red dot); in practice, that is the number of green points falling outside the red circle. The joined distribution of maximal vectors for incongruent trials is shown in **(F)**. **(G)** The computations in (C&F) were done for every tested frequency between 4-12 Hz in 0.1-Hz steps. The dark and light green lines depict the corrected p -values for congruent and incongruent trials, respectively. The black dotted line indicates $\alpha = 0.05$ (corrected for multiple comparisons). **(H)** As a post-hoc test, we split the congruent trials further into trials that contained a left target and trials that contained a right target and tested their phase relationship using circular statistics. At 9.4 Hz observers showed very strong phase coherence for both ears. Here, we plot the individual phase and amplitude vectors for congruent trials containing a target in the left ear. The thick red line indicates the direction of the mean vector (with unit length) which is close to 180° . **(I)** The individual phase and amplitude vectors at 9.4 Hz for congruent trials containing a right target. The direction of the mean vector is $\sim 300^\circ$.

As the serial dependence analysis showed strong 2-back effects (Fig. 2B), we hypothesised that the 9-Hz oscillation could be long-lasting, spanning at least two trials, which is also suggested by recent findings (Chang et al., 2017). To investigate whether events two trials back show measurable oscillations near 9.4 Hz, we separated the *1-back incongruent data* (which show no significant oscillations in Fig. 5C) further on the basis of whether the stimulus two trials back was congruent or incongruent. Given that the *1-back incongruent data* showed no oscillation in previous analyses, any memory trace should be relatively weak and confined to a very narrow frequency window 0.5 Hz wide, 9.1-9.6 Hz, the frequency range where the congruent 1-back data showed a significant oscillation (after correction for multiple comparison; grey shaded box in Fig. 6B). We used the same analysis as before, but instead of testing every frequency from 4-12 Hz, we allowed the curve-fitting algorithm to search for the best frequency within our limited window of interest (9.1-9.6 Hz) for both the original and shuffled data. The results, shown in Fig. 7 confirm a 9-Hz oscillation in the 2-back congruent data which was best fitted by a sinusoidal with frequency of ~ 9.2 Hz,

$R^2 = 0.08$. The permutation test confirmed that this fit is significant (Fig. 7B), $p = 0.037$ when compared with the best fits of shuffled data within the frequency of interest, 9.1-9.6 Hz. There was no modulation near this frequency in the 2-back incongruent data (Figs 2E&F): $R^2 = 0.02$, with $p = 0.7$ (Fig. 7D). No other frequency within the range of 4-12 Hz approached significance, after multiple comparison correction.

To examine the individual data and group coherence at 9.2 Hz, we applied the same regression analysis as before (Eq. 5). Figure 7C shows the individual vectors at 9.2 Hz based on the 2-back congruent trials and Fig. 7G the individual vectors based on the incongruent trials. The individual phases in the congruent condition cluster around a similar phase, $190^\circ \pm 18^\circ$, similar to that of the 1-back congruent data (Fig. 6B). The mean phase angle in the incongruent condition is $295^\circ \pm 15^\circ$ and bears no relation to either the mean phase in the 1-back congruent or incongruent condition (Figs. 6B&E). Figures 7D&H show the results of the 2D permutation test at 9.2 Hz, which are consistent with the results of the aggregate data analysis. Very few points (dark cyan dots in Fig. 7D) from the permutation distribution ($p = 0.013$) exceed the group mean vector (thick red dot) in the congruent condition, while many more points exceed the mean vector in the incongruent condition, $p = 0.36$ (Fig. 7H). Taken together, the results suggest that the 9-Hz oscillation lasts at least two trials. Although the reverberation is weaker 2-back than 1-back, as to be expected, it may be sufficient to induce the long-lasting serial dependence observed on average (Fig. 2).




Figure 7. Results of the two-back analysis for criterion with aggregate data. For this analysis, we separated the *incongruent* data from the 1-back analysis (Figs. 5&6) further into congruent and incongruent 2-back. **(A)** The dark cyan line shows the binned *congruent* trials. As in Fig. 4, we smoothed the data for display but conducted all statistical analyses on the non-smoothed data. The error bars indicate ± 1 SD obtained by bootstrap ($n = 2,000$). The dark grey thick line represents the best fitting sinusoid which for the *congruent trials* was at 9.2 Hz (the frequency was free to vary between 9.1 and 9.6 Hz). The binned incongruent data is shown in (E), fitted with the same frequency, 9.2 Hz. No other frequency in the 9.1-9.6 Hz range fitted better. **(B)** The R^2 obtained at 9.2 Hz (thick red line) was compared against the goodness-of-fit of the shuffled data (dark cyan histogram), binned and fitted as the original data. The test confirmed that the 9.2-Hz oscillation in the congruent data was significant. As shown in (F), the same test for the incongruent condition was not significant. **(C)** The individual vectors in the congruent condition at 9.2 Hz. The subjects are colour-coded as in Figs. 4&6. Their phases cluster around a similar phase as in the congruent 1-back (Fig. 6B). In contrast, (F) shows the incongruent phase cluster at 9.2 Hz and their mean phase bears no relation to either the congruent or incongruent mean phase in the 1-back (Figs. 6B&D). **(D)** The result of the 2D permutation test for the congruent condition, which is consistent with the result of the aggregate data analysis shown in (B). The 2D permutation result for the incongruent condition is shown in (H) and also consistent with the result in (F).

Discussion

To maintain a stable and coherent percept in a world that is naturally noisy and ambiguous, observers take advantage of past information to anticipate forthcoming sensory input. While there is a good deal of behavioural evidence in favour of this predictive account of perception, little is known about the underlying neural mechanisms. Here, we present evidence suggesting that predictive perception is implemented rhythmically through alpha-band oscillations, along the lines of the “perceptual echo” suggested by VanRullen and Macdonald (2012).

Our current study shows that identifying the ear of origin of a weak tone was strongly biased by previous stimuli, two or even three trials before the current trial. Although the immediately previous trial had no average serial effect, we observed a strong rhythmic fluctuation in bias (measured by *criterion*) at ~9.4 Hz, which was critically dependent on stimulus history: strong oscillations occurred only when the previous target had been presented to the same ear as the current one, and weaker oscillations could be elicited by stimuli in the same ear two trials previously. The results reinforce our previous study (Ho et al., 2017) showing oscillations in criterion (for a different task) at similar frequencies (around 7.5 and 8.7 Hz), and extend these in an important way by showing that the oscillations are contingent on past stimulus history.

Why do oscillations occur only when the previous trial was presented to the same ear as the current one? One possible mechanism, illustrated in Fig. 8, is based on the “perceptual echo”, as suggested by VanRullen and Macdonald (2012; see also Bowen, 1989). An auditory signal presented to one ear should elicit a neural reverberatory response that oscillates in the alpha range, and affects subsequent perceptual decisions. Our data further suggest that the perceptual echo is confined to the ear in which it is evoked. We assume that

the phase of this oscillation becomes aligned to the onset of the noise-burst of the following trial (dotted vertical black lines in Fig. 8), and that the alignment phase is opposite for each ear (180° for the left and 0° for the right ear). This echo could then bias the response to subsequent signals presented to the same ear, either by modulating the neural gain of the signals (Rahnev et al., 2011), or by causing a shift in the decision boundary.

To illustrate, Fig. 8 shows putative internal representations of noise and signal. Presentation of a stimulus may set up an internal “echo” at about 9 Hz, confined to that ear. The onset of the noise burst resets the phase of the echo, to about 180° in the left ear (Fig. 8A) and 0° in the right ear (Fig. 8B). If we assume that the echo modulates gain, then the amplitude of the response to stimuli presented to the same ear will depend on time of presentation, more amplified if presented at a peak than at a trough. This modulation will be reflected in response bias, but not average sensitivity, as the echoes in the two ears are assumed to be out of phase (discussed below). The new stimulus will in turn elicit a new echo in that ear (red trace), which reverberates to the next trial.




Figure 8. Possible mechanisms by which past sensory experience biases forthcoming perceptual decisions, causing oscillations in congruent but not incongruent trials. **(A)** The green oscillation represents the ‘echo’ evoked by a target in the left ear in the previous trial ($n-1$). The reverberation resonates at alpha rhythm and continues to the current trial (n) where

it synchronises with other ongoing brain oscillations (possibly related to perceptual and decisional processes) by the noise onset. In this example, the current target is *congruent* with the preceding target, i.e., both occur in the left ear. Because the echo is ear specific, it will bias the observer's decision towards a "left response" when the stimulus is presented in the excitatory period of the echo and toward a "right response" in the negative periods. **(B)** Here, the target is *incongruent* with the preceding target, i.e., it occurs in the opposite ear. In this case, the echo has no influence on the observer's decision. In both scenarios, the target of the current trial will elicit a new echo (red oscillation) that reverberates into the next trial.

How could cyclic increases in sensitivity or gain result in changes in *criterion* but not in d' ? As the sequence of stimuli was completely random, past information was not informative about the current trial, and therefore could not increase d' , either on average or in a cyclical manner; but it can change the trial-by-trial judgment of which ear carried the weak signal. It is important to point out that *criterion* changes do not necessarily reflect decision processes (such as shifts in decision boundaries), but can also reflect perceptual changes (Peters et al., 2016; Witt et al., 2015). In the current experimental design, calculations of d' can only be made after combination of hits and false alarms to stimuli presented to the two ears. If the modulation of activity in the two ears is in counterphase, they will cancel each other out when the output is combined for d' , but add together for *criterion*. Evidence from our previous (Ho et al., 2017) and present study (of Figs. 6H&I) suggests that this does occur: when the congruent data are separated for ear of origin of the signal, the oscillations in the left ear tend to be out of phase with those of the right ($\sim 115^\circ$ on average in this study).

Why did our experiments reveal no positive serial dependence on the immediately previous trial (1-back), as is normally observed in studies of serial dependence? There are (at least) two possible, non-mutually exclusive explanations. The first is a consequence of the forced-choice paradigm used here (while most studies of serial dependence use a reproduction technique). Forced-choice paradigms can lead to sequential response biases independent of the stimuli, such as alternation of responses (Abrahamyan et al., 2016), especially after a long run of similar responses (probably related to the gambler's fallacy

(Burns and Corpus, 2004; Tversky and Kahneman, 1971). As the responses were strongly correlated with stimuli (75% correct), a systematic alternation would tend to cancel out positive serial dependence based on the previous stimulus (but have no effect on average for trials two-back). Not all observers alternate: some show the opposite tendency, “sticking” with the current response. Figure 2C is consistent with this idea: while only one observer showed negative serial dependence for 2-back trials, the variability in 1-back trials was much higher: five out of 14 observers showed positive 1-back biases, showing a tendency for “sticking”, while the others could be switchers. Further evidence comes from the response-based analysis, which showed a strong negative serial dependency, meaning that responses tended to be *different* from the previous one (independent of the stimuli). As stimuli and responses were highly correlated, this response-dependent bias will also impact the dependence on previous stimuli, potentially cancelling any positive serial dependence that may have been present.

Another possible reason for the lack of positive 1-back effects is that the stimuli may have caused both positive serial dependence and negative adaptation aftereffects. As mentioned in the introduction, both types of effects have been reported in sequential judgments, sometimes within the same experiment (Abrahamyan et al., 2016; Chopin and Mamassian, 2012; Taubert et al., 2016). Negative aftereffects tend to be shorter lived than the positive dependencies, and should therefore affect only 1-back, not 2-back trials (Chopin and Mamassian, 2012). This seems to be less plausible given the strong dependence on response (while aftereffects are stimulus-driven), but we cannot rule out the possibility.

Whatever the reason for the lack of positive serial dependence in the averaged results, our study suggests that oscillations may be a more sensitive signature of memory-based perceptual effects than simply looking at average results. Many competing effects could

reduce or annul average serial dependence effects, without affecting rhythmic, time-dependent oscillations.

As both vision and audition have similar serial dependencies and oscillations of decision criteria at alpha rhythm, the mechanism described in Fig. 8 may generalise to other sensory modalities. However, considering the anatomical and organisational differences between the visual and auditory system, it is possible that this mechanism had to be adapted to the specific architecture of each sense. For instance, oscillatory effects that are robust in vision are not readily observed in audition (VanRullen et al., 2014; Zoefel et al., 2015; Zoefel and Heil, 2013; Zoefel and VanRullen, 2017). However, by using *dichotic* rather than *diotic* stimulation (as all previous studies had done) we were able to show that similar to visual sensitivity, auditory sensitivity also oscillates but in antiphase for the two ears (Ho et al., 2017). As a result, when we summed the ears (as would be the case in diotic stimulations), the sensitivity oscillations cancelled each other out. Our findings therefore show that the ears need to be stimulated separately in order to observe certain oscillatory effects.

In summary, we showed that when participants were asked to determine the ear of origin of a brief sinusoidal target masked by white noise, their decision criteria oscillated at an alpha rhythm. Further analyses revealed that the alpha oscillation in criteria was driven entirely by trials in which the target occurred in the same ear as that of the immediately preceding trial. When the previous and current targets occurred in opposite ears, we found no observable oscillations of criteria. To account for these findings, we proposed that every target elicits a long-lasting reverberation at alpha rhythm that is ear-specific and continues to the next trial, where it synchronises with other ongoing brain oscillations possibly related to both perceptual and decisional processes. Because this ‘echo’ is ear specific, it will only bias perceptual decisions (towards or away from the ear in which the echo was elicited), if the following target occurs in the same ear. The bias is not absolute but fluctuates over time, and

may become stronger as more evidence of the same stimuli occurs successively in the sequence. Although this model is very basic, we believe it could be elaborated to explain more complex serial dependence effects and provide a deeper understanding of how expectation or prediction guides perception.

Materials and Methods

Participants

Eighteen healthy participants took part in the experiment. All reported normal hearing. Four participants were male and two left-handed. After inspecting participants' auditory thresholds and (log-transformed) reaction times, we excluded four participants (one male) from the data analysis for the following reasons. Three exhibited large differences in mean auditory threshold between the left and right ear (1 standard deviation from the group mean) and one displayed an atypical distribution of very long reaction times (2.5 standard deviations from the group mean). The mean age of the remaining 14 participants was 21.14 ± 4.22 . All participants provided written, informed consent. The study was approved by the Human Research Ethics Committees of the University of Sydney. No power analysis was used to decide the number of subjects or the number of trials per subject. We based our sample size estimations on our previous study (Ho et al., 2017), which showed oscillations in auditory perceptual performance before, and other studies on similar behavioural rhythms in vision (Fiebelkorn et al., 2013; Landau and Fries, 2012).

Experimental procedure

The experimental design was kept largely the same to our earlier study (Ho et al., 2017). However, instead of the bilateral tone discrimination task (2×2 design) used in our previous

experiment, we asked participants to determine the *ear of origin* of a similarly brief sinusoidal tone of 10-ms duration and 1,000 Hz in frequency masked by dichotic white noise.

As in the previous experiment, participants sat in a dark room and listened to auditory stimuli via in-ear tube-phones (ER-2, Etymotic Research, Elk Grove, Illinois). They wore earmuffs on top of the tube-phones (3M Peltor 30 dBA) to isolate external noise. The broadband white-noise masks were 2 s long and randomly generated each trial. To ensure that the noise masks were both clearly lateralised and uncorrelated, the left- and right-ear maskers were in antiphase (each a time-reversed duplicate of the other). As illustrated in Fig. 1A, the noise segments were presented to both ears with simultaneous onset. The target tone was delivered randomly with equal probability in either ear during the 2-s noise masker with a stimulus-onset asynchrony (SOA) randomly drawn from an interval of 0.2–1.2 s post noise onset. Participants responded via button press on a response box (ResponsePixx, Vpixmap Technologies, Saint-Bruno, Quebec). All participants used their thumbs to respond were instructed to respond as soon as possible while the noise masker was still present. For each ear, the target's intensity was kept around individuals' thresholds (75 % accuracy) using an accelerated stochastic approximation staircase procedure (Faes et al., 2007; García-Pérez, 2011). The next trial started after a silent inter-trial interval (ITI) of random duration between 1.2–2.2 s.

Participants completed 2,800 trials (40 blocks of 70 trials) in total. Blocks lasted ~5 minutes and participants were allowed to rest after every block. Prior to the experiment, they completed a practice block of 20 trials with feedback but no feedback was provided during the experiment. Stimuli were presented using the software *PsychToolbox* (Brainard, 1997) in conjunction with *DataPixx* (Vpixmap Technologies, Saint-Bruno, Quebec) in MATLAB (Mathworks, Natick, Massachusetts).

Signal detection theory

Prior to the data analysis, trials in which the response occurred before the target onset or after the noise offset were excluded. Additionally, we eliminated trials with reaction time (RT) exceeding the 99 % confidence intervals of individuals' RT and with target intensity beyond the 95 % confidence intervals of individuals' thresholds.

We computed d' and c using Eq. 1 and 2 from *signal detection theory* (SDT) (Green and Swets, 1966; Macmillan and Creelman, 2004). As illustrated in Fig. 1B, the calculation of the hit rate (H_{right}) was based on the hits from the right target condition and the false alarm rate (FA_{left}) based on the false alarms from the left target condition. d' is given by the difference between z-transformed hit and false alarm rates:

$$d' = \frac{z(H_{right}) - z(FA_{left})}{\sqrt{2}} \quad (1)$$

Half the sum of the z-transformed hit and false alarm rates gives c , the bias towards one of the responses (left ear when positive and right ear when negative).

$$c = -0.5 \times (z(H_{right}) + z(FA_{left})) \quad (2)$$

Behavioural oscillations

Aggregate data analysis. To look for oscillations in sensitivity and response bias, we applied two different methods. The first is the same *curve fitting* approach we used in our previous study (Ho et al., 2017). For this analysis, we pooled across all 14 participants, sorted the trials by target SOA and grouped the data into hundred 10-ms bins, from 0.2 to 1.2 s post noise onset. The mean number of trials per bin was 151 ± 24 for the left-target condition and 152 ± 25 for the right-target condition. For each bin, we computed d' and c as above (Eq. 1&2). Using the standard MATLAB function *fit* included in the *Curve Fitting* toolbox, we

fitted the following Fourier series model (Eq. 3) to the resulting sequence of sensitivity and criterion (see Fig. 3A&B):

$$f(t) = a_0 + a_1 \sin(\omega t) + b_1 \cos(\omega t) = A \cos(\omega t + \phi), \quad (3)$$

where t is time ($t = 0.2, 0.3 \dots 1.2$ in seconds), ω is the angular frequency ($\omega = 2\pi f$) we want to test, a_0 is a constant and a_1 and b_1 are the sine and cosine coefficients respectively. A and ϕ represent amplitude and phase of the sinusoidal fit. We used a non-linear least-squares method (a standard implementation in MATLAB) whereby the summed squares of the residuals were minimised through successive iterations (400 iterations in total). As in our previous study (Ho et al., 2017), sensitivity displayed a decreasing non-linear trend over time which we removed before curve fitting using the same polynomial fit (*polyfit* in MATLAB). Detrending was not applied to the criterion curve. For each frequency between 4 and 12 Hz (in 0.1 Hz steps), we fitted the best sinusoid, allowing amplitude and phase to vary (two degrees of freedom). This yielded a measure of goodness of fit, R^2 , as a function of frequency (Fig. 3C). To test the significance of every fit, we applied a permutation procedure (Ernst, 2004) whereby we shuffled the responses of each individual trial over all SOAs to generate 2,000 surrogate datasets which we submitted to the same binning and curve fitting procedure as the original data. To correct for multiple comparisons, we determined the maximal R^2 for every surrogate dataset irrespective of frequency. This resulted in a distribution of 2,000 maximal R^2 (Figs. 3D&E) against which we compared each fit to the original dataset. Any frequency that exceeded the 95 percentile of the maximal- R^2 distribution (dotted line in Fig. 3C) was considered significant. We also estimated the variability in the aggregate data by applying the bootstrap method which involves the random selection of the same number of trials (with replacement) as in the original data and submitting the surrogate datasets to the same binning and curve fitting procedure as above.

Individual and group analysis. The curve fitting method described above requires a sufficient number of trials per time point to accurately estimate the oscillations underlying d' and c . For that reason, we had to pool across all subjects and bin the aggregated data. In order to examine the individual data for oscillations and evaluate their coherence across subjects, we used a different approach which does not require data binning but allows for an estimation of participants' sensitivity and response bias based on single trials (for similar approaches, see (Benedetto et al., 2018; Tomassini et al., 2017)). As illustrated in Fig. 1C, the response y_i ($i = 1, 2 \dots n$, where n is the total number of trials) to a target presented at time t_i (i.e., the interval from noise onset to target onset in seconds) can be modelled as the linear combination of harmonics at each tested (angular) frequency (compare with Eq. 3):

$$\hat{Y}_n = \beta_0 + \beta_1 \sin(\omega t_n) + \beta_2 \cos(\omega t_n), \quad (4)$$

where $\hat{Y}_n$ is the predicted responses and β_0 , β_1 and β_2 are fixed-effect regression parameters that can be estimated using the *linear* least-squares method implemented in MATLAB as the *fitlm* function from the *Statistics and Machine Learning* toolbox. To retain some consistency between the individual data and aggregate data analyses, we analysed sensitivity and response bias separately, that is, y_i reflected either individual accuracy or response bias. Therefore, the model we used is a special case of the *general linear model* (GLM) as it is restricted to one dependent variable (also called *multiple linear regression*). This model estimates the regression parameters adequately when the sampling rate is uniform across the time series. As this condition may not always be met, we applied the *full model* which includes a third independent regressor containing information about the stimulus:

$$\hat{Y}_n = \beta_0 + \beta_1 \sin(\omega t_n) + \beta_2 \cos(\omega t_n) + \beta_3 S(t_n), \quad (5)$$

where S is the stimulus at time t and takes the value -1 or $+1$ for left and right target respectively. We examined sensitivity based on participants' correct and incorrect responses

in which case $y_i = 1$ for correct and $y_i = 0$ for incorrect. To analyse response bias, y_i took the value 1 when participants made a ‘right’ response and 0 when they made a ‘left’ response.¹ To measure the group’s coherence in terms of both phase and amplitude at each frequency, we averaged the sine and cosine regression parameters across all participants, i.e., $B_1 = \frac{1}{n} \sum_i^n \beta_1$ and $B_2 = \frac{1}{n} \sum_i^n \beta_2$, where $n = 15$ and $i = 1, 2 \dots n$, and obtained an amplitude spectrum (Figs. 4A&B) by taking the vectorial average of the individual estimates given by the square root of their sum squared for every frequency:

$$A = \sqrt{B_1^2 + B_2^2} \quad (6)$$

We computed the mean phase θ by taking the arctangent of the ratio of the averaged sine to cosine regression parameter:

$$\theta = \tan^{-1} \frac{B_1}{B_2} \quad (7)$$

The significance of the model fit in Eq. 5 was evaluated using a two-dimensional (2D) permutation test. Specifically, we tested the null hypothesis that the individual response data contain no temporal structure and thus their time stamps should be interchangeable. To this end, we shuffled the SOAs (i.e., the interval between noise onset and target presentation), keeping the relationship between the response (‘left’ response or ‘right’ response) and stimulus (left target or right target) constant (Kennedy and in Statistics-Simulation, 1996; Winkler et al., 2014). The permutation was carried out at the individual subject level 2,000 times per dataset. Each surrogate dataset was fitted using the same model described in Eq. 5 and the resulting β_1 and β_2 were averaged across subjects for every frequency tested. This yielded a joined distribution of 2,000 surrogate means for each frequency from 4-12 Hz in

¹ Binary responses can be modelled using a *generalised* linear model with a logit link function. In an ideal scenario in which the observer has no internal noise, simulation results suggest that the *generalised* linear model indeed outperforms the linear model; in more realistic settings, however, the two models yield comparable results (Knoblauch and Maloney, 2008) which our own simulations corroborate (not reported here).

0.1-Hz steps. Similar to the *multiple comparison correction* we applied in the *aggregate data analysis*, we determined the *maximal vector* of each joined distribution, irrespective of the frequency. This resulted in a joined distribution of 2,000 maximal vectors, against which we compared the original mean (see Fig. 4F). This comparison entails determining the proportion of surrogate amplitude group means equal or exceeding the original group mean.

References

- Aagten-Murphy D, Burr D. 2016. Adaptation to numerosity requires only brief exposures, and is determined by number of events, not exposure duration. *Journal of Vision* **16**:22–22. doi:10.1167/16.10.22
- Abrahamyan A, Silva L, Dakin SC, Carandini M, Gardner JL. 2016. Adaptable history biases in human perceptual decisions. *Proceedings of the National Academy of Sciences* **113**:E3548–E3557. doi:10.1073/pnas.1518786113
- Addams R. 1834. An account of a peculiar optical phænomenon seen after having looked at a moving body. *Philosophical Mag Ser 3* **5**:373–374. doi:10.1080/14786443408648481
- Alais D, Leung J, der Burg E. 2017. Linear Summation of Repulsive and Attractive Serial Dependencies: Orientation and Motion Dependencies Sum in Motion Perception. *The Journal of Neuroscience* **37**:4381–4390. doi:10.1523/JNEUROSCI.4601-15.2017
- Alais D, Orchard-Mills E, der Burg E. 2015. Auditory frequency perception adapts rapidly to the immediate past. *Attention, Perception, & Psychophysics* **77**:896–906. doi:10.3758/s13414-014-0812-2
- Alexi J, Cleary D, Dommissie K, Palermo R, Kloth N, Burr D, Bell J. 2018. Past visual experiences weigh in on body size estimation. *Scientific Reports* **8**:215. doi:10.1038/s41598-017-18418-3
- Arzounian D, de Kerangal M, de Cheveigné A. 2017. Sequential dependencies in pitch judgments. *The Journal of the Acoustical Society of America* **142**:3047–3057. doi:10.1121/1.5009938
- Benedetto A, Burr D, Morrone M. 2018. Perceptual Oscillation of Audio-Visual Time Simultaneity. *eNeuro* **5**:ENEURO.0047-18.2018. doi:10.1523/ENEURO.0047-18.2018
- Benedetto A, Morrone M. 2017. Saccadic Suppression Is Embedded Within Extended Oscillatory Modulation of Sensitivity. *The Journal of Neuroscience* **37**:3661–3670. doi:10.1523/JNEUROSCI.2390-16.2016
- Benedetto A, Spinelli D, Morrone CM. 2016. Rhythmic modulation of visual contrast discrimination triggered by action. *Proc R Soc Lond, B, Biol Sci* **283**:20160692.

- 1 doi:10.1098/rspb.2016.0692
- 2
- 3 Berens P. 2009. CircStat: a MATLAB toolbox for circular statistics. *J Stat Softw* **31**:1–21.
- 4
- 5 Bestelmeyer P, Rouger J, DeBruine LM, Belin P. 2010. Auditory adaptation in vocal affect
- 6 perception. *Cognition* **117**:217–223. doi:10.1016/j.cognition.2010.08.008
- 7
- 8 Bowen R. 1989. Two pulses seen as three flashes: a superposition analysis. *Vision research*
- 9 **29**:409–17.
- 10
- 11 Brainard DH. 1997. The Psychophysics Toolbox. *Spat Vis* **10**:433–436.
- 12 doi:10.1163/156856897X00357
- 13
- 14 Burns BD, Corpus B. 2004. Randomness and inductions from streaks: “Gambler’s fallacy”
- 15 versus ”hot hand“. *Psychonomic Bulletin & Review* **11**:179–184. doi:10.3758/BF03206480
- 16
- 17 Burr D, Ross J. 2008. A Visual Sense of Number. *Current Biology* **18**:425–428.
- 18 doi:10.1016/j.cub.2008.02.052
- 19
- 20 Buzsáki G, Draguhn A. 2004. Neuronal oscillations in cortical networks. *Science (New York,*
- 21 *NY)* **304**:1926–9. doi:10.1126/science.1099745
- 22
- 23 Chambers C, Pressnitzer D. 2014. Perceptual hysteresis in the judgment of auditory pitch
- 24 shift. *Attention, Perception, & Psychophysics* **76**:1271–1279. doi:10.3758/s13414-014-0676-
- 25 5
- 26
- 27 Chang AY, Schwartzman DJ, VanRullen R, Kanai R, Seth AK. 2017. Visual Perceptual Echo
- 28 Reflects Learning of Regularities in Rapid Luminance Sequences. *The Journal of*
- 29 *neuroscience : the official journal of the Society for Neuroscience* **37**:8486–8497.
- 30 doi:10.1523/JNEUROSCI.3714-16.2017
- 31
- 32 Chopin A, Mamassian P. 2012. Predictive Properties of Visual Adaptation. *Current Biology*
- 33 **22**:622–626. doi:10.1016/j.cub.2012.02.021
- 34
- 35 Cicchini G, Anobile G, Burr D. 2014. Compressive mapping of number to space reflects
- 36 dynamic encoding mechanisms, not static logarithmic transform. *Proc National Acad Sci*
- 37 **111**:7867–7872. doi:10.1073/pnas.1402785111
- 38
- 39 Cicchini G, Mikellidou K, Burr DC. 2018. The functional role of serial dependence. *Proc R*
- 40 *Soc B* **285**:20181722. doi:10.1098/rspb.2018.1722
- 41
- 42 Craddock M, Poliakoff E, El-dereby W, Klepousniotou E, Lloyd DM. 2017. Pre-stimulus
- 43 alpha oscillations over somatosensory cortex predict tactile misperceptions.
- 44 *Neuropsychologia* **96**:9–18. doi:10.1016/j.neuropsychologia.2016.12.030
- 45
- 46 de Lange FP, Rahnev DA, Donner TH, Lau H. 2013. Prestimulus Oscillatory Activity over
- 47 Motor Cortex Reflects Perceptual Expectations. *The Journal of Neuroscience* **33**:1400–1410.
- 48 doi:10.1523/JNEUROSCI.1094-12.2013
- 49
- 50 Dong DW, Atick JJ. 1995. Statistics of natural time-varying images. *Network: Computation*

- 1 *in Neural Systems* **6**:345–358. doi:10.1088/0954-898X_6_3_003
- 2
- 3 Engel A, Fries P, Singer W. 2001. Dynamic predictions: oscillations and synchrony in top-
- 4 down processing. *Nature reviews Neuroscience* **2**:704–16. doi:10.1038/35094565
- 5
- 6 Ernst MD. 2004. Permutation methods: a basis for exact inference. *Statistical Science*
- 7 **19**:676–685. doi:10.1214/088342304000000396
- 8
- 9 Faes L, Nollo G, Ravelli F, Ricci L, Vescovi M, Turatto M, Pavani F, Antolini R. 2007.
- 10 Small-sample characterization of stochastic approximation staircases in forced-choice
- 11 adaptive threshold estimation. *Percept Psychophys* **69**:254–62.
- 12
- 13 Fiebelkorn IC, Foxe JJ, Butler JS, Mercier MR, Snyder AC, Molholm S. 2011. Ready, Set,
- 14 Reset: Stimulus-Locked Periodicity in Behavioral Performance Demonstrates the
- 15 Consequences of Cross-Sensory Phase Reset. *The Journal of Neuroscience* **31**:9971–9981.
- 16 doi:10.1523/JNEUROSCI.1338-11.2011
- 17
- 18 Fiebelkorn IC, Saalman YB, Kastner S. 2013. Rhythmic Sampling within and between
- 19 Objects despite Sustained Attention at a Cued Location. *Curr Biol* **23**:2553–2558.
- 20 doi:10.1016/j.cub.2013.10.063
- 21
- 22 Fischer J, Whitney D. 2014. Serial dependence in visual perception. *Nat Neurosci* **17**:738–
- 23 743. doi:10.1038/nn.3689
- 24
- 25 Friston K. 2005. A theory of cortical responses. *Philosophical transactions of the Royal*
- 26 *Society of London Series B, Biological sciences* **360**:815–36. doi:10.1098/rstb.2005.1622
- 27
- 28 Friston KJ, Bastos AM, Pinotsis D, Litvak V. 2015. LFP and oscillations—what do they tell
- 29 us? *Curr Opin Neurobiol* **31**:1–6. doi:10.1016/j.conb.2014.05.004
- 30
- 31 García-Pérez MA. 2011. A cautionary note on the use of the adaptive up-down method. *J*
- 32 *Acoust Soc Am* **130**:2098–107. doi:10.1121/1.3628334
- 33
- 34 Gibson J, Radner M. 1937. Adaptation, after-effect and contrast in the perception of tilted
- 35 lines. I. Quantitative studies. *Journal of Experimental Psychology* **20**:453/467.
- 36
- 37 Grantham WD, Wightman FL. 1979. Auditory motion aftereffects. *Perception &*
- 38 *Psychophysics* **26**:403–408. doi:10.3758/BF03204166
- 39
- 40 Gray C, König P, Engel A, Singer W. 1989. Oscillatory responses in cat visual cortex exhibit
- 41 inter-columnar synchronization which reflects global stimulus properties. *Nature* **338**:334–7.
- 42 doi:10.1038/338334a0
- 43
- 44 Green DM, Swets JA. 1966. Signal Detection Theory and Psychophysics, John Wiley. John
- 45 Wiley.
- 46
- 47 Gregory R. 1997. Knowledge in perception and illusion. *Philosophical transactions of the*
- 48 *Royal Society of London Series B, Biological sciences* **352**:1121–7.
- 49 doi:10.1098/rstb.1997.0095
- 50

- 1 Haegens S, Vázquez Y, Zainos A, Alvarez M, Jensen O, Romo R. 2014. Thalamocortical
2 rhythms during a vibrotactile detection task. *Proceedings of the National Academy of*
3 *Sciences* **111**:E1797–E1805. doi:10.1073/pnas.1405516111
4
- 5 Helmholtz H. 1867. Handbuch der physiologischen Optik, Voss. Voss.
6
- 7 Ho HT, Leung J, Burr DC, Alais D, Morrone M. 2017. Auditory Sensitivity and Decision
8 Criteria Oscillate at Different Frequencies Separately for the Two Ears. *Current Biology*.
9 doi:10.1016/j.cub.2017.10.017
10
- 11 Huang Q, Jia J, Han Q, Luo H. 2018. Fast-backward replay of sequentially memorized items
12 in humans. *eLife* **7**:e35164. doi:10.7554/eLife.35164
13
- 14 Iemi L, Busch NA. 2018. Moment-to-Moment Fluctuations in Neuronal Excitability Bias
15 Subjective Perception Rather than Strategic Decision-Making. *eNeuro* **5**.
16 doi:10.1523/ENEURO.0430-17.2018
17
- 18 Iemi L, Chaumon M, Crouzet SM, Busch NA. 2017. Spontaneous Neural Oscillations Bias
19 Perception by Modulating Baseline Excitability. *J Neurosci* **37**:807–819.
20 doi:10.1523/JNEUROSCI.1432-16.2017
21
- 22 Kanai R, Verstraten F. 2005. Perceptual manifestations of fast neural plasticity: Motion
23 priming, rapid motion aftereffect and perceptual sensitization. *Vision Research* **45**:3109–
24 3116. doi:10.1016/j.visres.2005.05.014
25
- 26 Kashino M, Nishida S. 1998. Adaptation in the processing of interaural time differences
27 revealed by the auditory localization aftereffect. *The Journal of the Acoustical Society of*
28 *America* **103**:3597–604. doi:10.1121/1.423064
29
- 30 Kennedy P, in Statistics-Simulation CB. 1996. Randomization tests for multiple regression.
31 doi:10.1080/03610919608813350
32
- 33 Klimesch W, ppelmayr, Russegger H, Pachinger T, Schwaiger J. 1998. Induced alpha band
34 power changes in the human EEG and attention. *Neuroscience Letters* **244**:73–76.
35 doi:10.1016/S0304-3940(98)00122-0
36
- 37 Knoblauch K, Maloney LT. 2008. Estimating classification images with generalized linear
38 and additive models. *Journal of Vision* **8**:10–10. doi:10.1167/8.16.10
39
- 40 Landau A, Fries P. 2012. Attention Samples Stimuli Rhythmically. *Curr Biol* **22**:1000–1004.
41 doi:10.1016/j.cub.2012.03.054
42
- 43 Lee TS, Mumford D. 2003. Hierarchical Bayesian inference in the visual cortex. *Journal of*
44 *the Optical Society of America A, Optics, image science, and vision* **20**:1434–48.
45
- 46 Leopold DA, O’Toole AJ, Vetter T, Blanz V. 2001. Prototype-referenced shape encoding
47 revealed by high-level aftereffects. *Nature Neuroscience* **4**:nn0101_89. doi:10.1038/82947
48
- 49 Liberman A, Fischer J, Whitney D. 2014. Serial Dependence in the Perception of Faces.
50 *Current Biology* **24**:2569–2574. doi:10.1016/j.cub.2014.09.025

- 1
- 2 Limbach K, Corballis PM. 2016. Prestimulus alpha power influences response criterion in a
- 3 detection task. *Psychophysiology* **53**:1154–1164. doi:10.1111/psyp.12666
- 4
- 5 Macmillan NA, Creelman DC. 2004. Detection theory: A user's guide, Lawrence Erlbaum
- 6 Associates. Lawrence Erlbaum Associates.
- 7
- 8 Mayer A, hwiedrzik C, Wibral M, Singer W, Melloni L. 2016. Expecting to See a Letter:
- 9 Alpha Oscillations as Carriers of Top-Down Sensory Predictions. *Cerebral Cortex* **26**:3146–
- 10 3160. doi:10.1093/cercor/bhv146
- 11
- 12 Neitz J, Carroll J, Yamauchi Y, Neitz M, Williams DR. 2002. Color Perception Is Mediated
- 13 by a Plastic Neural Mechanism that Is Adjustable in Adults. *Neuron* **35**:783–792.
- 14 doi:10.1016/S0896-6273(02)00818-8
- 15
- 16 Pantle AJ, Gallogly DP, Piehler OC. 2000. Direction biasing by brief apparent motion
- 17 stimuli. *Vision Research* **40**:1979–1991. doi:10.1016/S0042-6989(00)00071-7
- 18
- 19 Peters MA, Ro T, Lau H. 2016. Who's afraid of response bias? *Neuroscience of*
- 20 *Consciousness* **2016**:niw001. doi:10.1093/nc/niw001
- 21
- 22 Premereur E, Vanduffel W, Janssen P. 2012. Local Field Potential Activity Associated with
- 23 Temporal Expectations in the Macaque Lateral Intraparietal Area. *Journal of Cognitive*
- 24 *Neuroscience* **24**:1314–1330. doi:10.1162/jocn_a_00221
- 25
- 26 Rahnev D, Maniscalco B, Graves T, Huang E, de Lange FP, Lau H. 2011. Attention induces
- 27 conservative subjective biases in visual perception. *Nature Neuroscience* **14**:1513.
- 28 doi:10.1038/nn.2948
- 29
- 30 Reinhardt-Rutland A, Anstis S. 1982. Auditory adaptation to gradual rise or fall in intensity
- 31 of a tone. *Perception & Psychophysics* **31**:63–67. doi:10.3758/BF03206201
- 32
- 33 Rohenkohl G, Nobre A. 2011. Alpha Oscillations Related to Anticipatory Attention Follow
- 34 Temporal Expectations. *J Neurosci* **31**:14076–14084. doi:10.1523/JNEUROSCI.3387-
- 35 11.2011
- 36
- 37 Sanders LL, Aukstulewicz R, Hohlefeld FU, Busch NA, Sterzer P. 2014. The influence of
- 38 spontaneous brain oscillations on apparent motion perception. *NeuroImage* **102 Pt 2**:241–8.
- 39 doi:10.1016/j.neuroimage.2014.07.065
- 40
- 41 Schweinberger SR, Casper C, Hauthal N, Kaufmann JM, Kawahara H, Kloth N, Robertson
- 42 D, Simpson AP, Zäske R. 2008. Auditory Adaptation in Voice Perception. *Current Biology*
- 43 **18**:684–688. doi:10.1016/j.cub.2008.04.015
- 44
- 45 Sherman MT, Kanai R, Seth AK, VanRullen R. 2016. Rhythmic Influence of Top-Down
- 46 Perceptual Priors in the Phase of Prestimulus Occipital Alpha Oscillations. *J Cogn Neurosci*
- 47 **28**:1–13. doi:10.1162/jocn_a_00973
- 48
- 49 Summerfield C, de Lange FP. 2014. Expectation in perceptual decision making: neural and
- 50 computational mechanisms. *Nature Reviews Neuroscience* **15**:nrn3838. doi:10.1038/nrn3838

- 1 Summerfield C, Koechlin E. 2008. A neural representation of prior information during
2 perceptual inference. *Neuron* **59**:336–47. doi:10.1016/j.neuron.2008.05.021
3
4
- 5 Taubert J, Alais D, Burr D. 2016. Different coding strategies for the perception of stable and
6 changeable facial attributes. *Scientific Reports* **6**:32239. doi:10.1038/srep32239
7
- 8 Thompson P, Burr D. 2009. Visual aftereffects. *Current Biology* **19**:R11–R14.
9 doi:10.1016/j.cub.2008.10.014
10
- 11 Tomassini A, Ambrogioni L, Medendorp PW, Maris E. 2017. Theta oscillations locked to
12 intended actions rhythmically modulate perception. *eLife* **6**:e25618. doi:10.7554/eLife.25618
13
- 14 Tomassini A, Spinelli D, Jacono M, Sandini G, Morrone M. 2015. Rhythmic Oscillations of
15 Visual Contrast Sensitivity Synchronized with Action. *J Neurosci* **35**:7019–7029.
16 doi:10.1523/jneurosci.4568-14.2015
17
- 18 Tversky A, Kahneman D. 1971. Belief in the law of small numbers. *Psychological Bulletin*
19 **76**:105. doi:10.1037/h0031322
20
- 21 VanRullen R. 2017. Predictive Coding and Neural Communication Delays Produce Alpha-
22 Band Oscillatory Impulse Response FunctionsConference on Cognitive Computational
23 Neuroscience .
24
- 25 VanRullen R, Macdonald J. 2012. Perceptual Echoes at 10 Hz in the Human Brain. *Current*
26 *Biology* **22**:995–999. doi:10.1016/j.cub.2012.03.050
27
- 28 VanRullen R, Zoefel B, Ilhan B. 2014. On the cyclic nature of perception in vision versus
29 audition. *Philos Trans R Soc Lond, B, Biol Sci* **369**:20130214. doi:10.1098/rstb.2013.0214
30
- 31 Voss R. 1975. “1/f noise” in music and speech. *Nature* **258**:317–318.
32
- 33 Voss R, Clarke J. 1978. “1/f noise” in music: Music from 1/f noise. *The Journal of the*
34 *Acoustical Society of America* **63**:258–263.
35
- 36 Winkler AM, Ridgway GR, Webster MA, Smith SM, Nichols TE. 2014. Permutation
37 inference for the general linear model. *NeuroImage* **92**:381–397.
38 doi:10.1016/j.neuroimage.2014.01.060
39
- 40 Witt JK, Taylor J, Sugovic M, Wixted JT. 2015. Signal Detection Measures Cannot
41 Distinguish Perceptual Biases from Response Biases. *Perception* **44**:289–300.
42 doi:10.1068/p7908
43
- 44 Yoshimoto S, Uchida-Ota M, Takeuchi T. 2014. The reference frame of visual motion
45 priming depends on underlying motion mechanisms. *Journal of Vision* **14**:10–10.
46 doi:10.1167/14.1.10
47
- 48 Yuille A, Kersten D. 2006. Vision as Bayesian inference: analysis by synthesis? *Trends in*
49 *Cognitive Sciences* **10**:301–308. doi:10.1016/j.tics.2006.05.002
50

1 Zoefel B, Heil P. 2013. Detection of Near-Threshold Sounds is Independent of EEG Phase in
2 Common Frequency Bands. *Frontiers in Psychology* **4**:262. doi:10.3389/fpsyg.2013.00262
3
4 Zoefel B, Pasham N, Brüers S, VanRullen R. 2015. The ability of the auditory system to cope
5 with temporal subsampling depends on the hierarchical level of processing. *NeuroReport*
6 **26**:773. doi:10.1097/WNR.0000000000000422
7
8 Zoefel B, VanRullen R. 2017. Oscillatory Mechanisms of Stimulus Processing and Selection
9 in the Visual and Auditory Systems: State-of-the-Art, Speculations and Suggestions. *Front*
10 *Neurosci* **11**:296. doi:10.3389/fnins.2017.00296
11