

Table S2. Secondary antibodies used in this study

Secondary antibody	Host species	Dilution	Supplier
Alexa Fluor 488 goat anti-mouse IgG (H+L)	Goat anti-mouse	1:500	Molecular Probes Europe BV
Alexa Fluor 488 goat anti-rabbit IgG (H+L)	Goat anti-rabbit	1:200	Molecular Probes Europe BV
Alexa Fluor 546 goat anti-mouse IgG (H+L)	Goat anti-mouse	1:500	Molecular Probes Europe BV
Alexa Fluor 568 goat anti-rabbit IgG (H+L)	Goat anti-rabbit	1:500	Molecular Probes Europe BV
Alexa Fluor 568 donkey anti-goat IgG (H+L)	Donkey anti-goat	1:500	Molecular Probes Europe BV
Alexa Fluor 546 goat anti-mouse IgG (H+L)	Goat anti-mouse	1:300	Molecular Probes Europe BV
Biotinylated anti-goat IgG (H+L)	Rabbit anti-goat	1:100	Vector Laboratories
Biotinylated anti-mouse IgG (H+L)	Goat anti-mouse	1:100	Vector Laboratories
Biotinylated anti-rabbit IgG (H+L)	Goat anti-rabbit	1:100	Vector Laboratories
Cy2-conjugated streptavidin	—	1:500	Amersham
Cy3-conjugated streptavidin	—	1:1000	Amersham