

SUPPLEMENTARY TABLE S2. DEMOGRAPHIC CHARACTERISTICS AND CANNABIS USE PATTERNS ACROSS AGE GROUPS

	<i>Young, %</i>	<i>Middle age, %</i>	<i>Older, %</i>	
Gender (male)	56.7	51.3	52.3	$\chi^2 = 7.82, p = 0.02$
Ethnicity (White)	82.2 ^a	88.2 ^b	89.4 ^b	$\chi^2 = 23.64, p < 0.001$
Cannabis user type				
Medical	17.8 ^a	37.3 ^b	52.1 ^c	$\chi^2 = 227.85, p < 0.001$
Recreational	49.8 ^a	33.9 ^b	17.8 ^c	$\chi^2 = 170.62, p < 0.001$
Mixed	32.3	28.8	30.1	$\chi^2 = 3.48, p = 0.17$
Education				
High school/GED or less	37.6 ^a	25.4 ^b	16.5 ^c	$\chi^2 = 90.92, p < 0.001$
Technical or associate degree	22.7 ^a	26.7 ^a	34.7 ^b	$\chi^2 = 27.17, p < 0.001$
Bachelor's degree	33.4 ^a	28.5 ^{a,b}	25.0 ^b	$\chi^2 = 14.16, p = 0.001$
Graduate degree	6.3 ^a	19.5 ^b	23.8 ^b	$\chi^2 = 128.21, p < 0.001$
Income				
<\$20,000	24.5 ^a	15.9 ^b	19.4 ^{a,b}	$\chi^2 = 26.30, p < 0.001$
\$20,000–60,000	42.4	40.0	35.9	$\chi^2 = 6.30, p = 0.04$
\$60,000–80,000	9.0 ^a	13.8 ^b	12.7 ^{a,b}	$\chi^2 = 13.80, p = 0.001$
>\$80,000	24.0 ^a	30.2 ^b	32.0 ^b	$\chi^2 = 16.39, p < 0.001$
Employment				
Full-time	51.7 ^a	64.9 ^b	33.1 ^c	$\chi^2 = 139.40, p < 0.001$
Part-time	27.8 ^a	14.4 ^b	11.3 ^b	$\chi^2 = 92.96, p < 0.001$
Not working	20.4 ^a	20.7 ^a	55.6 ^b	$\chi^2 = 259.05, p < 0.001$
Relationship status				
Married/domestic	28.6 ^a	62.8 ^b	65.2 ^b	$\chi^2 = 347.46, p < 0.001$
Single	61.3 ^a	25.9 ^b	16.4 ^c	$\chi^2 = 447.31, p < 0.001$
Other	10.1 ^a	11.3 ^a	18.4 ^b	$\chi^2 = 24.35, p < 0.001$
Frequency of use				
Daily	61.5	64.1	67.5	$\chi^2 = 5.62, p = 0.06$
Weekly	26.1	25.4	25.2	$\chi^2 = 0.22, p = 0.89$
Monthly or less	12.3	10.5	7.3	$\chi^2 = 9.45, p = 0.009$
Quantity of use, %				
≥1 oz	4.1 ^a	7.2 ^b	9.9 ^b	$\chi^2 = 23.05, p < 0.001$
1/4 oz	20.1	22.0	20.9	$\chi^2 = 1.25, p = 0.53$
3–5 g	33.4	27.8	26.3	$\chi^2 = 12.50, p = 0.002$
<2 g	42.5	43.2	42.9	$\chi^2 = 0.89, p = 0.96$
Age of first use				
<14	12.4	17.4	13.2	$\chi^2 = 12.62, p = 0.002$
14–16	40.4	33.4	35.5	$\chi^2 = 12.71, p = 0.002$
17–20	38.0 ^a	28.3 ^b	31.0 ^{a,b}	$\chi^2 = 26.31, p < 0.001$
21+	9.2 ^a	20.9 ^b	20.3 ^b	$\chi^2 = 72.14, p < 0.001$
Method of use				
Inhalation	97.0 ^a	90.6 ^b	81.2 ^c	$\chi^2 = 123.35, p < 0.001$
Oral	2.9 ^a	8.5 ^b	17.8 ^c	$\chi^2 = 117.03, p < 0.001$
Method of selection				
High THC	44.5 ^a	37.1 ^b	41.4 ^{a,b}	$\chi^2 = 13.19, p = 0.001$
High CBD	29.5 ^a	31.7 ^{a,b}	39.4 ^b	$\chi^2 = 16.81, p < 0.001$
Terpenoids	8.6	9.7	10.1	$\chi^2 = 1.30, p = 0.52$
Smell	51.9 ^a	40.8 ^b	30.6 ^c	$\chi^2 = 74.44, p < 0.001$

Percentages represent overall raw percentages. Bolded chi-square results indicate an overall significant difference across the three groups. Different superscripts represent specific group differences with $p \leq 0.001$.

CBD, cannabidiol; THC, delta-9 tetrahydrocannabinol.