

SUPPLEMENTARY TABLE S5. DIFFERENCES IN WITHDRAWAL SYMPTOMS AND BELIEFS ABOUT ADDICTIVENESS OF CANNABIS AMONG RECREATIONAL, MEDICAL, AND MIXED CANNABIS USERS

	<i>Medical</i> (n = 891), %	<i>Recreational</i> (n = 1110), %	<i>Mixed</i> (n = 877), %	
Withdrawal symptoms				
Not applicable	34.8	40.7	29.3	$\chi^2 = 6.25, p = 0.04$
Anxiety	30.5 ^a	12.2 ^b	27.7 ^a	$\chi^2 = 67.88, p < 0.001$
Irritability	35.4	27.6	39.3	$\chi^2 = 9.70, p = 0.008$
Insomnia/interrupted sleep	33.0 ^{a,b}	22.9 ^b	37.3 ^a	$\chi^2 = 18.88, p < 0.001$
Tiredness	8.8	7.3	9.0	$\chi^2 = 6.57, p = 0.04$
Vivid dreams	14.0	14.0	19.6	$\chi^2 = 2.35, p = 0.31$
Loss of productivity	17.3 ^a	6.7 ^b	14.6 ^a	$\chi^2 = 32.77, p < 0.001$
Improved productivity	1.8 ^a	6.9 ^b	5.1 ^{a,b}	$\chi^2 = 15.85, p < 0.001$
Nausea	10.7 ^a	3.2 ^b	7.9 ^{a,b}	$\chi^2 = 20.98, p < 0.001$
Loss of appetite	21.0 ^a	13.4 ^b	23.4 ^a	$\chi^2 = 24.77, p < 0.001$
Weight loss	4.4	3.2	4.3	$\chi^2 = 5.60, p = 0.06$
Tremor	2.7	0.6	1.0	$\chi^2 = 12.40, p = 0.002$
Sweating	4.7	3.4	3.6	$\chi^2 = 5.04, p = 0.08$
Salivation	0.6	0.3	0.9	$\chi^2 = 4.41, p = 0.11$
Addictiveness				
Trouble stopping	13.8	18.8	16.6	$\chi^2 = 4.14, p = 0.13$
Cannabis is addictive	12.7	21.9	15.2	$\chi^2 = 6.71, p = 0.03$
Cannabis not addictive	71.7	64.2	69.8	$\chi^2 = 4.98, p = 0.08$
Don't know if addictive	15.6	13.9	15.0	$\chi^2 = 0.52, p = 0.77$

Percentages represent overall raw percentages (without effects of covariates removed). Bolded chi-square results indicate an overall significant difference ($p \leq 0.001$) across the three groups with the effects of the covariates (bolded in Supplementary Table S1) statistically removed. Different superscripts represent specific group differences with effects of covariates removed and $p \leq 0.001$.