

Impact of cannabis use on brain metabolism using ³¹P and ¹H magnetic resonance spectroscopy

Maximilian Fenzl¹ (ORCID 0000-0003-1011-2754) · Martin Backens¹ (ORCID 0000-0002-3414-696X) · Silviu Bodea² · Miriam Wittemann³ · Florian Werler⁴ · Jule Brielmaier⁵ · Robert Christian Wolf⁴ (ORCID 0000-0002-5358-5212) · Wolfgang Reith¹

1. Institute of Neuroradiology, Saarland University, 66421 Homburg, Germany
2. Helmholtz Zentrum Munich, German Research Center for Environmental Health Institute of Biological and Medical Imaging, 85748 Munich, Germany
3. Department of Psychiatry and Psychotherapy, Saarland University, 66421 Homburg, Germany
4. Department of General Psychiatry at the Center for Psychosocial Medicine, Heidelberg University, 69115 Heidelberg, Germany
5. Department of Obstetrics and Gynecology, RKH Clinic Ludwigsburg, 71640 Ludwigsburg, Germany

Suppl_Table1: Demographics

Demographics and clinical scores	fN percentile			mN percentile			mC percentile			p value ΔCN
	25	median	75	25	median	75	25	median	75	
number of subjects	21			26			40			
Age, years	21.7	22.9	23.8	23.3	24.6	26.5	21.2	23.1	25.7	0.172
Years of education	12.0	13.0	16.0	12.0	12.8	16.5	12.0	13.0	15.8	0.991
Tobacco use, py	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.2	5.0	0.001
BDI	1.0	3.0	5.0	0.0	2.0	4.0	1.5	4.0	8.0	0.143
CUDIT							8.5	12.0	18.5	
Duration of use, years							0.0	2.0	4.5	
Onset use, age							15.8	17.8	20.0	
lifetime joints (x1000)							0.4	1.0	5.5	
Current use, days/week							1.50	4.0	7.00	
Current use, g/week							0.33	2.6	5.25	

Tobacco py: number of packs of cigarettes smoked per day multiplied by the number of years the person has smoked

BDI: Beck depression inventory

CUDIT: Cannabis use identification test

p-values resulting from Mann-Whitney U test between mN and mC groups (ΔCN). p-values < 0.05 are marked in bold.

Demographics and clinical assessment scores of study subjects