

Original Article

The Early Psychosis Working Group

includes:

General Directorate of Health and Social Policies: Sandra Ventura, Maristella Salaris; DMH-PA Piacenza: Silvia Chiesa, Massimo Rossetti, Ornella Bettinardi; DMH-PA Parma: Emanuela Leuci, Lorenzo Pelizza, Antonella Squarcia; DMH-PA Reggio Emilia: Lorenzo Berardi, Francesca Fontana, Sara Paderni; DMH-PA Modena: Francesca Vacca, Nicoletta Nigro, Virginia Guerra; DMH-PA Imola: Vincenza Giannini, Noemi Alagia, Pietro Papili; DMH-PA Bologna: Caterina Bruschi, Lorenzo Gammino, Stefania Artioli; DMH-PA Ferrara: Maria Ferrara, Luigi Grassi, Spyridon Zotos; DMH-PA Ravenna: Sara Sternini, Rosaria Rispoli, Francesca Catena Egitto; DMH-PA Forlì-Cesena: Forlani Claudia, Enrico Meregalli, Elena Soldati; DMH-PA Rimini: Laura Mulazzani, Elena Vagnoni, Michela Pratelli.

Cite this article: Belvederi Murri M *et al.* (2025) The impact of socioeconomic factors on the incidence and characteristics of first-episode psychosis. *Epidemiology and Psychiatric Sciences* **34**, e45, 1–14. <https://doi.org/10.1017/S2045796025100206>

Received: 9 July 2024

Revised: 8 April 2025

Accepted: 23 July 2025

Keywords:



epidemiology; minority issues and cross cultural psychiatry; psychoactive substance use disorder; psychosis; social factors

Corresponding author:

Martino Belvederi Murri;
Email: martino.belvederimurri@unife.it

© The Author(s), 2025. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

The impact of socioeconomic factors on the incidence and characteristics of first-episode psychosis

Martino Belvederi Murri¹ , Alice Onofrio¹, Chiara Punzi¹, Nicola Caranci², Enrico Rubolino³, Francesco Giovinazzi⁴, Danila Azzolina⁵, Federica Folesani¹, Luigi Grassi¹ , Ilaria Tarricone⁶ and Fabrizio Starace⁴

¹Department of Neuroscience and Rehabilitation, Institute of Psychiatry, University of Ferrara, Ferrara, Italy;

²Department of Innovation in Health and Social Care, General Directorate of Health and Welfare,

Emilia-Romagna Region, Bologna, Italy; ³Department of Business and Economics, University of Lausanne,

Lausanne, France; ⁴Department of Mental Health and Pathological Addiction, AUSL Modena, Modena, Italy;

⁵Department of Environmental and Preventive Science, University of Ferrara, Ferrara, Italy and ⁶Department of Medical and Surgical Sciences, Bologna Transcultural Psychosomatic Team (BoTPT), University of Bologna, Bologna, Italy

Abstract

Aim. The environment shapes the risk of psychosis. In particular, urbanicity, deprivation or inequality, migrant density and cannabis availability may not only influence psychosis incidence, but also the characteristics of individuals who arrive at clinical services. This study examined how socioeconomic factors influence the incidence and characteristics of cases of First-Episode Psychosis (FEP).

Methods. We analysed prospective data collected from the FEP early detection programme of Emilia-Romagna, a high-income Italian region. Participants were 1240 individuals aged 18–35 years, who presented at the public healthcare services for a FEP. Exposures were derived from area-level data of 331 municipalities. We used population density, socioeconomic deprivation, educational deprivation, economic inequality, migrant density (proportion of migrants), frequent cannabis use (proportion of people aged 15–19 years old who reported frequent cannabis use). Outcome measures were FEP incidence (cases/100 000 inhabitants at risk per year) and characteristics (age of onset, migrant status, unemployment, substance use, treatment lag [DUP], family and resource problems). We reviewed pertinent literature, and formulated a Directed Acyclic Graph to present causal assumptions and provide adjustment sets for Bayesian spatial and multilevel models of social causation. To compare the effects of different exposures, we computed Average Marginal Effects and report the outcome changes that correspond to one standard deviation change of the exposure, incidence rate ratios (IRR) or odds ratios (OR).

Results. The exposures and incidence of FEP displayed heterogeneous spatial distribution, with no spatially organized pattern. Accordingly, incidence and characteristics were best modelled as non-spatial, three-level hierarchical models. The incidence of FEP was influenced by population density (IRR, 1.14; 95% CrI, 1.03; 1.29), educational deprivation (IRR, 1.15; 95% CrI, 1.02; 1.31) and frequent cannabis use (IRR, 1.31; 95% CrI, 0.98; 1.82), more than socioeconomic deprivation. Higher migrant density in an area shortened the DUP on average by 3.4 months (95% CrI, –1.122; 0.76), while an increase of cannabis use of one standard deviation increased the DUP of 12.9 months (95% CrI, –2.86; 6229). Socioeconomic deprivation increased the likelihood of FEP cases being substance users (OR, 1.12; 95% CrI, 1.01; 1.26), while population density decreased it (OR, 0.91; 95% CrI, 0.83; 1.00).

Conclusions. Area-level socioeconomic features affect both the incidence and the characteristics of FEP, including the probability of individual being migrants, substance users or having a different DUP. Educational deprivation may function as a proxy for culture- or cognitive-related factors. Area-level socioeconomic data may inform public healthcare strategies for early identification and availability of tertiary clinical services.

Introduction

The socioeconomic environment influences the incidence of psychosis, and the clinical profiles of affected individuals. Nonetheless, research on this topic is still warranted, given its clinical and societal importance.

Psychotic disorders have a multifaceted aetiology, at the intersection between genetic predisposition and environmental factors. Particular attention is needed to how social, economic and cultural factors shape their impact across the globe (D'Andrea *et al.*, 2024; Jongasma *et al.*, 2018; Kirkbride *et al.*, 2024). Research began to delineate how specific socioeconomic factors may influence the presentation of psychotic disorders: some trends have emerged in relation to the urban vs. rural settings (Kirkbride *et al.*, 2017), population ethnic composition (Chung *et al.*, 2023), absolute societal deprivation as well as relative economic inequalities (Kirkbride *et al.*, 2014; Logeswaran *et al.*, 2023).

Demonstrating these effects, however, is challenging because of inherent geographic differences, which imply methodological constraints and complex causal interpretation (Grover *et al.*, 2024). For instance, the link between urbanicity and increased incidence of First-Episode Psychosis (FEP) is not consistent worldwide, and exhibits large geographic differences (Jongasma *et al.*, 2018) partly related to the level of analysis (neighbourhood or larger levels), partly because of distinct historical and sociocultural features of each area, that delineate unique 'etiological' landscapes (DeVylder *et al.*, 2018; Kirkbride *et al.*, 2018). In addition, research needs to address the complex causal pathways that underlie social causation, the reciprocal interplay between factors, both at the area-level and individual level, and their distributions and non-linear patterns (González-Valderrama *et al.*, 2022). Notably, recent studies robustly support social causation pathways of psychosis, even accounting for the role of social immobility, social constraints (Logeswaran *et al.*, 2023), and genetic factors (Maxwell *et al.*, 2021). The synthesis of data across distinct domains proves particularly effective to predict the onset of psychosis within communities: a recent study harnessed a range of social, economic, and demographic factors—including age, gender, ethnicity, small-area-level deprivation, social fragmentation, and regional cannabis use—and accurately predicted the incidence of FEP cases across a broad UK region (McDonald *et al.*, 2021). A similar approach could also be applied to explore the impact of socioeconomic factors on the characteristics of FEP users, offering valuable insights for resource allocation and the customization of care.

Aims

The aim of this study was to examine the effects of socioeconomic factors on the treated incidence of first-episode psychosis, and user characteristics. We hypothesized that area-level socioeconomic indices would predict not only the variability of treated incidence, but also key features of presenting users, such as, age of onset, migrant status, unemployment, substance use, treatment lag, family or resource problems.

Methods

Study design and setting

This study uses data of FEP cases collected for the Regional Program for the detection and care of FEP (RER-FEP) between 1 January 2013 and 30 December 2019, first 7 years of activity. The programme was developed within the public health system for the early detection of FEP cases and multi-component care (Regione Emilia Anglin *et al.*, 2023; Belvederi Murri *et al.*, 2021; Romagna, 2016). Data were linked with those from the regional administrative database of the Emilia-Romagna Regional Mental Healthcare

Office, in the same timeframe, using a unique anonymized regional identifier assigned to each user across health records.

Emilia-Romagna is a region in northern Italy with 4.5 million inhabitants and one of the highest GDP per capita in the country, ranking among the most affluent regions in Europe. It includes large urban areas such as Bologna, Modena and Parma, and a mix of smaller towns and rural municipalities. The region has a well-developed public health system and a high standard of living. Around 12% of the population is of foreign origin.

Inclusion criteria

Eligibility was based on the presence of a FEP. Admissible ICD-9 diagnoses were affective psychoses (295.34, 296.24, 296.44, 296.14, 296.54, 296.64) and non-affective psychosis (295.0–295.95, 299.9, 297, 298), excluding substance-induced psychosis. Diagnoses were checked regardless of whether they were recorded as primary or secondary. Other criteria were age between 18 and 35 years; absence of prior treatment with antipsychotics lifetime, absence of intellectual disability. Data include also individuals who were ineligible for the RER-FEP programme and were referred to the CMHC for treatment as usual. Thus, individuals without Italian language proficiency who were excluded from the program, were included in this study.

Ethical approval was obtained by the Area Vasta Emilia Centro Ethical Committee (CE-AVEC); the study conforms to principles expressed in the Declaration of Helsinki. Participants gave their written informed consent prior to their inclusion. This study followed the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) reporting guideline.

Assessments and outcome definitions

Treated incidence was estimated as the number of FEP cases who had access to care per 100 000 persons at risk (aged 18–35 years) per year, per gender, within each municipality. Yearly data of at-risk population were retrieved from the national census (ISTAT). Participants underwent the collection of basic sociodemographic and clinical data, including migrants status, first-generation (born outside Italy) or second-generation (born in Italy from parents born abroad). Presence of substance use (any problematic use of psychoactive substance use) was rated as present by clinicians during clinical contacts, in most cases aided by toxicologic screens. Similarly, clinicians estimated the duration of untreated psychosis (DUP) in months, defined as the time elapsing between the emergence of clear psychotic symptoms and initiation of antipsychotic treatment. The date of onset of psychosis was derived from retrospective clinical interviews. Participants were assessed by psychiatrists using the Italian version of the Health of the Nation Outcome Scales (HoNOS) (Belvederi Murri *et al.*, 2021; Lora *et al.*, 2001). This study analysed in particular items pertaining *Problems with relationships* (item 9), *Activities of daily living and (or) self-care* (10), *Living conditions and (or) family life* (11) and *Occupation and (or) school attendance* (12) to obtain measures of individual social functioning and resource availability (*Supplementary Methods, paragraph S1.3*).

Predictors

We examined the role of population density (log transformed), global deprivation, educational deprivation, economic inequality,

migrant density (indexed by proportion of migrants in each municipality, on the total resident population), cannabis use. Data on density and percentage of people born outside Italy over the total population was obtained from the Italian Institute of statistics and those on deprivation from the ISTAT 2011 census. The set of deprivation indices from the social, economic, housing and education domains, at the level of municipality, was obtained from ISTAT data as defined by a recent study (Caranci *et al.*, 2010; Rosano *et al.*, 2020). We used data of the global deprivation index and an educational deprivation index (percentage of the population aged 15–60 years having elementary education or less). The economic inequality index was computed as the ratio between the subjects in the highest income brackets and subjects in the lowest bracket (Piketty and Saez, 2014). Cannabis use was estimated using 2019 ESPAD survey data on high school students (aged 15–19 years), reporting the provincial-level percentage of individuals using cannabis ≥ 20 times in the past 30 days (Benedetti *et al.*, 2021). Further details on predictors, outcomes and data reduction is reported in the supplement (S1.1–S1.4).

Data analysis

Causal inference

First, we conducted analyses on the effects of socioeconomic variables on incidence, then, on demographic and clinical characteristics of participants. Data analysis was based on a causal inference framework (McElreath, 2018) to support estimation of the effects of interest and identification of variables for model adjustments. We developed a Directed Acyclic Graph (DAG, par. S1.4, Supplementary Fig. 1, SF1) to depict assumptions on causal relationships between socioeconomic factors, psychosis incidence and individual characteristics in the study, supported by a literature review. We then used the DAG to categorize variables as confounders, colliders or mediators for each model. For clarity, we define confounders as variables that causally influence both exposure and outcome, colliders as variables influenced by both, and mediators as variables lying on the causal pathway between exposure and outcome. We adjusted each predictor–outcome analysis for the confounders that were identified by the DAG (Supplementary Table 1, ST1). Unless otherwise specified, we included random slopes for adjustment variables to account for potential variations in their effects across municipalities, in line with recommendations (Barr *et al.*, 2013).

Effect estimation

Data was analysed with Bayesian Generalized Multilevel Models in *brms* (Bürkner, 2017), with outcome-specific distributions: incidence was modelled using a negative binomial distribution to estimate incidence rate ratios (IRRs), since overdispersion was expected (Belvederi Murri *et al.*, 2023). Binary outcomes (e.g., substance use, migrant status) were modelled using a Bernoulli distribution with logit link to estimate odds ratios (ORs); DUP was modelled with a lognormal distribution to account for its positive skew. (Supplement, Par. 1.5). The association between each couple of exposure and outcome was analysed in distinct models of increasing complexity, with municipalities nested in provinces, plus a spatial modified Besag York Mollié (BYM2) model, which accounts for spatial autocorrelation and can identify spatial clusters of outcome distribution (Morris *et al.*, 2019).

The paper reports the exposure–outcome associations that had high credibility, i.e. where the posterior distribution suggested 95% or higher probability of the effect being non-zero (Probability of

Direction, PD). The supplement reports those with lower level of credibility (89–95% PD). We estimated the average marginal effects with their 95% credible intervals (CrI) using *marginal effects* (Arel-Bundock, 2023). In particular, the entity of change of the outcome that would correspond (all else equal) to a change of the exposure spanning: (a) the whole exposure value (minimum to maximum level based on the observed range in the data); (b) one standard deviation around the mean (1SD). Effects are reported both on the response scale and as ratios (IRRs and ORs). Of note, estimates from the observed minimum–maximum range may enhance interpretability, but be more sensitive to outliers.

The *marginal effects* package reports 95% Equal-Tailed Credible Intervals (ETIs), which are symmetric around the median and model-agnostic. However, to better capture where the bulk of the posterior probability lies we additionally computed the probability of direction using the *bayestestR* package, which is based on high-density intervals (HDIs). Credibility assessments are based on HDIs (Makowski *et al.*, 2019).

Results

Causal assumptions and development of the DAG

After the literature review, we built the DAG representing assumptions on the causal relationships between socioeconomic indices and FEP incidence/characteristics (Supplement, Par. S2.1–S2.2). We assumed that population density would drive general deprivation, educational deprivation, inequality and area-level migrant density; such indices would in turn increase the incidence of FEP through increases of unemployment, individual problems, educational attainment, area-level and individual substance use, increased severity of symptoms. In addition, we assumed that socioeconomic disadvantage would also increase the incidence of FEP through a relative lack of resources in the mental health system and increased stigma (Fig. 1). Analysis of the DAG provided the minimal adjustment sets for each model (ST1).

Characteristics of municipalities in the Emilia-Romagna region

The distribution of socioeconomic indices was highly variable across the 331 municipalities and nine provinces (Table 1 and Fig. 2). As expected, socioeconomic indices were inter-correlated: population density correlated with global deprivation ($R = 0.64$), inequality ($R = 0.54$), proportion of migrants ($R = 0.42$) and was weakly negatively correlated with educational deprivation ($R = -0.10$, so that higher density municipalities displayed higher education, consistent with the presence of universities and higher concentrations of younger, more educated residents in larger municipalities in this region). The global deprivation index had a weak positive correlation with inequality ($R = 0.13$), educational deprivation ($R = 0.28$) and migrant density (0.38, all $P < 0.05$). In turn, educational deprivation was negatively correlated with economic inequality, so that areas with lower education had greater inequality ($R = 0.30$). Frequent cannabis use only correlated with economic inequality ($R = 0.26$). The full correlation matrix is reported in SF3, the distribution across provinces in SFs 4–11.

Analysis of clinical data

In the study period 1240 cases of FEP met inclusion criteria (70.5% males, Table 2). Considering an average population at risk of 797

Table 1. Characteristics of municipalities grouped by province

Characteristic	Bologna, N = 55 ¹	Ferrara, N = 23 ¹	Forlì-cesena, N = 30 ¹	Modena, N = 47 ¹	Parma, N = 45 ¹	Piacenza, N = 46 ¹	Ravenna, N = 18 ¹	Reggio emilia, N = 42 ¹	Rimini, N = 25 ¹
Area, mean (SD), km ²	69 (43)	118 (105)	81 (66)	59 (38)	79 (48)	58 (33)	106 (153)	56 (53)	35 (29)
Population at risk, mean (SD), thousands	3.2 (9.9)	2.4 (4.3)	2.3 (4.6)	2.8 (5.3)	1.9 (5.6)	1.1 (2.8)	3.7 (6.2)	2.4 (5.1)	2.5 (5.4)
Population density, Mean (SD), No./km ²	44 (79)	20 (19)	43 (61)	48 (52)	20 (24)	19 (26)	37 (16)	46 (32)	84 (118)
Deprivation indices^a									
Global deprivation index	0.6 (1.8)	0.0 (2.4)	1.2 (2.6)	0.3 (2.8)	-1.6 (3.3)	-2.2 (3.4)	-1.0 (1.6)	0.5 (2.3)	3.7 (2.3)
Educational deprivation index	-0.48 (0.75)	1.00 (1.34)	-0.18 (0.70)	0.30 (0.90)	0.00 (0.94)	-0.09 (1.11)	-0.30 (0.57)	0.47 (0.91)	-0.62 (0.60)
Unemployment index	-0.09 (0.66)	0.40 (0.87)	-0.08 (1.05)	-0.11 (0.76)	-0.45 (0.68)	-0.41 (1.41)	-0.33 (0.80)	0.29 (0.73)	1.38 (0.88)
House occupation index	0.38 (0.67)	-0.54 (0.55)	0.78 (0.72)	0.13 (0.75)	-0.70 (0.95)	-0.74 (1.02)	-0.44 (0.68)	0.13 (0.63)	1.40 (0.72)
Housing index	0.36 (0.76)	-0.34 (0.60)	0.43 (0.85)	-0.04 (1.03)	-0.09 (1.34)	-0.06 (1.30)	0.04 (0.57)	0.00 (0.75)	-0.44 (0.73)
Single-parent index	0.45 (0.56)	-0.56 (0.58)	0.22 (0.72)	0.03 (0.81)	-0.38 (0.77)	-0.86 (0.75)	0.02 (0.40)	-0.37 (0.56)	1.99 (1.08)
Income 0 to 10 000 euros, mean (SD) ^b	20.3 (2.5)	27.6 (8.3)	26.4 (3.3)	24.2 (3.7)	25.2 (6.2)	27.2 (6.1)	24.3 (2.4)	22.8 (2.9)	32.4 (3.5)
Income > 75 000 euros, mean (SD) ^b	2.01 (0.94)	0.92 (0.60)	1.11 (0.56)	1.65 (0.76)	1.69 (0.89)	1.66 (0.94)	1.48 (0.48)	1.87 (0.76)	1.04 (0.62)
Inequality index, mean (SD) ^c	0.10 (0.06)	0.04 (0.03)	0.04 (0.02)	0.07 (0.04)	0.08 (0.05)	0.07 (0.04)	0.06 (0.02)	0.09 (0.04)	0.03 (0.02)
Migrant density, proportion of migrants over total population, mean % (SD)	9.6 (2.2)	7.7 (2.7)	10.3 (3.3)	11.3 (3.2)	10.6 (3.9)	10.1 (4.5)	11.9 (2.4)	10.4 (3.2)	8.2 (2.5)

(Continued)

Table 1. (Continued.)

Characteristic	Bologna, <i>N</i> = 55 ¹	Ferrara, <i>N</i> = 23 ¹	Forlì-cesena, <i>N</i> = 30 ¹	Modena, <i>N</i> = 47 ¹	Parma, <i>N</i> = 45 ¹	Piacenza, <i>N</i> = 46 ¹	Ravenna, <i>N</i> = 18 ¹	Reggio emilia, <i>N</i> = 42 ¹	Rimini, <i>N</i> = 25 ¹
Lifetime cannabis use, % ^d	39.00	33.80	35.10	33.80	34.00	32.10	37.80	34.10	35.30
Frequent cannabis use, males and females, %	6.0 (m), 1.7 (f)	4.8 (m), 1.7 (f)	4.3 (m), 1.4 (f)	4.9 (m), 1.1 (f)	5.5 (m), 2.2 (f)	3.7 (m), 0.9 (f)	6.4 (m), 2.0 (f)	5.0 (m), 1.4 (f)	3.9 (m), 0.6 (f)
Total FEP cases over 7 years, No.	268	101	83	197	220	65	35	197	74
FEP cases/municipality over 7 years, mean (SD)	5 (17)	4 (9)	3 (6)	4 (7)	5 (16)	1 (4)	2 (5)	5 (8)	3 (8)

^az-Scores.^bIncome brackets, percentage of individuals in the bracket, over the total number of taxpayers.^cRatio of the previous income brackets.^dData from the ESPAD survey of cannabis use among 15–19 students in the area.




Figure 1. Directed Acyclic Graph representing the causal assumptions of the study.

thousand individuals per year, the crude average incidence rate of FEP was 21.7 new cases per 100 000 persons/year (95% Confidence Interval [CrI], 19.3–24.1). As expected the incidence rate was higher for males (30.9 cases; 95%CrI: 26.5–35.2) than females (12.5 cases; 95% CrI: 10.4–14.5). The characteristics of participants with FEP displayed a substantial variability across the provinces and municipalities of the region (Table 2).

Effect of socioeconomic factors on FEP incidence

In all instances, multilevel models accounting for municipality and province, including random intercepts and slopes for adjustment variables, performed better than both the BYM2 models and simpler multilevel models with random intercepts only (Table 3). Each analysis were adjusted for confounders identified through the DAG (Supplementary Table 1). Models with cannabis use as the main exposure included only random intercepts, as random slopes did not improve model fit or convergence in this case.

There was a high probability (>95%) that higher population density determined an increase of FEP incidence, so that an increase of population density of the entire range was associated with 10.6 additional cases for females (95% Credible Interval [CrI]: 1.47–18.7) and 22.9 for males (95% CrI: 1.43–40.1). Whereas, an increase of population density of a SD around the mean, there

were 1.77 additional cases per 100 000 subjects per year for females (95% CrI: 0.29–3.07), and 3.81 for males (95% CrI: 0.45–6.63). On average, this corresponds to an IRR of 1.14 (95% CrI: 1.03; 1.29).

A change of the entire range of educational deprivation also corresponded to a significant increase of incidence (Table 3, 98% posterior probability): 18.5 cases per 100 000 subjects per year for females (95% CrI: 1.38; 52.78), and of 37.83 males (95% CrI: 1.00; 104). An increase of one standard deviation of educational deprivation was associated with an increase of 2.00 female cases per 100 000 subjects per year (95% CrI: –0.07; 4.46), and 4.11 for males (95% CrI: 0.58; 9.48). On average this corresponds to an IRR of 1.15 (95% CrI: 1.02; 1.31).

There was a high probability (97%) that higher levels of province-level cannabis use determined an increase of FEP incidence, after adjusting for population density, global and educational deprivation, migrant density, gender, and year, so that an increase of the entire range of cannabis use corresponded to a large increase of incidence (Table 3): 17.2 cases per 100 000 subjects per year for females (95% CrI: –1.07; 61.39), and of 23.67 males (95% CrI: –2.51; 44.14). An increase of one standard deviation of province-level cannabis use was associated with an increase of 5.40 female cases per 100 000 subjects per year (95% CrI: –0.35; 16.04), and 7.16 for males (95% CrI: –0.82; 11.54). On average this corresponds to an IRR of 1.31 (95% CrI: 0.98; 1.82).




Figure 2. Distribution of socioeconomic factors in the Emilia-Romagna region.

Increases of the global deprivation index were associated with an increase of incidence, although with a lower level of credibility (ST 2, posterior probability: 89%, IRR: 1.13; 95% CrI: 0.96; 1.34). Whereas, economic inequality and migrant density were not consistently associated with changes of incidence.

Effect of socioeconomic factors on FEP incidence

Among the most credible effects (Table 3; Fig. 3), economic inequality increased the probability of observing FEP among migrants than among native individuals (OR for a 1SD increase of exposure: 1.28; 95% CrI: 0.95; 1.73). The migrant density in the area was associated with an increase of the probability of observing migrant FEP (OR for a 1sd increase: 1.31; 95% CrI: 1.08; 1.60). Population density was associated with reduced

likelihood of substance use (OR for 1sd increase: 0.91; 95% CrI: 0.83; 1.00). Whereas, global deprivation increased the likelihood of using substances (OR: 1.12; 95%CrI: 1.01; 1.26). Finally, the DUP was affected by migrant density in the area, so that a change of a standard deviation increase of the proportion of migrants corresponded to a DUP 3.3 months shorter and by area-level frequent cannabis use, so that a change of the standard deviation of cannabis use increased the DUP of 13 months (95% probability).

Discussion

This study is one of the few to examine the joint effects of multiple socioeconomic factors on the epidemiology of FEP. In a high-income large European region, with mixed urban and rural

Table 2. Characteristics of participants with FEP







	F, N = 366 ^a	M, N = 874 ^a	P-value ^b
Age			0.049
Mean (SD)	24.4 (5.2)	23.6 (4.6)	
Education			0.005
Elementary/None	8 (2.6%)	20 (2.7%)	
Lower secondary	101 (33%)	293 (40%)	
Upper secondary	155 (50%)	362 (50%)	
Graduate	44 (14%)	55 (7.5%)	
Unknown	58	144	
Civil status			<0.001
Single	265 (85%)	716 (96%)	
Married/Cohabiting	41 (13%)	27 (3.6%)	
Separated/Divorced	4 (1.3%)	2 (0.3%)	
Unknown	56	129	
1st or 2nd generation migrant	90 (25%)	209 (24%)	0.8
Occupation			<0.001
Other	156 (50%)	263 (35%)	
Unemployed	109 (35%)	326 (44%)	
Non-working	2 (0.6%)	3 (0.4%)	
Employed	47 (15%)	156 (21%)	
Unknown	52	126	
Living arrangement			0.3
Other co-living	34 (11%)	83 (11%)	
Other	3 (1.0%)	9 (1.2%)	
Acquired family	70 (23%)	131 (17%)	
Family of origin	177 (57%)	482 (64%)	
Homeless	1 (0.3%)	3 (0.4%)	
Non-psychiatric facilities	4 (1.3%)	16 (2.1%)	
Living alone	19 (6.2%)	33 (4.4%)	
Unknown	58	117	
Substance use			<0.001
Yes	78 (22%)	410 (48%)	
Unknown	11	25	
DUP			0.014
Mean (SD)	8 (12)	9 (12)	
Unknown	19	67	

^an (%).^bWilcoxon rank-sum test; Fisher's exact test.

settings we found that area-level educational deprivation, population density and frequent cannabis use increased the incidence of psychosis. These, and other socioeconomic factors also influenced treatment delay, the probability of observing FEP cases who used illicit substances, and who were migrants. These findings may allow customization of care resources and components to meet the specific needs of the populations with firstFEP.




Our results are largely consistent with those of other recent studies detecting an effect of population density across the rural-urban gradient, and of area-level cannabis use on the incidence of FEP (Brink *et al.*, 2024; Grover *et al.*, 2024; Kirkbride *et al.*, 2017; McDonald *et al.*, 2021). Whereas, unlike other reports, in our study the effect of educational deprivation was stronger than those of economic deprivation. The effect of education is unlikely

Table 3. Effect of socioeconomic factors on FEP incidence and characteristics

Outcome	Exposure	Estimate	Est. Error	l-95% CrI	u-95% CrI	Rhat	Probability coefficient over/below zero ¹	Plot	Average marginal effect ²
Incidence	Population density	0.10	0.04	0.02	0.19	1.00	99% (+)		Min-max: 8.4 cases (0.81; 14.7); IRR 2.54 (1.25; 6.60) 1SD: 1.40 cases (0.20; 2.42); IRR 1.14 (1.03; 1.29)
	Educational deprivation	0.13	0.06	0.00	0.26	1.00	98% (+)		Min-max: 14.2 cases (0.68; 39.41); IRR 3.22 (1.26; 14.3) 1SD: 1.54 cases (0.15; 3.48); IRR 1.15 (1.02; 1.31)
	Area cannabis use	0.15	0.08	-0.01	0.32	1.00	97% (+)		Min-max: 10.39 cases (-0.89; 26.32) IRR 2.34 (0.934; 6.32) 1SD: 3.15 cases (-0.29; 6.86) IRR 1.31 (0.98; 1.82)
Migrant (yes/ no)	Economic inequality	5.03	3.12	-1.23	11.10	1.00	95% (+)		Min-max: 28% (-6%; 59%); OR 3.18 (0.52; 14.5) 1SD: 5% (-1%; 11%); OR 1.28 (0.95; 1.73)
	Migrant density (% migrants)	0.10	0.04	0.03	0.18	1.00	100% (+)		Min-max: 31% (10%; 50%); OR 4.81 (1.00; 23.1) 1SD: 6% (2%; 10%); OR 1.31 (1.08; 1.60)
Substance use (yes/no)	Population density	-0.14	0.07	-0.28	0.01	1.00	97% (-)		Min-max: - 18% (-35%; 0.7%); OR 0.65 (0.44; 1.08) 1SD: -3% (-7%; 0.2%); OR 0.91 (0.83; 1.00)

(Continued)

Table 3. (Continued.)

Outcome	Exposure	Estimate	Est. Error	l-95% CrI	u-95% CrI	Rhat	Probability coefficient over/below zero ¹	Plot	Average marginal effect ²
	Deprivation	0.09	0.04	0.01	0.18	1.00	98% (+)		Min-max: 18% (2%; 50%); OR 3.44 (1.21; 21.8) 1SD: 4% (0.3%; 8%); OR 1.12 (1.01; 1.26)
DUP (months)	Migrant density (% migrants)	-0.04	0.02	-0.08	0.00	1.00	97% (-)		Min-max: - 21.4 months (-7179; 4.16) 1sd: - 3.35 months (-1122; 0.76)
	Area cannabis use	0.18	0.11	-0.03	0.39	1.00	95% (+)		Min-max: 39.6 months (-9.73; 19261) 1SD: 12.9 months (-2.86; 6229)

The table reports the association between each exposure and outcome, only where the exposure had a 95% or higher posterior probability of being non-zero. Each row reports the median parameter value of the exposure (fixed effect), its 95% 95% CrI, Rhat value. 1. The 'Probability coefficient non-zero' column reports the posterior probability of direction, i.e. the probability of the hypothesis that the coefficient of the fixed effect is above (+) or below (-) zero. For instance, population density had a 99% posterior probability of being above zero. 2. The column 'Average Marginal Effect' reports the marginal effect, i.e. the effect on the outcome for modifications of the exposure. Marginal effects are reported in the response scale (i.e. for incidence: number of cases per 100 000 persons year; for DUP: months; for dichotomous outcomes: probability of 'yes'), as OR or IRR. They correspond to a change of the predictor from the minimum to maximum value ('min-max') or to a change of one standard deviation around the mean value (1SD). Predictors (operational definition): Population density (number of residents per km² (log-transformed)); Educational deprivation (% of residents aged 15-60 years with only primary education); Global deprivation (Composite index of social and material disadvantage); Economic inequality (Ratio of population in highest vs lowest income deciles); Migrant density (% of residents born outside Italy); Cannabis use (% of 15- to 19-year-old students reporting ≥ 20 uses in past 30 days in the province); Outcomes: Incidence (new FEP cases per 100 000 at-risk individuals/year); DUP (months elapsing from psychosis emergence to first antipsychotic treatment); Migrant status (FEP case born outside Italy); Substance use (clinically assessed problematic use of any psychoactive substance).

Effects




Figure 3. Average Marginal Effects of socioeconomic factors on FEP incidence and characteristics.

to depend on differences in the DUP or substance use among cases, as we did not detect associations between education deprivation and these outcomes. Emilia-Romagna has a relatively high income and low educational attainment compared with other European regions, and these two indices were weakly correlated in our study, unlike in others (Sirin, 2005). Thus, the effect of education may have emerged because it reflects other social factors than economic ones, which elevate psychosis risk. Educational deprivation may

increase FEP incidence through various mechanisms. First, areas with lower educational levels may be characterized by lower mental health literacy, higher discrimination and higher social pressure towards individuals at risk for psychosis, potentially hastening its onset (Belvederi Murri *et al.*, 2016; Sum *et al.*, 2022; Takizawa *et al.*, 2021). Second, educational deprivation may be linked with worse cognitive performance, as has been recently shown (Lewis *et al.*, 2020). In the study by Lewis and colleagues, this effect was deemed

independent from genetic factors (Lewis *et al.*, 2020). Arguably, societal and individual educational deprivation may lead to worse inferential abilities and more frequent cognitive biases (Corlett and Fletcher, 2021) that might increase psychosis risk. Lastly, individuals with lower education may be subject to overdiagnosis of psychosis because of cultural bias (Morgan *et al.*, 2019).

We identified an association between area-level cannabis use and psychosis incidence, which was not limited to affective psychoses, unlike in recent analyses of the EU-GEI study (Brink *et al.*, 2024). However, direct comparisons are limited by differences in study design, the operationalization of cannabis exposure, and the geographic scale of analysis—including variation in urban/rural composition. These methodological and contextual distinctions likely contribute to the divergent findings and underscore the need for further research on how cannabis use and local sociodemographic environments interact in shaping psychosis risk (Belvederi Murri *et al.*, 2025; Brink *et al.*, 2024).

This study is one of the few, to our knowledge, to link area-level socioeconomic factors with individual characteristics. Few other reports suggest that population density, ethnic composition and income inequality may not only affect incidence rates, but also differences in the clinical presentation (Tibber *et al.*, 2019) and, in some cases, clinical outcomes (Griffiths *et al.*, 2023). In our study, individuals with FEP were more likely to have engaged in substance use if they came from areas with higher deprivation and decreasing population density. The lower prevalence of substance use within high-density areas is different from findings of other studies (Morgan and Mall, 2019; Schifano, 2008), and confirms that the effect of socioeconomic risk factors may present specific patterns and modalities across different settings. In Emilia-Romagna higher-density municipalities are generally wealthy, socially connected, and may offer greater access to health education and preventive services, potentially leading to healthier lifestyles and less harmful use patterns (Belvederi Murri *et al.*, 2023; Brink *et al.*, 2024).

Cannabis use among FEP individuals commonly represent an attempt at self-medication, and requires targeted care strategies, similar to the tailored management required for FEP patients who are either first- or second-generation migrants (Tarricone *et al.*, 2021) or domestic migrants (Tarricone *et al.*, 2016). This study also suggests that mental health services in areas with greater economic disparity should be specifically equipped to care for cases of FEP who are migrants (D'Andrea *et al.*, 2023). Finally, the DUP was increased by area-level cannabis use, and decreased by higher area-level density of migrants. Of note, associations with factors at the area-level should be interpreted differently from those detected at the individual level, as in other studies (Hastrup *et al.*, 2018). Effects of area-level features may reflect differences in the societal appraisal of symptoms. In areas with high cannabis use, behavioural symptoms of psychosis may be more frequently normalized and help-seeking delayed, or may be more frequently linked with paranoid or avoidant psychopathology (Belvederi Murri *et al.*, 2025). In areas with higher migrant density, protective effects of lower stigmatization and better social cohesion may lead to earlier detection of psychosis (Anglin *et al.*, 2023; Morgan *et al.*, 2019; Veling *et al.*, 2008).

Strengths and limitations

The large sample size and long observation period, over a region with mixed urban and rural area suggests that study results are representative of high-income populations of FEP seen in clinical

services. Exposures include a comprehensive array of environmental factors that were measured prospectively, except for deprivation indices. Coupled with a rigorous approach to analyses, our study ought to reduce bias due to confounding and provide reliable estimates of total assumed causal effects (McElreath, 2018).

Limitations include a possible underestimation of incidence (e.g. patients entirely seen within the private sector), despite the consolidated, proactive methodology for early detection (Belvederi Murri *et al.*, 2021). Information on some exposures or confounders were not available, such as detailed ethnic composition, social cohesion, type of cannabis availability, use in other age groups, or other detailed data on cannabis use (e.g. potency), as well as other substance use (Di Forti *et al.*, 2019), diagnosis, individual income or familial characteristics – this may have prevented us from detecting some associations with sufficient level of confidence. While clinicians used a shared, standardized definition of DUP, we did not include a standardized rating instrument and did not collect IRR data. Finally, we did not directly consider hypotheses related to reverse-causation, namely the link between genetic vulnerability and drift towards selected social contexts (Logeswaran *et al.*, 2023; Maxwell *et al.*, 2021). Our DAG reflects a contextual (structural) perspective, where population density shapes the sociodemographic composition of areas. To account for the possibility of compositional confounding, we ran a sensitivity analysis for the density–incidence association, including mean age of FEP and proportion of migrants among FEP cases in each municipality. The results showed sign reversal and poor convergence, potentially indicating a collider structure. Given the rarity of FEP and the high number of municipalities with zero cases, our ability to isolate contextual effects may have been limited. These results do not exclude that compositional mechanisms may also contribute, and further research integrating both perspectives may help to clarify how population structure and area-level factors interact to influence psychosis risk.

Clinical relevance

This study, supporting the ability to predict incidence and characteristics, enables resource allocation, tailoring of clinical services for FEP based on service user needs (McDonald *et al.*, 2021), clinical prediction or support to decisions re. policy. In addition, it may help to target awareness campaigns on psychosis, which might reduce the DUP (Connor *et al.*, 2016) or improve stigma attitudes and help seeking (Belvederi Murri and Amore, 2018; Xu *et al.*, 2018).

In conclusion, we found that the incidence of psychosis was influenced by population density, low population education levels and cannabis use. Area-level deprivation, economic inequality, cannabis use and migrant density influenced the DUP and other individual features of patients with FEP presenting in public mental healthcare services.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S2045796025100206>.

Availability of data and materials. Data are not publicly available, due to inability of the funding agency to share healthcare data with third parties.

Acknowledgements. The supporting organizations had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Author contributions. Prof. B.M., Dr F.G. and Prof D.A. had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis. Concept and design: B.M. and other authors. Acquisition, analysis, or interpretation of data: All authors. Drafting of the manuscript: B.M., A.O., C.P. and I.T. I.T. and F.S. are joint last authors.

Financial support. Early Intervention Service Implementation has been funded by Regione Emilia Romagna since 2012 (Delibera Giunta Regionale Nun. 2155 del 2011). Last funding granted by Delibera Giunta Regionale Num. 1170 del 23/07/2018. Seduta n.32.

Prof. Grassi work has been supported by #NEXTGENERATIONEU (NGEU) and funded by the Italian Ministry of University and Research (MUR), National Recovery and Resilience Plan (NRRP), project MNESYS (PE0000006) – A Multiscale integrated approach to the study of the nervous system in health and disease (DN. 1553 11.10.2022).

Competing interests. None.

Ethical standards. The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Declaration of Helsinki 1975, as revised in 2000.

References

- Anglin DM, Espinosa A, Addington J, Cadenhead KS, Cannon TD, Cornblatt BA, Keshavan M, MATHALON DH, Perkins DO, Stone W, Tsuang M, Woods SW, Walker E, Bearden CE and Ku BS (2023) Association of childhood area-level ethnic density and psychosis risk among ethnoracial minoritized individuals in the US. *Journal of the American Medical Association Psychiatry* **80**(12), 1226–1234. doi:10.1001/jamapsychiatry.2023.2841.
- Arel-Bundock V (2023) *margineffects: Predictions, Comparisons, Slopes, Marginal Means, and Hypothesis Tests*. <https://margineffects.com/> (accessed 1 January 2025).
- Barr DJ, Levy R, Scheepers C and Tily HJ (2013) Random effects structure for confirmatory hypothesis testing: keep it maximal. *Journal of Memory and Language* **68**(3), 255–278. doi:10.1016/j.jml.2012.11.001.
- Belvederi Murri M and Amore M (2018) The Multiple Dimensions of Insight in Schizophrenia-Spectrum Disorders. *Schizophrenia Bulletin* **45**(2), 277–283. doi:10.1093/schbul/sby092.
- Belvederi Murri M, Amore M, Calcagno P, Respino M, Marozzi V, Masotti M, Bugliani M, Innamorati M, Pompili M, Galderisi S and Maj M (2016) The ‘Insight Paradox’ in Schizophrenia: magnitude, Moderators and Mediators of the Association Between Insight and Depression. *Schizophrenia Bulletin* **42**(5), 1225–1233. doi:10.1093/schbul/sbw040.
- Belvederi Murri M, Bertelli R, Carozza P, Berardi L, Cantarelli L, Croce E, Antenora F, Curtarello EMA, Simonelli G, Recla E, Girotto B and Grassi L and , Psychosis, F.W.G. on F.-E. & Team, T.F.W.G. on F.-E.P. in F. includes: I. coordinating (2021) First-episode psychosis in the Ferrara Mental Health Department: incidence and clinical course within the first 2 years. *Early Intervention in Psychiatry* **15**(6), 1738–1748. doi:10.1111/eip.13095.
- Belvederi Murri M, Catania S, Centra S, Folesani F, Muscettola A, Zerbini L, Toffanin T, Ferrara M, Ossola P, Rossi R, Jannini T, Caruso R, Nanni MG and Grassi L (2025) The association between cannabis use and paranoia: meta-analysis of experimental and observational studies. *Neuroscience and Biobehavioral Reviews* **176**, 106269. doi:10.1016/j.neubiorev.2025.106269.
- Belvederi Murri M, Ferrara M, Imbesi M, Leuci E, Marchi M, Musella V, Natali A, Neri A, Ragni S, Saponaro A, Tarricone I, Tullini A and Starace F and Early Psychosis Working Group (for Group Authorship) (2023) A public early intervention approach to first-episode psychosis: treated incidence over 7 years in the Emilia-Romagna region. *Early Intervention in Psychiatry* **17**(7), 724–736. doi:10.1111/eip.13437.
- Benedetti E, Resce G, Brunori P and Molinaro S (2021) Cannabis Policy Changes and Adolescent Cannabis Use: evidence from Europe. *International Journal of Environmental Research & Public Health* **18**(10), 5174. doi:10.3390/ijerph18105174.
- Brink V, Andleeb H, Gayer-Anderson C, Arango C, Arrojo M, Berardi D, Bernardo M, Bobes J, Del-Ben CM, Ferraro L, De Haan L (2024) The Role of Social Deprivation and Cannabis Use in Explaining Variation in the Incidence of Psychotic Disorders: findings From the EU-GEI Study. *Schizophrenia Bulletin* **50**(5), 1039–1049. doi:10.1093/schbul/sbae072.
- Bürkner P-C (2017) {brms}: an {R} Package for {Bayesian} Multilevel Models Using {Stan}. *Journal of Statistical Software* **80**(1), 1–28. doi:10.18637/jss.v080.i01.
- Caranci N, Biggeri A, Grisotto L, Pacelli B, Spadea T and Costa G (2010) The Italian deprivation index at census block level: definition, description and association with general mortality. *Epidemiologia E Prevenzione* **34**(4), 167–176.
- Chung W, Jiang S-F, Milham MP, Merikangas KR and Paksarian D (2023) Inequalities in the Incidence of Psychotic Disorders Among Racial and Ethnic Groups. *American Journal of Psychiatry* **180**(11), 805–814. doi:10.1176/appi.ajp.20220917.
- Connor C, Birchwood M, Freemantle N, Palmer C, Channa S, Barker C, Patterson P and Singh S (2016) Don't turn your back on the symptoms of psychosis: the results of a proof-of-principle, quasi-experimental intervention to reduce duration of untreated psychosis. *BioMed Central Psychiatry* **16**, 127. doi:10.1186/s12888-016-0816-7.
- Corlett PR and Fletcher PC (2021) Modelling delusions as temporally-evolving beliefs (Commentary on Coltheart and Davies). *Cognitive Neuropsychiatry* **26**(4), 231–241. doi:10.1080/13546805.2021.1938984.
- D'Andrea G, Lal J, Tosato S, Gayer-Anderson C, Jongsma HE, Stilo SA, Van Der Ven E, Quattrone D, Velthorst E, Berardi D, Menezes PR (2023) Child maltreatment, migration and risk of first-episode psychosis: results from the multinational EU-GEI study. *Psychological Medicine* **53**(13), 6150–6160. doi:10.1017/S003329172200335X.
- D'Andrea G, Quattrone D, Malone K, Tripoli G, Trotta G, Spinazzola E, Gayer-Anderson C, Jongsma HE, Sideli L, Stilo SA, La Cascia C (2024) Variation of subclinical psychosis across 16 sites in Europe and Brazil: findings from the multi-national EU-GEI study. *Psychological Medicine* 1–14. doi:10.1017/S0033291723003781.
- DeVylder JE, Kelleher I, Lalane M, Oh H, Link BG and Koyanagi A (2018) Association of Urbanicity With Psychosis in Low- and Middle-Income Countries. *Journal of the American Medical Association Psychiatry* **75**(7), 678–686. doi:10.1001/jamapsychiatry.2018.0577.
- Di Forti M, Quattrone D, Freeman TP, Tripoli G, Gayer-Anderson C, Quigley H, Rodriguez V, Jongsma HE, Ferraro L, La Cascia C, La Barbera D (2019) The contribution of cannabis use to variation in the incidence of psychotic disorder across Europe (EU-GEI): a multicentre case-control study. *The Lancet Psychiatry* **6**(5), 427–436. doi:10.1016/S2215-0366(19)30048-3.
- González-Valderrama A, Jongsma HE, Mena C, Castañeda CP, Nachar R, Undurraga J, Crossley N, Aceituno D, Iruretagoyena B, Gallardo C, Mondaca P, Monje M, Irrazaval M, Zavala C, Valmaggia L and Kirkbride JB (2022) The incidence of non-affective psychotic disorders in Chile between 2005 and 2018: results from a national register of over 30 000 cases. *Psychological Medicine* **52**(5), 914–923. doi:10.1017/S0033291720002664.
- Griffiths SL, Bogatsu T, Longhi M, Butler E, Alexander B, Bandawar M, Everard L, Jones PB, Fowler D, Hodgkins J, Amos T, Freemantle N, McCrone P, Singh SP, Birchwood M and Upthegrove R (2023) Five-year illness trajectories across racial groups in the UK following a first episode psychosis. *Social Psychiatry & Psychiatric Epidemiology* doi:10.1007/s00127-023-02428-w.
- Grover S, Varadharajan N and Venu S (2024) Urbanization and psychosis: an update of recent evidence. *Current Opinion in Psychiatry* **37**(3), 191–201. doi:10.1097/YCO.0000000000000931.
- Hastrup LH, Haahr UH, Jansen JE and Simonsen E (2018) Determinants of duration of untreated psychosis among first-episode psychosis patients in Denmark: a nationwide register-based study. *Schizophrenia Research* **192**, 154–158. doi:10.1016/j.schres.2017.05.026.
- Jongsma HE, Gayer-Anderson C, Lasalvia A, Quattrone D, Mulè A, Szöke A, Selten JP, Turner C, Arango C, Tarricone I, Berardi D (2018) Treated incidence of psychotic disorders in the multinational EU-GEI study. *Journal*

- of the American Medical Association Psychiatry. 75(1), 36–46. doi:10.1001/jamapsychiatry.2017.3554.
- Kirkbride JB, Anglin DM, Colman I, Dykxhoorn J, Jones PB, Patalay P, Pitman A, Sonesson E, Steare T, Wright T and Griffiths SL** (2024) The social determinants of mental health and disorder: evidence, prevention and recommendations. *World Psychiatry: Official Journal of the World Psychiatric Association (WPA)* 23(1), 58–90. doi:10.1002/wps.21160.
- Kirkbride JB, Hameed Y, Ankireddypalli G, Ioannidis K, Crane CM, Nasir M, Kabacs N, Metastasio A, Jenkins O, Espandian A, Spyridi S, Ralevic D, Siddabattuni S, Walden B, Adeoye A, Perez J and Jones PB** (2017) The epidemiology of first-episode psychosis in early intervention in psychosis services: findings from the social epidemiology of psychoses in east Anglia [SEPEA] Study. *American Journal of Psychiatry*. doi:10.1176/appi.ajp.2016.16010103.
- Kirkbride JB, Jones PB, Ullrich S and Coid JW** (2014) Social Deprivation, Inequality, and the Neighborhood-Level Incidence of Psychotic Syndromes in East London. *Schizophrenia Bulletin*. 40(1), 169–180. doi:10.1093/schbul/sbs151.
- Kirkbride JB, Keyes KM and Susser E** (2018) City Living and Psychotic Disorders—Implications of Global Heterogeneity for Theory Development. *Journal of the American Medical Association Psychiatry* 75(12), 1211–1212. doi:10.1001/jamapsychiatry.2018.2640.
- Lewis G, Dykxhoorn J, Karlsson H, Khandaker GM, Lewis G, Dalman C and Kirkbride JB** (2020) Assessment of the Role of IQ in Associations Between Population Density and Deprivation and Nonaffective Psychosis. *Journal of the American Medical Association Psychiatry* 77(7), 729–736. doi:10.1001/jamapsychiatry.2020.0103.
- Logeswaran Y, Dykxhoorn J, Dalman C and Kirkbride JB** (2023) Social Deprivation and Population Density Trajectories Before and After Psychotic Disorder Diagnosis. *Journal of the American Medical Association Psychiatry*. 80(12), 1258–1268. doi:10.1001/jamapsychiatry.2023.3220.
- Lora A, Bai G, Bianchi S, Bolongaro G, Civenti G, Erlicher A, Maresca G, Monzani E, Panetta B, Von Morgen D, Rossi F, Torri V and Morosini P** (2001) The Italian version of HoNOS (Health of the Nation Outcome Scales), a scale for evaluating the outcome and the severity in mental health services. *Epidemiologia E Psichiatria Sociale* doi:10.1017/s1121189x00005339.
- Makowski D, Ben-Shachar MS and Lüdtke D** (2019) bayestestR: describing Effects and their Uncertainty, Existence and Significance within the Bayesian Framework. *Journal of Open Source Software* 4(40), 1541. doi:10.21105/joss.01541.
- Maxwell JM, Coleman JRI, Breen G and Vassos E** (2021) Association Between Genetic Risk for Psychiatric Disorders and the Probability of Living in Urban Settings. *Journal of the American Medical Association Psychiatry*. 78(12), 1355–1364. doi:10.1001/jamapsychiatry.2021.2983.
- McDonald K, Ding T, Ker H, Dliwayo TR, Osborn DPJ, Wohland P, Coid JW, French P, Jones PB, Baio G and Kirkbride JB** (2021) Using epidemiological evidence to forecast population need for early treatment programmes in mental health: a generalisable Bayesian prediction methodology applied to and validated for first-episode psychosis in England. *The British Journal of Psychiatry* 219(1), 383–391. doi:10.1192/bjp.2021.18.
- McElreath R** (2018) *Statistical Rethinking: A Bayesian Course with Examples in R and Stan*. New York: Chapman and Hall/CRC.
- Morgan C, Knowles G and Hutchinson G** (2019) Migration, ethnicity and psychoses: evidence, models and future directions. *World Psychiatry: Official Journal of the World Psychiatric Association (WPA)* 18(3), 247–258. doi:10.1002/wps.20655.
- Morgan N and Mall S** (2019) Pathways between urbanization and harmful substance use. *Current Opinion in Psychiatry* 32(3), 218–223. doi:10.1097/YCO.0000000000000488.
- Morris M, Wheeler-Martin K, Simpson D, Mooney SJ, Gelman A and DiMaggio C** (2019) Bayesian hierarchical spatial models: implementing the Besag York Mollie model in stan. *Spatial and Spatio-temporal Epidemiology* 31, 100301. doi:10.1016/j.sste.2019.100301.
- Piketty T and Saez E** (2014) Inequality in the long run. *Science* 344(6186), 838–843. doi:10.1126/science.1251936.
- Regione Emilia Romagna** (2016) *Raccomandazioni regionali per la promozione della salute e del benessere in persone all'esordio psicotico*. 2016. Bologna: Centro stampa della Regione Emilia Romagna. <https://salute.regione.emilia-romagna.it/normativa-e-documentazione/linee-di-indirizzo/archivio-documenti-tecnici/linee-guida/raccomandazioni-regionali-esordio-psicotico-2016> (accessed 1 January 2025).
- Rosano A, Pacelli B, Zengarini N, Costa G, Cislighi C and Caranci N** (2020) Update and review of the 2011 Italian deprivation index calculated at the census section level. *Epidemiologia E Prevenzione* 44(2–3), 162–170. doi:10.19191/EP20.2-3.P162.039.
- Schifano F** (2008) Is urbanization a risk factor for substance misuse? *Current Opinion in Psychiatry* 21(4), 391–397. doi:10.1097/YCO.0b013e328303e2b7.
- Sirin SR** (2005) Socioeconomic status and academic achievement: a meta-analytic review of research. *Review of Educational Research* 75(3), 417–453.
- Sum MY, Chan SKW, Suen YN, Cheung C, Hui CLM, Chang WC, Lee EHM and Chen EYH** (2022) The role of education level on changes in endorsement of medication treatment and perceived public stigma towards psychosis in Hong Kong: comparison of three population-based surveys between 2009 and 2018. *BioMed Central Psychiatry* 22(1), 641. doi:10.1186/s12888-022-04288-1.
- Takizawa N, Melle I, Barrett EA, Nerhus M and Ottesen AA** (2021) The Influence of Mental Health Literacy, Migration, and Education on the Duration of Untreated Psychosis. *Frontiers in Public Health* 9, 705397. doi:10.3389/fpubh.2021.705397.
- Tarricone I, Boydell J, Kokona A, Triolo F, Gamberini L, Sutti E, Marchetta M, Menchetti M, Di Forti M, Murray RM, Morgan C and Berardi D** (2016) Risk of psychosis and internal migration: results from the Bologna First Episode Psychosis study. *Schizophrenia Research* 173(1–2), 90–93. doi:10.1016/j.schres.2016.02.032.
- Tarricone I, D'Andrea G, Storbini V, Braca M, Ferrari S, Reggianini C, Rigatelli M, Gramaglia C, Zeppego P, Gambaro E, Luciano M** (2021) First-episode Psychosis and Migration in Italy: results from a Study in the Italian Mental Health Services (Pep-Ita Study). *Journal of Immigrant & Minority Health* 23(3), 519–527. doi:10.1007/s10903-021-01168-w.
- Tibber MS, Kirkbride JB, Mutsatsa S, Harrison I, Barnes TRE, Joyce EM and Huddy V** (2019) Are socioenvironmental factors associated with psychotic symptoms in people with first-episode psychosis? A cross-sectional study of a West London clinical sample. *BMJ Open* 9(9), e030448. doi:10.1136/bmjopen-2019-030448.
- Velting W, Susser E, Van Os J, Mackenbach JP, Selten J-P and Hoek HW** (2008) Ethnic Density of Neighborhoods and Incidence of Psychotic Disorders Among Immigrants. *American Journal of Psychiatry* 165(1), 66–73. doi:10.1176/appi.ajp.2007.07030423.
- Xu Z, Huang F, Kösters M, Staiger T, Becker T, Thornicroft G and Rüschen N** (2018) Effectiveness of interventions to promote help-seeking for mental health problems: systematic review and meta-analysis. *Psychological Medicine* 48(16), 2658–2667. doi:10.1017/S0033291718001265.