



Published in final edited form as:

*Acta Neuropsychiatr.* 2024 October ; 36(5): 320–324. doi:10.1017/neu.2023.30.

## Cannabidiol negatively modulates adenosine A<sub>2A</sub> receptor functioning in living cells

Nuria Sánchez-Fernández<sup>1,2</sup>, Laura Gómez-Acero<sup>1,2</sup>, Laura I. Sarasola<sup>1,2</sup>, Josep Argerich<sup>1,2</sup>, Andy Chevigné<sup>3</sup>, Kenneth A. Jacobson<sup>4</sup>, Francisco Ciruela<sup>1,2</sup>, Víctor Fernández-Dueñas<sup>1,2</sup>, Ester Aso<sup>1,2</sup>

<sup>1</sup>Pharmacology Unit, Department of Pathology and Experimental Therapeutics, School of Medicine and Health Sciences, Institute of Neurosciences, University of Barcelona, L'Hospitalet de Llobregat, Spain;

<sup>2</sup>Neuropharmacology and Pain Group, Neuroscience Program, Institut d'Investigació Biomèdica de Bellvitge, IDIBELL, L'Hospitalet de Llobregat, Spain;

<sup>3</sup>Immuno-Pharmacology and Interactomics, Department of Infection and Immunity, Luxembourg Institute of Health (LIH), Esch-sur-Alzette, Luxembourg

<sup>4</sup>Molecular Recognition Section, Laboratory of Bioorganic Chemistry, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, MD, USA

### Abstract

**Objectives:** Cannabidiol (CBD) is a phytocannabinoid with great potential in clinical applications. The mechanism(s) of action of CBD require further investigation. Previous studies suggested that adenosine A<sub>2A</sub> receptors (A<sub>2A</sub>Rs) could play a role in CBD-induced effects. Here, we evaluated the ability of CBD to modify the function of A<sub>2A</sub>R.

**Methods:** We used HEK-293T cells transfected with the cDNA encoding the human A<sub>2A</sub>R and G<sub>αs</sub> protein, both modified to perform bioluminescence-based assays. We first assessed the effect of CBD on A<sub>2A</sub>R ligand binding using an A<sub>2A</sub>R NanoLuciferase sensor. Next, we evaluated whether CBD modified A<sub>2A</sub>R coupling to mini-G<sub>αs</sub> proteins using the NanoBiT™ assay. Finally, we further assessed CBD effects on A<sub>2A</sub>R intrinsic activity by recording agonist-induced cAMP accumulation.

**Results:** CBD did not bind orthosterically to A<sub>2A</sub>R but reduced the coupling of A<sub>2A</sub>R to G<sub>αs</sub> protein and the subsequent generation of cAMP.

**Conclusion:** CBD negatively modulates A<sub>2A</sub>R functioning.

---

This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

**Corresponding authors:** Ester Aso; ester.aso@ub.edu; Víctor Fernández-Dueñas; vfernandez@ub.edu.

**Author contribution.** EA, VFD, and FC conceived and designed the study and wrote the manuscript. NSF performed all the experiments and contributed to the manuscript preparation. FC and AC designed the cDNA constructs used in this study. LGA, LIS, and JA cloned and validated cDNA constructs. KAJ provided the fluorescent selective A<sub>2A</sub>R antagonist.

**Competing interests.** None of the authors declare any conflict of interest.

**Ethical standard.** The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on experimentation with cells and DNA constructs.

## Keywords

cannabidiol; adenosine 2A receptor; negative allosteric regulation; competitive binding; cyclic AMP; luminescence-based assays

---

## Introduction

Cannabidiol (CBD) is a phytocannabinoid isolated from *Cannabis sativa* without psychoactive properties, but with potential benefits against multiple pathological conditions (ElSohly *et al.*, 2017). Several preclinical reports demonstrated protective and anti-inflammatory effects of CBD in a wide spectrum of neurodegenerative diseases, neuroinflammatory processes, stroke, colitis, liver, kidney injury, cardiovascular disease, arthritis, sepsis, diabetes, cancer, and epilepsy models (Pacher *et al.*, 2020). Furthermore, CBD exerted positive effects in experimental models of other neuropsychiatric disorders such as epilepsy, anxiety, schizophrenia, dementia, addiction, and neonatal hypoxic-ischemic encephalopathy (Devinsky *et al.*, 2014). Although its translation to clinical trials is somewhat limited to date, the successful case of Epidiolex<sup>®</sup>, an oral solution based on a botanical extract containing purified CBD, is notable. Epidiolex<sup>®</sup> was approved by the US Food and Drug Administration in 2018 for the treatment of Lennox-Gastaut and Dravet syndromes, two rare and debilitating genetic forms of epilepsy in children. Additionally, CBD is currently under clinical evaluation for other conditions, including different forms of pain, obsessive-compulsive disorders, and behavioural problems associated with intellectual disability or autism, among others ([ClinicalTrials.gov](https://clinicaltrials.gov) database).

Despite growing interest in its potential clinical applications, the mechanism(s) of action of CBD require further exploration. CBD has a very low affinity for the orthosteric site of CB<sub>1</sub> and CB<sub>2</sub> receptors, the main G protein-coupled receptors (GPCRs) that belong to the endogenous cannabinoid system (McPartland *et al.*, 2015). Alternatively, CBD can act on multiple targets, including TRPV1 channels and PPAR $\gamma$ , adenosine A<sub>2A</sub>, 5-HT<sub>1A</sub>,  $\alpha_3$ -glycine,  $\alpha_1$ -adrenal, dopamine D<sub>2</sub>, GABA<sub>A</sub>,  $\mu$ - and  $\delta$ -opioid receptors (McPartland *et al.*, 2015). Additionally, CBD can inhibit the activity of GPR55 (Ryberg *et al.*, 2007), an effect that has been associated with its antiepileptic activity (Sylantsev *et al.*, 2013). In the present study, we probed the putative direct effects of CBD on adenosine A<sub>2A</sub> receptors (A<sub>2A</sub>Rs). The relevant role that A<sub>2A</sub>Rs play in several of the neuropsychiatric disorders in which CBD could offer beneficial effects (i.e. dementia, schizophrenia, epilepsy, depression, anxiety) supports this interest (Domenici *et al.*, 2019). Furthermore, previous preclinical evidence supports the participation of A<sub>2A</sub>R in CBD-mediated effects. Thus, A<sub>2A</sub>R antagonists blocked the anti-inflammatory effects of CBD (Liou *et al.*, 2008; Ribeiro *et al.*, 2012; Mecha *et al.*, 2013; Oláh *et al.*, 2014), or the ability of CBD to blunt <sup>9</sup>-THC-induced cognitive impairment (Aso *et al.*, 2019). Similarly, the genetic deletion of A<sub>2A</sub>R reduced the CBD-induced potentiation of the cataleptic and anxiolytic properties of <sup>9</sup>-THC (Stollenwerk *et al.*, 2021). This A<sub>2A</sub>R-dependent activity of CBD was proposed to depend on the ability of CBD to bind to the equilibrative nucleoside transporter (ENT). Thus, inhibition of adenosine uptake would lead to indirect activation of A<sub>2A</sub>R (Pandolfo *et al.*, 2011). However, a direct effect of CBD on A<sub>2A</sub>R has not been further investigated. Here

we aimed to evaluate the capacity of CBD to bind to the orthosteric site of A<sub>2A</sub>R and/or to modify its intrinsic activity by using state-of-the-art luminescence-based assays.

## Materials and methods

### Reagents

The ligands used were CGS21680, ZM241385, and CBD (Tocris Bioscience, Bristol, United Kingdom). MRS7396, a fluorescent selective A<sub>2A</sub>R orthosteric antagonist derived from SCH442416 and containing a BODIPY630/650 fluorophore, was previously described (Duroux *et al.*, 2017). Other reagents used were Dulbecco's modified Eagle's medium (DMEM; Sigma Aldrich, St Louis, MO, USA), geneticin (Santa Cruz Biotechnology, Dallas, TX, USA), adenosine deaminase (ADA; Roche Diagnostics GmbH, Mannheim, Germany), zardaverine (Calbiochem, San Diego, CA, USA), and coelenterazine 400a (NanoLight Technologies, Pinetop, AZ, USA).

### Plasmid constructs

To perform bioluminescence resonance energy transfer (BRET) experiments and cAMP accumulation assays, we used the A<sub>2A</sub>R NanoLuciferase (NanoLuc) sensor (A<sub>2A</sub>R<sup>NL</sup>), previously described (Lanznaster *et al.*, 2019). To perform the NanoBiT™ assay, the cDNA encoding human A<sub>2A</sub>R was cloned at the BamHI/EcoRV restriction enzyme sites of pIRESHyg3-SmBiT (Promega, Madison, WI, USA), as previously described (Sarasola *et al.*, 2022). The construct (A<sub>2A</sub>R<sup>SmBiT</sup>) was verified by DNA sequencing. The plasmid encoding the mini-Gαs (engineered GTPase domain of Gα subunit; LgBiTmini-Gαs) linked to LgBiT was previously described (Wan *et al.*, 2018; Meyrath *et al.*, 2021).

### Cell culture and transfection

Human embryonic kidney (HEK)-293T cells were grown in DMEM supplemented with 1 mM sodium pyruvate (Biowest, Nuaille, France), 2 mM L-glutamine (Biowest), 100 U/mL streptomycin (Biowest), 100 mg/mL penicillin (Biowest), and 5% (v/v) foetal bovine serum (Invitrogen Corporation, Camarillo, CA, USA) at 37°C and in an atmosphere of 5% CO<sub>2</sub>. Cells were transiently transfected with the indicated cDNA construct using polyethylenimine (PEI, 1 mg/mL, Sigma Aldrich), as previously described (Longo *et al.*, 2013). Finally, HEK-293T cells stably expressing A<sub>2A</sub>R<sup>NL</sup> were grown in the presence of geneticin (1 mg/mL).

### NanoBRET experiments

The NanoBRET assay was performed as previously described (Lanznaster *et al.*, 2019). Briefly, HEK-293T cells expressing the A<sub>2A</sub>R<sup>NL</sup> construct were resuspended in Hank's balanced salt solution (HBSS; Thermo Fisher, Waltham, MA, USA) containing ADA (0.5 U/mL) and plated on white 96-well plates coated with poly-ornithine (Corning, Corning, NY, USA) at a density of 20,000 cells/well. After 24 h, cells were challenged with the fluorescent A<sub>2A</sub>R antagonist (MRS7396) in the absence/presence of ZM241385 or CBD and incubated for 1 h at 37°C. Subsequently, coelenterazine 400a was added at a final concentration of 1 μM, and the readings were performed after 15 min using a CLARIOStar microplate reader (BMG Labtech, Durham, NC, USA). Donor and acceptor emission were

measured at  $490 \pm 10$  nm and  $650 \pm 40$  nm, respectively. The raw NanoBRET ratio was calculated by dividing the 650 nm emission by the 490 nm emission and the values fitted by nonlinear regression using GraphPad Prism 9 (GraphPad Software, La Jolla, CA, USA). The results were expressed as a percentage of the maximum signal obtained (mBU; miliBRET units).

### NanoBiT assay

The NanoBiT™ assay (Promega) was performed as previously described (Sarasola *et al.*, 2022). Briefly, transiently transfected HEK-293T cells with A<sub>2A</sub>R<sup>SmBiT</sup> and LgBiT<sup>mini-Gαs</sup> were resuspended in HBSS containing ADA (0.5 U/mL) and transferred (90 μl) into white 96-well plates (Corning). Subsequently, coelenterazine 400a was added (1 μM) to each well. After 15-minute incubation, basal luminescence was determined using a CLARIOstar plate reader (BMG Labtech). Immediately after the initial measurement (basal), the ligands were added, and the luminescent signal was measured every 5 min for 30 min. The luminescence signal (RLU) was normalised as follows:  $(RLU_{\text{sample}} - RLU_{\text{basal}}) / RLU_{\text{basal}}$ .

### cAMP assay

cAMP accumulation was measured using the LANCE® Ultra cAMP Kit (PerkinElmer, Waltham, MA, USA) as previously described (Lanznaster *et al.*, 2019). Briefly, HEK-293T cells stably expressing the A<sub>2A</sub>R<sup>NL</sup> construct were first incubated for 1 h at 37°C with stimulation buffer (BSA 0.1%, ADA 0.5 units/mL, zardaverine 2 μM; in serum-free DMEM) and later with CGS21680 (100 nM) and increasing concentrations of ZM241385 or CBD for 30 min at 37°C. Subsequently, cells were transferred (1000 cells/well) into white 384-well plates (Corning), in which reagents were added following the manufacturer's instructions. After 1 h at room temperature, time-resolved fluorescence resonance energy transfer (TR-FRET) was determined by measuring light emission at 620 nm and 665 nm using a CLARIOstar plate reader (BMG Labtech).

### Statistics

Data are represented as mean ± standard error of mean (SEM) with statistical significance set at  $P < 0.05$ . The number of samples (n) in each experimental condition is indicated in the legend of the corresponding figure. Outliers were assessed using the ROUT method (Motulsky & Brown, 2006); thus, no sample was excluded assuming a  $Q$  value of 1% in GraphPad Prism 9. Comparisons between experimental groups were performed using one-way factor analysis of variance (ANOVA) followed by Dunnett's multiple comparisons *post hoc* test using GraphPad Prism 9 as indicated.

### Results

To assess the impact of CBD on A<sub>2A</sub>R functionality, we initially evaluated whether CBD modified the binding of MRS7396, a fluorescent A<sub>2A</sub>R antagonist. To this end, we took advantage of a previously reported NanoBRET-based A<sub>2A</sub>R binding assay (Fig. 1a) (Lanznaster *et al.*, 2019). HEK-293T cells permanently expressing the A<sub>2A</sub>R<sup>NL</sup> construct were challenged with increasing concentrations of MRS7396, which upon binding to the receptor can act as a compatible acceptor in a BRET process (Fig. 1a). As expected, a



protein at sub-micromolar concentrations (100 nM), thus reducing receptor signalling (i.e. cAMP generation). Therefore, we disclose a new non-competitive interaction of CBD with A<sub>2A</sub>R.

The effect of CBD on A<sub>2A</sub>R could operate through a new allosteric site at the receptor. However, further experiments (i.e. using labelled CBD) would be needed to confirm this hypothesis. On the other hand, we cannot rule out other mechanisms of action for CBD different from classical allosteric drugs. In this sense, previous evidence indicates that other lipids, including the endogenous cannabinoid anandamide at micromolar concentrations, might act as allosteric modulators of other GPCRs through a membrane-perturbing effect that is sensitive to receptor conformation (Lanzafame *et al.*, 2004; Van der Westhuizen *et al.*, 2015). Further studies are needed to assess this putative CBD-mediated membrane effect on A<sub>2A</sub>R-G $\alpha$ s protein coupling. Similarly, CBD could indirectly modify A<sub>2A</sub>R functioning by interacting with equilibrative nucleoside transporter 1 (ENT1), as was previously demonstrated in striatal synaptosomes (Pandolfo *et al.*, 2011). However, this hypothetical CBD effect on ENT seems not to play a relevant role *in vivo*, since a recent study demonstrated that CBD lacks the ability to substantially raise endogenous adenosine levels by using the hypothermia mouse model (Xiao *et al.*, 2023). These discrepancies between *in vitro* and *in vivo* studies could be also explained by the fact that A<sub>2A</sub>R can form heteromers with other GPCRs, including CB<sub>1</sub>R (Carriba *et al.*, 2007; Ferré *et al.*, 2010; Aso *et al.*, 2019), in physiological conditions different from that obtained in heterologous expression systems. The assembly of A<sub>2A</sub>R-containing heteromers leads to changes in the agonist recognition, signalling, and trafficking, which might result in different A<sub>2A</sub>R activity in the presence of CBD.

Although we evaluated the effects of CBD in cultured cells expressing A<sub>2A</sub>R, these results could be relevant for many disorders in which A<sub>2A</sub>R activity increases. For example, in certain inflammatory processes and cardiovascular diseases, but also in pathological conditions that affect the central nervous system, such as Alzheimer's disease, Parkinson's disease, attention deficit hyperactivity disorder, fragile X syndrome, depression, or anxiety (Domenici *et al.*, 2019). A<sub>2A</sub>Rs, which are widely expressed both in neurons and glia, are mainly found in the dorsal and ventral striatum and other nuclei of the basal ganglia, where they play a key role in the control of voluntary movements, as well as in motivational, emotional and cognitive processes (Sebastião and Ribeiro, 2009). In this way, A<sub>2A</sub>Rs are involved in regulating the release of neurotransmitters and contribute to the homeostatic control of synaptic transmission and brain function (Sebastião and Ribeiro, 2009). In general, our results are consistent with the positive effects reported for CBD in various brain disorders that can be associated with an exacerbated A<sub>2A</sub>R function, where CBD would tone down A<sub>2A</sub>R hyperactivity.

Overall, the present study provides evidence on the ability of CBD to negatively modulate A<sub>2A</sub>R signalling. The CBD-mediated negative modulation of A<sub>2A</sub>R function is restricted to the receptor-effector coupling and does not interfere with the binding of the orthosteric ligand. Accordingly, we provide a new and genuine pharmacological way to modulate the adenosinergic system in pathological conditions in which A<sub>2A</sub>R function is increased.

## Acknowledgements.

We thank Centres de Recerca de Catalunya Programme/Generalitat de Catalunya for IDIBELL institutional support and Maria de Maeztu MDM-2017-0729 to Institut de Neurociències, Universitat de Barcelona. We are also grateful to the CanaLatan network members (CYTED-Programa Iberoamericano de Ciencia y Tecnología para el Desarrollo) for the fruitful discussion about the results.

### Financial support.

This study was supported by Ministerio de Ciencia, Innovación y Universidades–Agencia Estatal de Investigación–FEDER funds/European Regional Development Fund – ‘a way to build Europe’ grant RTI2018-097773-A-I00 to EA and PID2020-118511RB-I00 to FC. Founded by MCIN/AEI /10.13039/501100011033 ‘ESF Investing in your future’ grant PRE2018–084480 to JA, grant PRE2019-088153 to NSF and grant FPU19/03142 to LGA. Also supported by ‘Acció instrumental de formació de científics i tecnòlegs’ (SLT017/20/000114) of the Departament de Salut de la Generalitat de Catalunya to LIS. The study was also supported by the Luxembourg Institute of Health (LIH), Luxembourg National Research Fund (INTER/FNRS grants 20/15084569 to AC) and the National Institute of Diabetes and Digestive and Kidney Diseases NIDDK Intramural Research Program (ZIADK031117 to KAJ).

## References

- Aso E, Fernández-Dueñas V, López-Cano M, Taura J, Watanabe M, Ferrer I, Luján R and Ciruela F (2019) Adenosine A<sub>2A</sub>-cannabinoid CB<sub>1</sub> receptor heteromers in the hippocampus: cannabidiol blunts <sup>9</sup>-tetrahydrocannabinol-induced cognitive impairment. *Molecular Neurobiology* 56(8), 5382–5391. [PubMed: 30610611]
- Carriba P, Ortiz O, Patkar K, Justinova Z, Stroik J, Themann A, Müller C, Woods AS, Hope BT, Ciruela F, Casadó V, Canela EI, Lluís C, Goldberg SR, Moratalla R, Franco R and Ferré S (2007) Striatal adenosine A<sub>2A</sub> and cannabinoid CB<sub>1</sub> receptors form functional heteromeric complexes that mediate the motor effects of cannabinoids. *Neuropsychopharmacology* 32(11), 2249–2259. [PubMed: 17356572]
- ClinicalTrials.gov database. The National Library of Medicine. Available at <https://clinicaltrials.gov/> (accessed 16 December 2022).
- Devinsky O, Cilio MR, Cross H, Fernández-Ruiz J, French J, Hill C, Katz R, Di Marzo V, Jutras-Aswad D, Notcutt WG, Martínez-Orgado J, Robson PJ, Rohrback BG, Thiele E, Whalley B and Friedman D (2014) Cannabidiol: pharmacology and potential therapeutic role in epilepsy and other neuropsychiatric disorders. *Epilepsia* 55(6), 791–802. [PubMed: 24854329]
- Domenici MR, Ferrante A, Martire A, Chiodi V, Peponi R, Tebano MT and Popoli P (2019) Adenosine A<sub>2A</sub> receptor as potential therapeutic target in neuropsychiatric disorders. *Pharmacological Research* 147, 104338. [PubMed: 31276772]
- Durox R, Ciancetta A, Mannes P, Yu J, Boyapati S, Gizewski E, Yous S, Ciruela F, Auchampach JA, Gao ZG and Jacobson KA (2017) Bitopic fluorescent antagonists of the A<sub>2A</sub> adenosine receptor based on pyrazolo [4,3-: E] [1,2,4]triazolo[1,5- c] pyrimidin-5-amine functionalized congeners. *Medicinal Chemistry Communications* 8(8), 1659–1667. [PubMed: 29250307]
- ElSohly MA, Radwan MM, Gul W, Chandra S and Galal A (2017) Phytochemistry of Cannabis sativa L. In Kinghorn AD, Falk H, Gibbons S and Kobayashi J (ed), *Phytocannabinoids: unraveling the complex chemistry and pharmacology of Cannabis sativa*. Cham, Switzerland: Springer, pp. 1–36.
- Ferré S, Lluís C, Justinova Z, Quiroz C, Orru M, Navarro G, Canela EI, Franco R and Goldberg SR (2010) Adenosine-cannabinoid receptor interactions. Implications for striatal function. *British Journal Pharmacology* 160(3), 443–453.
- Lanzafame A, Guida E and Christopoulos A (2004) Effects of anandamide on the binding and signaling properties of M1 muscarinic acetylcholine receptors. *Biochemical Pharmacology* 68(11), 2207–2219. [PubMed: 15498511]
- Lanznaster D, Massari CM, Marková V, Šimková T, Durox R, Jacobson KA, Fernández-Dueñas V, Tasca CI and Ciruela F (2019) Adenosine A<sub>1</sub>-A<sub>2A</sub> receptor-receptor interaction: contribution to guanosine-mediated effects. *Cells* 8(12), 1630. [PubMed: 31847113]
- Liou GI, Auchampach JA, Hillard CJ, Zhu G, Yousufzai B, Mian S, Khan S and Khalifa Y (2008) Mediation of cannabidiol anti-inflammation in the retina by equilibrative nucleoside transporter

and A<sub>2A</sub> adenosine receptor. *Investigative Ophthalmology & Visual Science* 49(12), 5526–5531. [PubMed: 18641283]

- Longo PA, Kavran JM, Kim MS and Leahy DJ (2013) Transient mammalian cell transfection with polyethylenimine (PEI). *Methods Enzymology* 529, 227–240.
- McPartland JM, Duncan M, Di Marzo V and Pertwee RG (2015) Are cannabidiol and (9)-tetrahydrocannabinol negative modulators of the endocannabinoid system? A systematic review. *British Journal Pharmacology* 172(3), 737–753.
- Mecha M, Feliú A, Iñigo PM, Mestre L, Carrillo-Salinas FJ and Guaza C (2013) Cannabidiol provides long-lasting protection against the deleterious effects of inflammation in a viral model of multiple sclerosis: a role for A<sub>2A</sub> receptors. *Neurobiology Disease* 59, 141–150.
- Meyrath M, Palmer CB, Reynders N, Vanderplasschen A, Ollert M, Bouvier M, Szpakowska M and Chevigné A (2021) Proadrenomedullin N-terminal 20 peptides (PAMPs) are agonists of the chemokine scavenger receptor ACKR3/CXCR7. *ACS Pharmacology Translational Science* 4(2), 813–823. [PubMed: 33860204]
- Motulsky HJ, Brown RE (2006) Detecting outliers when fitting data with nonlinear regression - a new method based on robust nonlinear regression and the false discovery rate. *BMC Bioinformatics* 7, 123. [PubMed: 16526949]
- Oláh A, Tóth BI, Borbíró I, Sugawara K, Szöllösi AG, Czifra G, Pál B, Ambrus L, Klopper J, Camera E, Ludovici M, Picardo M, Voets T, Zouboulis CC, Paus R and Bíró T (2014) Cannabidiol exerts sebostatic and antiinflammatory effects on human sebocytes. *Journal Clinical Investigation* 124(9), 3713–3724.
- Pacher P, Kogan NM and Mechoulam R (2020) Beyond THC and endocannabinoids. *Annual Review Pharmacology Toxicology* 60, 637–659.
- Pandolfo P, Silveirinha V, dos Santos-Rodrigues A, Venance L, Ledent C, Takahashi RN, Cunha RA and Köfalvi A (2011) Cannabinoids inhibit the synaptic uptake of adenosine and dopamine in the rat and mouse striatum. *European Journal Pharmacology* 655(1–3), 38–45.
- Ribeiro A, Ferraz-de-Paula V, Pinheiro ML, Vitoretti LB, Mariano-Souza DP, Quinteiro-Filho WM, Akamine AT, Almeida VI, Quevedo J, Dal-Pizzol F, Hallak JE, Zuardi AW, Crippa JA and Palermo-Neto J. (2012) Cannabidiol, a non-psychoactive plant-derived cannabinoid, decreases inflammation in a murine model of acute lung injury: role for the adenosine A<sub>2A</sub> receptor. *European Journal Pharmacology* 678(1–3), 78–85.
- Ryberg E, Larsson N, Sjögren S, Hjorth S, Hermansson NO, Leonova J, Elebring T, Nilsson K, Drmota T and Greasley PJ (2007) The orphan receptor GPR55 is a novel cannabinoid receptor. *British Journal Pharmacology* 152(7), 1092–1101.
- Sarasola LI, Del Torrent CL, Pérez-Arévalo A, Argerich J, Casajuana-Martín N, Chevigné A, Fernández-Dueñas V, Ferré S, Pardo L and Ciruela F (2022) The ADORA1 mutation linked to early-onset Parkinson's disease alters adenosine A<sub>1</sub>-A<sub>2A</sub> receptor heteromer formation and function. *Biomedicine and Pharmacotherapy* 156, 113896. [PubMed: 36279718]
- Sebastião AM, Ribeiro JA (2009) Adenosine receptors and the central nervous system. *Handbook Experimental Pharmacology* 193, 471–534.
- Stollenwerk TM, Pollock S and Hillard CJ (2021) Contribution of the adenosine 2A receptor to behavioral effects of tetrahydrocannabinol, cannabidiol and PECS-101. *Molecules* 26(17), 5354. [PubMed: 34500787]
- Sylantsev S, Jensen TP, Ross RA and Rusakov DA (2013) Cannabinoid- and lysophosphatidylinositol-sensitive receptor GPR55 boosts neurotransmitter release at central synapses. *Proceedings of the National Academy of Sciences of the United States of America* 110(13), 5193–5198. [PubMed: 23472002]
- Van der Westhuizen ET, Valant C, Sexton MC and Christopoulos A (2015) Endogenous allosteric modulators of G protein-coupled receptors. *Journal Pharmacology Experimental Therapeutics* 353(2), 246–260.
- Wan Q, Okashah N, Inoue A, Nehmé R, Carpenter B, Tate CG and Lambert NA (2018) Mini G protein probes for active G protein-coupled receptors (GPCRs) in live cells. *Journal Biological Chemistry* 293(19), 7466–7473.

Xiao C, Gavrilova O, Liu N, Lewicki SA, Reitman ML and Jacobson KA (2023) *In vivo* phenotypic validation of adenosine receptor-dependent activity of non-adenosine drugs. *Purinergic Signal* (in press).

Author Manuscript

Author Manuscript

Author Manuscript


Author Manuscript

**Significant outcomes**

- Cannabidiol does not bind orthosterically to A<sub>2A</sub>R.
- Cannabidiol reduces the functionality of A<sub>2A</sub>R


### Limitations

- This is an *in vitro* study, and results cannot be directly extrapolated to *in vivo* conditions.
- Putative allosteric binding of CBD to A<sub>2A</sub>R cannot be confirmed or ruled out with the luminescence-based techniques employed in this study.



**Figure 1.**

Determination of CBD effects on A<sub>2A</sub>R ligand binding affinity. **(a)** Schematic representation of the NanoBRET assay. A nanoluciferase is linked to the N-terminal part of the A<sub>2A</sub>R (A<sub>2A</sub>R<sup>NL</sup>). When the nanoluciferase substrate coelenterazine is added, A<sub>2A</sub>R<sup>NL</sup> (donor) emits light at 490–10 nm. Light excites the fluorescent selective A<sub>2A</sub>R ligand, MRS7396 (acceptor), which subsequently emits fluorescence at 650–80 nm. **(b)** NanoBRET saturation binding curves obtained by challenging A<sub>2A</sub>R<sup>NL</sup> expressing HEK-293T cells with increasing concentrations of MRS7396 in the absence/presence of CBD (1 μM) or ZM241385 (1 μM). **(c)** NanoBRET signals obtained by challenging A<sub>2A</sub>R<sup>NL</sup> expressing HEK-293T cells with a fixed concentration of MRS7396 (30 nM, normalised to 100%) in the presence of increasing concentrations of CBD or ZM241385. The represented data are mean ± SEM of three independent experiments each performed in triplicate.



**Figure 2.**

Assessment of CBD effects on A<sub>2A</sub>R intrinsic activity. **(a)** Schematic representation of the NanoBIT<sup>TM</sup>-based assay. The two fragments of nanoluciferase, small (SmBiT) and large (LgBiT), are fused to A<sub>2A</sub>R and mini-Gαs protein, respectively. Then, upon agonist binding, A<sub>2A</sub>R intrinsic activity is assessed by receptor recruitment of Gαs, which induces an increase on luminescence due to nanoluciferase reconstitution. **(b)** Representative time-course of A<sub>2A</sub>R agonist-mediated Gαs recruitment. The selective A<sub>2A</sub>R agonist CGS21680 was challenged to A<sub>2A</sub>R<sup>SmBiT</sup> and LgBiT mini-Gαs expressing HEK-293T cells in the absence/presence of increasing concentrations of CBD or ZM241385. The luminescent signal obtained after reconstitution of the nanoluciferase was assessed by calculating the area under the curve for each condition. Data are shown as mean ± SEM of three independent experiments with five replicates. \**P* < 0.05, \*\*\**P* < 0.001, one-way ANOVA with Dunnett's *post hoc* test. **(c)** cAMP accumulation was assessed on HEK-293T cells permanently expressing the A<sub>2A</sub>R<sup>NL</sup>. Cells were challenged with the selective A<sub>2A</sub>R agonist CGS21680 (100 nM, normalised to 100% of effect) in the absence/presence of increasing concentrations of CBD. Data are expressed as mean ± SEM of four independent experiments performed in triplicates. \**P* < 0.05, one-way ANOVA with Dunnett's *post hoc* test.