

PLOS ONE

Interventions to address empathy-based stress in mental health workers: A scoping review and research agenda --Manuscript Draft--

Manuscript Number:	PONE-D-23-33814
Article Type:	Research Article
Full Title:	Interventions to address empathy-based stress in mental health workers: A scoping review and research agenda
Short Title:	Interventions for empathy based stress in mental health workers
Corresponding Author:	Hannah May Avon and Wiltshire Mental Health Partnership NHS Trust Bristol, UNITED KINGDOM
Keywords:	empathy based stress; compassion fatigue; Burnout; secondary traumatic stress; mental health workers; Scoping review; research agenda
Abstract:	Consistently engaging with client distress can negatively impact mental health workers (MHWs). Various terms are used to describe this phenomenon, which can all be captured by the concept of empathy-based stress (EBS). Previous reviews have not addressed MHWs as a distinct group, despite evidence suggesting they are particularly vulnerable to EBS. In the context of rising mental health needs triggered by the COVID-19 pandemic, it is especially important to understand how to mitigate the impact of EBS on MHWs. This review therefore aimed to identify and describe available interventions. A systematic scoping review of the literature concerning interventions for EBS in MHWs between 1970 and 2022 was undertaken using five electronic databases. 51 studies were identified, which varied significantly with regards to: interventions used; study methodology and theoretical underpinnings. Studies were grouped according to the level at which they aimed to intervene, namely: individual; team or organisational. Most studies intervened at the level of the individual, despite the proposed causes of EBS being predominantly organisational. Furthermore, theoretical links to the origins of EBS were largely unclear. A dedicated research agenda is set out to address these and other issues identified in the field and to guide future research.
Order of Authors:	Hannah May Josie Millar Emma Griffiths Chris Gillmore Michael West Mhairi Kristoffersen Ross Robinson
Additional Information:	
Question	Response
Financial Disclosure Enter a financial disclosure statement that describes the sources of funding for the work included in this submission. Review the submission guidelines for detailed requirements. View published research articles from PLOS ONE for specific examples.	The author(s) received no specific funding for this work.

This statement is required for submission and **will appear in the published article** if the submission is accepted. Please make sure it is accurate.

Funded studies

Enter a statement with the following details:

- Initials of the authors who received each award
- Grant numbers awarded to each author
- The full name of each funder
- URL of each funder website
- Did the sponsors or funders play any role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript?

Did you receive funding for this work?

Competing Interests

Use the instructions below to enter a competing interest statement for this submission. On behalf of all authors, disclose any [competing interests](#) that could be perceived to bias this work—acknowledging all financial support and any other relevant financial or non-financial competing interests.

This statement is **required** for submission and **will appear in the published article** if the submission is accepted. Please make sure it is accurate and that any funding sources listed in your Funding Information later in the submission form are also declared in your Financial Disclosure statement.

View published research articles from [PLOS ONE](#) for specific examples.

The authors have declared that no competing interests exist.

NO authors have competing interests

Enter: *The authors have declared that no competing interests exist.*

Authors with competing interests

Enter competing interest details beginning with this statement:

I have read the journal's policy and the authors of this manuscript have the following competing interests: [insert competing interests here]

* typeset

Ethics Statement

N/A

Enter an ethics statement for this submission. This statement is required if the study involved:

- Human participants
- Human specimens or tissue
- Vertebrate animals or cephalopods
- Vertebrate embryos or tissues
- Field research

Write "N/A" if the submission does not require an ethics statement.

General guidance is provided below. Consult the [submission guidelines](#) for detailed instructions. **Make sure that all information entered here is included in the Methods section of the manuscript.**

Format for specific study types

Human Subject Research (involving human participants and/or tissue)

- Give the name of the institutional review board or ethics committee that approved the study
- Include the approval number and/or a statement indicating approval of this research
- Indicate the form of consent obtained (written/oral) or the reason that consent was not obtained (e.g. the data were analyzed anonymously)

Animal Research (involving vertebrate animals, embryos or tissues)

- Provide the name of the Institutional Animal Care and Use Committee (IACUC) or other relevant ethics board that reviewed the study protocol, and indicate whether they approved this research or granted a formal waiver of ethical approval
- Include an approval number if one was obtained
- If the study involved *non-human primates*, add *additional details* about animal welfare and steps taken to ameliorate suffering
- If anesthesia, euthanasia, or any kind of animal sacrifice is part of the study, include briefly which substances and/or methods were applied

Field Research

Include the following details if this study involves the collection of plant, animal, or other materials from a natural setting:

- Field permit number
- Name of the institution or relevant body that granted permission

Data Availability

Authors are required to make all data underlying the findings described fully available, without restriction, and from the time of publication. PLOS allows rare exceptions to address legal and ethical concerns. See the [PLOS Data Policy](#) and [FAQ](#) for detailed information.

Yes - all data are fully available without restriction

A Data Availability Statement describing where the data can be found is required at submission. Your answers to this question constitute the Data Availability Statement and **will be published in the article**, if accepted.

Important: Stating 'data available on request from the author' is not sufficient. If your data are only available upon request, select 'No' for the first question and explain your exceptional situation in the text box.

Do the authors confirm that all data underlying the findings described in their manuscript are fully available without restriction?

Describe where the data may be found in full sentences. If you are copying our sample text, replace any instances of XXX with the appropriate details.

- If the data are **held or will be held in a public repository**, include URLs, accession numbers or DOIs. If this information will only be available after acceptance, indicate this by ticking the box below. For example: *All XXX files are available from the XXX database (accession number(s) XXX, XXX).*
- If the data are all contained **within the manuscript and/or Supporting Information files**, enter the following: *All relevant data are within the manuscript and its Supporting Information files.*
- If neither of these applies but you are able to provide **details of access elsewhere**, with or without limitations, please do so. For example:

Data cannot be shared publicly because of [XXX]. Data are available from the XXX Institutional Data Access / Ethics Committee (contact via XXX) for researchers who meet the criteria for access to confidential data.

The data underlying the results presented in the study are available from (include the name of the third party

All relevant data are within the manuscript and its Supporting Information files.

and contact information or URL).

- This text is appropriate if the data are owned by a third party and authors do not have permission to share the data.

* typeset

Additional data availability information:

Interventions to address empathy-based stress in mental health workers: A scoping review and research agenda

1 **Hannah May^{1,2*}, Emma Griffith^{1,2}, Josie Millar¹, Chris Gillmore², Mhairi Kristoffersen¹, Ross**
2 **Robinson², Michael West³**

3 ¹ University of Bath, Department of Psychology, Bath, UK

4 ²Avon and Wiltshire Mental Health Partnership NHS Trust, Bath, UK

5 ³ University of Lancaster, Management School, Lancaster, UK

6 *** Correspondence:**

7 Hannah May

8 hannahrebeccamay@hotmail.co.uk

9 **Word count** (excluding abstract, tables, references and appendices): 9698

10 **Tables:** 7 in body of text; 3 additional in appendices (all tables are found at the end of this draft)

11 **Figures:** 3

12 **Keywords:** empathy based stress¹, compassion fatigue², burnout³, secondary traumatic stress⁴,
13 mental health workers⁵, scoping review⁶, research agenda⁷

14

15

16

17

18

19

20

21

22

23

Abstract

24
25 Consistently engaging with client distress can negatively impact mental health workers (MHWs).
26 Various terms are used to describe this phenomenon, which can all be captured by the concept of
27 empathy-based stress (EBS). Previous reviews have not addressed MHWs as a distinct group, despite
28 evidence suggesting they are particularly vulnerable to EBS. In the context of rising mental health
29 needs triggered by the COVID-19 pandemic, it is especially important to understand how to mitigate
30 the impact of EBS on MHWs. This review therefore aimed to identify and describe available
31 interventions. A systematic scoping review of the literature concerning interventions for EBS in
32 MHWs between 1970 and 2022 was undertaken using five electronic databases. 51 studies were
33 identified, which varied significantly with regards to: interventions used; study methodology and
34 theoretical underpinnings. Studies were grouped according to the level at which they aimed to
35 intervene, namely: individual; team or organisational. Most studies intervened at the level of the
36 individual, despite the proposed causes of EBS being predominantly organisational. Furthermore,
37 theoretical links to the origins of EBS were largely unclear. A dedicated research agenda is set out to
38 address these and other issues identified in the field and to guide future research

39

Introduction



41 **What is Empathy-based Stress?**

42 Empathy is the interpersonal capacity to be sensitive to others and understand their feelings by
43 actively entering into their experiences (Elliot et al., 2018). It is thought to be a key component of
44 compassion (Atkins & Parker, 2012; Strauss et al., 2016), wherein engagement with the suffering of
45 another facilitates actions to prevent and alleviate that suffering (Gilbert, 2014). People accessing
46 mental health services are found to have better outcomes in terms of mental health recovery if they

47 experience compassionate (Spandler & Stickley, 2011) and empathic (Elliot et al., 2018)
 48 relationships with staff caring for them. This may be especially important in mental health services
 49 given the stigma that persists in both society and health services regarding mental distress
 50 (Ahmedani, 2011; Henderson et al., 2014). Clinicians also stand to benefit from such practice, with
 51 evidence highlighting that compassionate practice can protect against the harmful effects of empathic
 52 distress (Klimecki et al., 2014); lead to positive emotions (Klimecki et al., 2013); and reduce
 53 psychological distress (Kirby et al., 2017).

54 Despite the benefits associated with compassionate and empathic care, research has also identified
 55 potential costs to clinicians working with distressed clients. These include: burnout (Maslach, 1993);
 56 secondary traumatic stress (STS; Figley, 1995; Newell & MacNeil, 2010); compassion fatigue (CF;
 57 Figley, 1995); and vicarious trauma (VT; Pearlman & Saakvitne, 1995). Whilst distinct definitions
 58 exist for these terms (see Table 1), the extent to which they differ from one another is unclear and
 59 they are often used interchangeably in the literature (Turgoose & Maddox, 2017; Rauvola et al.,
 60 2019).

61 **Table 1**

62 *Definitions of terms included in the Empathy-based stress construct*

Term	Definition
Burnout (Maslach, 1993).	A psychological response to chronic interpersonal stressors at work, resulting in exhaustion, detachment and feelings of incompetence. First identified in caregiving professions where services are provided via a relationship with people in need (Maslach 1982; 2006). These relationships (whilst rewarding) necessitate prolonged and intense emotional contact with clients which over time can produce burnout (Maslach, 2006).
Secondary Traumatic Stress (STS)	A form of second-hand post-traumatic stress disorder (PTSD). In STS, symptoms (e.g. hypervigilance, avoidance, intrusive

Term	Definition
(Figley, 1995; Newell & MacNeil, 2010).	thoughts/memories) arise out of exposure to the trauma of others, rather than through direct personal experience.
Compassion Fatigue (CF) Figley (1995).	Figley (1995) substituted STS with the term CF as a less stigmatising way of referring to the impact of these experiences on healthcare professionals. CF is “a state of exhaustion and dysfunction biologically, psychologically, and socially as a result of prolonged exposure to compassion stress” (p. 253). CF has an inverse relationship with CS; leading to the suggestion that overwhelming levels of CF may interfere with the capacity to experience CS (Rossi et al., 2013; Stamm, 2002).
Vicarious Trauma (VT) (Pearlman & Saakvitne, 1995).	VT is defined as “the transformation that occurs in the inner experience of the therapist [or worker] [...] as a result of empathic engagement with clients’ trauma material” (Pearlman & Saakvitne, 1995, p. 31). Whilst VT may result in PTSD-type responses (flashbacks, intrusions, dissociation) (Beaton & Murphy, 1995), definitions emphasise the negative impact on the clinician’s cognitive schemas (Pearlman & Saakvitne, 1995).

63 *Note.* STS = secondary traumatic stress; CF= compassion fatigue; VT= vicarious trauma; PTSD=
64 post-traumatic stress disorder
65

66 These inconsistencies led Rauvola et al. (2019) to propose an alternative conceptualisation in which
67 VT, burnout, CF and STS are all highly related aspects of ‘empathy-based stress’ (EBS; Figure 1).
68 EBS is described as applicable to anyone whose occupation exposes them to the distress of others
69 and requires that they respond with engagement and empathy. It is defined as “a stressor–strain-based
70 process of trauma at work, wherein exposure to secondary or indirect trauma, combined with
71 empathic experience, results in empathy-based strain and additional outcomes (i.e., other
72 occupational health/strain outcomes; work affect, behaviours, and cognitions)” (Rauvola et al., 2019,
73 p299). In the EBS model; VT, STS and CF are the three varying forms of empathy-based strain that
74 result from empathic engagement with the traumatic experiences of others. The state of burnout is

75 conceptualised as a reaction to this strain, though the authors acknowledge there are common features
76 between all four constructs.

77 EBS is a recently conceptualised model which has not yet been applied within the wider literature;
78 therefore, its feasibility and usefulness as a framework remain untested. However, EBS brings
79 together established pre-existing concepts in a way that allows for distinct but inter-related terms to
80 co-exist, thereby offering a potential solution to the conceptual ambiguities outlined above.
81 Furthermore, it allows that anyone who empathically engages with second-hand distress in a work
82 setting can consequently experience undesirable outcomes, which can interact with pre-existing
83 individual factors to make EBS more or less likely/ severe. It is therefore dynamic, process-driven
84 and applicable across professional fields. In this way it is akin to established models in the
85 occupational health literature, including the Job Demands- Resources model which posits that all
86 jobs comprise interacting sets of psychologically taxing demands and rewarding job resources
87 (Bakker & Demerouti, 2007) . This review will henceforth use the term EBS, in reference to the
88 dynamic model of “traumatic stressor exposure, empathic experience, and adverse reactions” (p. 297)
89 described by Rauvola et al.

90 **Why Prevent/Treat Empathy Based Stress in Mental Health Workers?**

91 There is evidence to suggest that EBS impacts care providers from diverse professional backgrounds
92 including nurses, physicians, social workers and mental health workers (Cavanagh et al., 2020; Rossi
93 et al., 2013). Mental health workers (MHWs) however, are often required to engage explicitly with
94 their clients’ distressing experiences during care provision (Canfield, 2005). For example, successful
95 psychotherapy is thought to depend in part on the clinician’s ability to empathise with and to some
96 degree absorb the pain that clients are experiencing (Rogers, 1967; Rothschild & Rand, 2006).
97 Compared to GPs and non-medical personnel, MHWs have been found to show greater empathic

98 concern for others and higher discomfort at witnessing suffering (Santamaría-García et al., 2017).
99 This empathic engagement predicts better client outcomes (Elliot et al., 2018); however, leaves
100 MHWs at increased risk of EBS (Canfield, 2005; Jachens et al., 2018).

101 The rates of trauma experiences are found to be higher amongst people accessing mental health
102 services, with estimates suggesting that roughly half have experienced physical abuse, whilst over a
103 third have experienced sexual abuse (Mauritz et al., 2013). As services become increasingly aware of
104 the importance of trauma-informed approaches to health and social care (Sweeney et al., 2016)
105 MHWs are being encouraged to explicitly ask clients about potential traumatic experiences (Read et
106 al., 2007). While undoubtedly a positive step, this seems likely to increase the quantity of traumatic
107 material MHWs are exposed to. In their review, Turgoose and Maddox (2017) found that greater
108 exposure to client trauma/distress was associated with higher CF in MHWs. EBS is associated with a
109 range of adverse outcomes including poor job satisfaction, absenteeism and poor patient care (Figley
110 2002; Mathieu, 2007). It can lead to what Baranowsky (2002) termed ‘the silencing response’,
111 whereby health workers begin to minimise, shutdown or divert away from traumatic material brought
112 to them by the client.

113 **Current Review and Aims**

114 Previous reviews targeting healthcare workers suggest that intervention/training may be effective in
115 reducing EBS (e.g. Cocker & Joss, 2016), however, existing reviews have either amalgamated
116 MHWs with other healthcare workers as one group (e.g. Conversano et al., 2020) or excluded them
117 (e.g. Cocker & Joss, 2016). Bercier and Maynard (2014) attempted to review EBS interventions for
118 MHWs, however were unable to identify relevant studies and called for more research. Sutton et al
119 (2022) examined the impact of organisational factors on some specific forms of EBS, however did

120 not consider other forms of intervention and looked only at particular professional groups working in
121 mental health.

122 Prompted by these challenges, this paper aimed to review available interventions for preventing or
123 treating EBS in MHWs specifically. Given the urgency indicated by staff stress levels (NHS, 2023)
124 and workforce shortages it seems timely to ask; “What interventions exist to prevent or treat EBS in
125 MHWs?”. Using systematic scoping methodology, we aimed to answer this question via the
126 following objectives:

- 127 1) To assess and summarise the available interventions for preventing or treating EBS in MHWs
- 128 2) To ascertain the theoretical underpinnings and assumptions of these interventions
- 129 3) To assess how EBS has been measured in these studies
- 130 4) To make recommendations for future research into the treatment and prevention of EBS in
131 MHWs

132 Once the authors had conducted the initial review and assessed the nature of the available literature, a
133 second aim was set: to provide a dedicated research agenda that summarises the existing state of the
134 literature on interventions for EBS in MHWs and sets out key issues and directions for future
135 research.

136 NHS England (2018) warned that the ever-growing need for healthcare services leaves NHS staff
137 vulnerable to CF, which has been linked to depression, stress and chronic illness (Barr, 2017;
138 Canfield, 2005; Demerouti, 2000; Mathieu, 2007). These are serious concerns for the workforce, as
139 demonstrated in a recent NHS staff survey which found that 44.8% of workers felt unwell due to
140 work-related stress; a percentage that has been climbing steadily since 2016 (NHS, 2023). This has
141 led West and Chowla (2017) to call for a culture change in the NHS to enable senior staff to lead
142 compassionately and sensitively rather than being solely driven by achieving targets and reducing



143 costs. However, Dean et al. (2019) highlight that such shifts require changes in the framework of our
144 healthcare systems that will not be simple or quick to achieve.

145 **Method**

146 **Scoping Review**

147 This review employed a systematic scoping review method. Scoping reviews offer a means to map
148 out the scope of the literature in a given area and to provide an overview of the type of evidence
149 available (Mays et al., 2001). Whilst they may be a precursor to systematic reviews, they can also be
150 conducted as self-contained reviews when there is insufficient knowledge about a subject to generate
151 sufficiently specific questions for a systematic review (Armstrong et al., 2011).

152 To this end, scoping reviews: have broader inclusion criteria compared to systematic reviews; **do not**
153 **assess bias in included studies;** do not attempt to synthesise evidence or arrive at precise
154 recommendations. Instead, they aim to outline the research that is available, how it has been
155 conducted and what theory/evidence has informed it. Despite these differences, scoping reviews are
156 still systematic in nature and involve an a priori protocol and a methodical, exhaustive, and replicable
157 search strategy (Arksey & O'Malley, 2005; Munn et al., 2018).

158 Munn et al (2018) provide guidance for authors choosing whether to conduct a systematic or scoping
159 review. They give six indications for conducting scoping reviews, four of which are relevant to the
160 current review: “To identify the types of available evidence in a given field; To clarify key concepts/
161 definitions in the literature; To examine how research is conducted on a certain topic or field... To
162 identify and analyse knowledge gaps” (pp.2). The review was conducted in accordance with the
163 Joanna Briggs Scoping Review Framework (Peters et al., 2017) and written up in line with the
164 Prisma Extension for Scoping Reviews (Tricco et al., 2018). See Appendix A for a completed

165 Preferred Reporting Items for Systematic reviews and Meta-Analyses extension for Scoping Reviews
166 (PRISMA-ScR) Checklist.

167 **An a priori protocol** was developed and registered on Open Science Framework: <https://osf.io/b7kcr/>

168 **Electronic searches and Search Terms**

169 Searches were conducted in the following databases: PubMed, PsychINFO (including APA PsycInfo,
170 APA PsycArticles, APA PsycExtra and APA PsycTests), Embase, PTSDPubs. Grey literature was
171 accessed via PsycExtra (theses) and the Cochrane Central Register of Controlled Trials (CENTRAL;
172 unpublished trials). Studies available in English and conducted between 1970 (when the term burnout
173 was introduced; Schaufeli et al., 2009) and 2021 were included in the initial search; in the updated
174 search the included date range was 2021 to 2022.

175 The search terms were determined by considering the core elements of the research question, in
176 consultation with a research librarian (see Table 2). Given the inconsistent/overlapping definitions of
177 terminology noted in the literature (Turgoose & Maddox, 2017; Rauvola et al., 2019), EBS search
178 terms incorporated the empathy-based strain constructs identified by Rauvola et al. (VT, STS, CF) as
179 well as burnout (proposed to be an adverse reaction to empathy-based strain). Precise terminology,
180 boolean operators and truncation were adapted/applied in accordance with each database (see
181 Appendix B for full search strategy). When full texts could not be retrieved via inter-library loan,
182 effort was made to locate/contact authors. Initial searches began in December 2020 and were
183 completed in January 2021. An updated search was subsequently carried out in September 2022. All
184 identified papers were uploaded to Covidence: <https://app.covidence.org/reviews/130454>

185

186

187 **Table 2**

188 *Example of Search Terms*

189

Search item of interest	Provisional search terms
Empathy based stress	Burnout OR Compassion fatigue OR secondary traumatic stress OR STS OR vicarious trauma OR VT OR compassion stress
Mental health workers	mental health nurs* OR forensic nurs* OR counsellor* OR psychologist* OR psychiatrist* OR psychotherapist* OR therapist* OR mental health support worker* OR mental health social worker* OR support time and recovery worker* OR family therapist* OR therapist* OR CBT therapist* OR mental health occupational therapist*
Intervention	Treat* OR prevent* OR train* OR program* OR interven*

190

191 **Study selection**

192 Titles and abstract screening as well as full-text review was conducted by two independent reviewers
193 (HM and MK/RR) according to the inclusion/exclusion criteria shown in Table 3. Whilst the mental
194 health worker search terms in Table 2 were used to guide the searches, some identified studies used
195 different terms to describe participant job roles, or else more general terms e.g. doctor, nurse. In these
196 cases the studies were carefully read in order to identify the nature of the client group participants
197 were working with, and/or the study setting. Such studies were included only if they clearly
198 referenced client groups with mental health needs e.g. clients with psychosis, or healthcare settings
199 that clearly had a mental health focus e.g. psychiatric wards.

200

201 **Table 3**

202 *Criteria for study inclusion/exclusion*

	Inclusion Criteria	Exclusion Criteria
Population	At least 50% of participants are mental health workers (see Table 2 for included professions/synonyms).	Studies including other healthcare staff whose role does not have a mental health focus/the general public will be excluded.
		Children, adolescents
Concept	Studies involving any intervention (including organisational strategies e.g. increased supervision) or training program that aims to prevent or reduce EBS (including STS, CF, VT & burnout).	Studies that only assess MHW's individual differences and/or self-care strategies in relation to EBS.
	Studies that in some way evaluate the impact of the intervention/training on EBS.	
Context	Study settings may include any setting where mental health workers are employed.	
Source	Primary research studies, systematic reviews and meta-analyses of any publication status.	Theoretical papers, opinion pieces or letters to the editor
	Qualitative or quantitative studies.	Books, book chapters

203 *Note.* EBS = empathy-based stress; MHWs = mental health workers; STS = secondary traumatic stress;
 204 CF = compassion fatigue; VT = vicarious trauma.

205

206 **Charting the Data**

207 Information from included studies was charted by reviewers HM and MK, using the following
208 headings:

209 *Characteristics of Included Studies*

210 **Study characteristics.** Authors, year of publication, study location, methodology, study
211 design, study aims, EBS construct targeted, type of **intervention/training**, method of evaluating EBS.

212 **Sample characteristics.** Sample size, age, gender, setting, % of MHWs in sample, specific
213 profession/role of MHWs.

214 *Characteristics of Intervention/Training*

215 This included: intervention/training type; the aims and rationales for the interventions/training
216 programs administered; duration of intervention; and method of delivery. Information about the
217 theoretical underpinnings of the intervention/training provided were recorded.

218 *Outcomes*

219 Measures used to assess EBS and, where possible, outcomes on these measures were recorded. Any
220 recommendations for practice or further research on training/interventions of this nature were
221 documented.



222 **Results**


223 Searches of all sources retrieved 20,552 records. After duplicates ($n = 10964$) and other ineligible
224 sources (see Figure 2) were removed, 9,590 titles and abstracts were screened of which 9,436 were
225 excluded. Of the 228 full text records screened, 177 were excluded. Therefore, 51 studies that met
226 eligibility criteria (Figure 2) were included in the review. In this section, the characteristics of the
227 included studies and their participants are first described. Thereafter, the EBS interventions used in the

228 studies are summarised with reference to their: nature; length; theoretical underpinnings and
 229 assumptions and; means of measurement. Though limited by considerable heterogeneity in the
 230 identified studies, the authors also attempt to place the findings in the context of relevant occupational
 231 health research by considering how they map on to the three types of organisational intervention for
 232 improving staff wellbeing and mental health described by Tetrick and Quick (2011).

233

234 **Figure 2**

235 *PRISMA Flow Chart of the Screening Process*



258

259 **Characteristics of Included Studies**

260 Table 4 provides an overview of the key features of the 51 included studies. The year of publication
261 ranged from 1981 to 2021. The majority of studies ($n = 28$) were conducted in the United States of
262 America (USA). Most studies ($n = 43$) used a quantitative methodology, but qualitative ($n = 2$) and
263 mixed methods ($n = 6$) were also used. Study designs varied, ranging from case studies ($n = 1$) to
264 randomised controlled trials (RCTs) ($n = 5$) with repeated measures designs ($n = 25$) being the most
265 common. Samples varied in size from $N = 2$ to $N = 296$ and consisted of a diverse range of MHWs,
266 most commonly nurses who comprised 100% of the sample in nine studies, and part of the sample in
267 eight. Fourteen studies had sample populations consisting of multiple types of MHWs. Of the 40
268 studies that provided gender breakdowns: 27 comprised over 50% women; and seven were 100%
269 women. Only 19 studies provided information on participant ethnicity (see Table 5), most of which
270 had majority ($n = 14$) or entirely ($n = 1$) White samples. Other ethnicities represented included Black,
271 Hispanic, Latino/Latina, Asian, Indian, Chinese, Malay, Mauritian, multi-racial, American Indian and
272 Pacific Islander. Eight of the 20 studies reporting ethnicity described a proportion of participants as
273 having ‘other’ or ‘non-White’ ethnicity, with no further information provided. A further four studies
274 reported a percentage of White participants but provided no indication of the ethnic backgrounds of
275 the rest of the sample. Settings included: community mental health ($n = 14$); inpatient ($n = 7$);
276 outpatient/clinic ($n = 4$); multiple settings ($n = 3$); forensic ($n = 3$); mental health centres ($n = 2$);
277 substance abuse services ($n = 2$); veteran’s services ($n = 1$); primary school ($n = 1$); domestic abuse
278 shelter ($n = 1$); child advocacy centre ($n = 1$). Most studies targeted a single EBS aspect of EBS ($n =$
279 47), whilst some targeted multiple ($n = 4$). Burnout was by far the most common outcome targeted
280 targeted ($n = 41$) although STS ($n = 7$) and CF ($n = 8$) were also addressed. According to the EBS
281 model (Rauvola et al., 2019), this means that over 80% of identified studies were targeting one of the
282 adverse outcomes of empathy-based strain, rather than addressing empathy-based strain itself. Most



283 studies ($n = 46$) were classified as treatment studies, in that, as far as could be determined, they aimed
284 to examine the ability of an intervention to treat/impact upon EBS, with the assumption apparently
285 being that EBS was already present. Five studies aimed to both treat and prevent EBS. No studies
286 targeted prevention alone.

287

288

289

290

291

292

293

294

295

296

297

298

299

300

301

302

303 **Table 4**
 304 *Characteristics of Included Studies*
 305

ID	Study	Country	Publication Status	Study Design	EBS Construct Targeted	Setting	N	% of MHWs (specific role)	Gender	Intervention
01	Haynos et al., 2016	USA	PRJ	Quantitative Repeated measures	Burnout	CYP inpatient	22	100% (MH nurses)	Not Given	DBT skills coaching training
02	Roberts, 2018	UK	UT/D	Quantitative Repeated measures	Burnout	Not given	141 (39 at FU)	Not given	79.4% women	DBT skills training
03	Clarke et al., 2015	UK	PRJ	Quantitative RCT	Burnout	Inpatient, outpatient	140 (57 at FU 1; 36 FU 2).	100% (Staff working with clients with BPD)	Not given	ACT Training vs psycho-ed. BPD training ¹
04	Doyle et al., 2007	UK	PRJ	Quantitative Experimental design	Burnout	Forensic	26	61.50% (Forensic nurses)	73.1% women	Psychosocial intervention training
05	Hallberg, 1994	Sweden	PRJ	Mixed methods Repeated measures	Burnout	CYP inpatient	11	100% (Child MH nurses)	63.3% women	Systematic clinical supervision

¹ The primary aim of this RCT was to compare the effects of these two interventions on stigma towards clients with BPD; however clinician burnout was also measured

ID	Study	Country	Publication Status	Study Design	EBS Construct Targeted	Setting	N	% of MHWs (specific role)	Gender	Intervention
06	Hunnicutt & MacMillan, 1983	USA	PRJ	Quantitative Experimental design (2 intervention groups; 1 control)	Burnout	Community	251	Not given	Not given	Burnout workshop standalone vs with staff consultation.
07	Landis, 2010	USA	UT/D	Mixed methods Repeated measures	ST	Community	5	100% (SWs)	80% women	CF/ST workshop; 'Sharevision' meetings
08	Perseius et al., 2007	Sweden	PRJ	Mixed methods Repeated measures	Burnout	Adult & CYP psychiatry clinics	22	100% (Drs; psychol; MH nurses/care assistants; therapist, OT)	86.4% women	DBT training
09	Raab et al., 2015	Canada	PRJ	Quantitative Repeated measures	Burnout	Community	22	100 % (MH care workers)	100% women	MBSR
10	Razzaque & Wood, 2015	UK	PRJ	Mixed methods Repeated measures	Burnout	Community	26	100% (Psychr)	65.4% women	Mindfulness Retreat

ID	Study	Country	Publication Status	Study Design	EBS Construct Targeted	Setting	N	% of MHWs (specific role)	Gender	Intervention
11	Rollins et al., 2016	USA	PRJ	Quantitative Experimental design	Burnout	Inpatient & outpatient, (mostly veterans services)	145	100% (Behavioural health providers)	71% women	'BREATHE' burnout reduction workshop
12	Melchior et al., 1996	The Netherlands	PRJ	Quantitative Experimental design	Burnout	Inpatient	161	100% (MH nurses)	72% women	Primary nursing
13	Duckworth, 2012	USA	UT/D	Quantitative Repeated measures	STS	CYP community	16	100% (psychol; nurses; SWs, therapists; admin)	93.8% women	STS education program and prevention plan
14	Boone, 2006	USA	UT/D	Quantitative Experimental design	STS	Not given	97 (53 at FU)	100% (Domestic violence counsellors)	88.7% women	Poetry writing therapy
15	Flarity et al., 2016	USA	PRJ	Quantitative Repeated measures	CF	Inpatient (emergency department)	7	100 % (Forensic nurses)	100% women	CF prevention and resilience training
16	Suyi et al., 2017	Singapore	PRJ	Quantitative	Burnout	Not given	37	100%	81.1% women	MBSR

ID	Study	Country	Publication Status	Study Design	EBS Construct Targeted	Setting	N	% of MHWs (specific role)	Gender	Intervention
				Repeated measures				(Psychol; Psychr; SWs)		
17	Kovač et al., 2016	Slovenia	PRJ	Quantitative Experimental design	Burnout	School	30	100% (School counsellors)	100% women	Supervision (relational family model)
18	Van Kirk, 2020	USA	PRJ	Quantitative Repeated measures	Burnout, STS & CS	Veteran MH services	57	100% (nurses, SWs, psychol, psychr, counsellors, SA therapists; OTs; assistants)	64% women	Employee wellness program
19	Alenezi et al., 2019	Saudi Arabia	PRJ	Quantitative Experimental design	Burnout	Inpatient	296	100% (MH nurses)	49.3% women	Burnout prevention programme
20	Eriksson et al., 2018	Sweden	PRJ	Quantitative RCT	Burnout	Not given	101 (81 at FU)	100% (Psychol)	96% women (plus one NB person)	Web-based compassion program
21	Wymer, 2009	USA	UT/D	Quantitative	STS	Child advocacy centres	3	100%	100% women	Affective check-in supervision

ID	Study	Country	Publication Status	Study Design	EBS Construct Targeted	Setting	N	% of MHWs (specific role)	Gender	Intervention
				Single case research design				(CSA counsellors)		
22	Chilton et al., 2020	USA	PRJ	Qualitative Case study	CF & STS	Addiction treatment centre.	2	100% (art/expressive arts therapists)	100% women	El Duende ‘Process Painting’
23	Askey-Jones, 2018	UK	PRJ	Quantitative Repeated measures	Burnout	Primary & secondary	69 (43 at FU)	52% (MH Nurses)	76% women	MBCT group therapy
24	Ewers et al., 2001	UK	PRJ	Quantitative Experimental design	Burnout	secure forensic settings	33 (20 at FU)	100% (MH nurses)	51.5 % women	Psychosocial Intervention Training
25	Finamore et al., 2020	UK	PRJ	Quantitative Repeated measures	Burnout	Not given	253 (201 at FU)	100% Mixed	Not given	Knowledge & Understanding Framework PD Training
26	Reyes Ortega et al., 2019	Mexico	PRJ	Quantitative Experimental design	Burnout	BPD Clinic	6	100% (Psychol; psychr)	50% women	‘Helping the Helper’ social connectedness intervention
27	Brady et al., 2011	USA	PRJ	Quantitative	Burnout	Inpatient	16	100% (Psychr; Psychol; MH technicians)	Not given	MBSR

ID	Study	Country	Publication Status	Study Design	EBS Construct Targeted	Setting	N	% of MHWs (specific role)	Gender	Intervention
				Repeated measures				Activity Therapists)		
28	Walker, 2018	USA	UT/D	Quantitative	Burnout	Can't access-preview	34	100% (Can't access-preview)	Can't access-preview	MBSR
				Repeated measures						
29	Kiley et al., 2018	USA	PRJ	Quantitative RCT	CF	Community	69 (56 at FU)	100 (SWs; counsellors; psychr; CMs, support staff and management)	Not given	Guided Imagery
30	Riley et al., 2017	USA	PRJ	Quantitative 2 experimental conditions, no control	STS, CF & Burnout	Multisite MH centre	38 (28 at FU 1; 25 at FU 2; 19 at 6 month FU)	100% (Not given)	84.2% women	Yoga-based stress management vs cognitive-behavioural stress management
31	Rosada et al., 2015	USA	PRJ	Quantitative Repeated measures cross-over design	Burnout	Community	45	100 % (MH clinicians)	73.3% women	Reiki (reiki vs 'sham')

ID	Study	Country	Publication Status	Study Design	EBS Construct Targeted	Setting	N	% of MHWs (specific role)	Gender	Intervention
32	Ifrach & Miller, 2016	USA	PRJ	Quantitative Repeated measures	CF	Domestic abuse shelters	30	100% (DV counsellors)	100% women	Social action art therapy
33	Carmel et al., 2014	USA	PRJ	Quantitative Repeated measures	Burnout	Community	9	100% (MH practitioners; SA counsellors)	88% women	DBT training
34	Newman et al., 2015	South Africa	PRJ	Qualitative Retrospective	Burnout	CYP Community	30	100% (Child Psychr, psychr, Drs, nurses, clinical psychol, OTS, admins, interpreter)	70.6% women	Drumming
35	Paulson et al., 2015	USA	PRJ	Mixed methods Repeated measures	Burnout	Community (rural)	6	100% (psychol; counsellors)	Not given	Online peer consultation network (support group)
36	Luoma & Vilardaga, 2013	USA	PRJ	Quantitative Experimental design	Burnout	Community	20	80 (Psychol; Counsellors)	40% women	ACT Training
37	Little, 2000	USA	UT/D	Quantitative	Burnout	Not given	37	100%	75.7% women	DBT training

ID	Study	Country	Publication Status	Study Design	EBS Construct Targeted	Setting	N	% of MHWs (specific role)	Gender	Intervention
				Repeated measures				(Therapists; CMs; admins)		
38	Scarnera et al., 2006	Italy	PRJ	Quantitative	Burnout	Not given	25	100%	44% women	Interpersonal relationship management training
				Repeated measures pilot study				(Nurses, educators; psychol; psychr; managers)		
39	Ray, 1981	USA	UT/D	Quantitative	Burnout	Community	36	100%	75% women	Stress management training program
				Experimental design				(SWs; psychol; therapists; attendant; nurses; admin)		
40	Jones, 2009	UK	PRJ	Quantitative	Burnout	Inpatient	72 (62 at 3 rd FU, 49 at 4 th FU)	100%	57% women	Interventions for psychosis training + clinical practice development
				Repeated measures				(MH nurses; HCAs; OTs ward managers; CNSs)		
41	Redhead et al., 2011	UK	PRJ	Quantitative RCT	Burnout	Forensic inpatient	42	100	Not specified	Psychosocial intervention training
								(Forensic nurses)		
42	Salyers et al.,	USA	PRJ	Quantitative	Burnout	Substance abuse	84 (74 at FU)	Not given	87% women	BREATHE Burnout

ID	Study	Country	Publication Status	Study Design	EBS Construct Targeted	Setting	N	% of MHWs (specific role)	Gender	Intervention
	2011			Experimental design						reduction retreat
43	Weingardt et al., 2009	USA	PRJ	Quantitative RCT (2 experimental conditions, no control)	Burnout	Outpatient & controlled settings	147	100% (SA counsellors)	62.1% women	Online CBT training + supervision meetings (high vs low fidelity)
44	Leykin et al., 2011	USA	PRJ	Quantitative 2 experimental conditions, no control	Burnout	Community	149 (112 at 1 st FU 81 at 2 nd FU)	100% (SA counsellors)	62.6% women	Online training in CBT for substance abuse + supervision meetings (high vs low fidelity)
45	Gentry et al., 2004	USA	PRJ	Quantitative Repeated measures	Burnout & CF	Not given	83	100% (SWs; counsellors; psychols)	Not given	Certified CF Specialist Training
46	Hayes et al., 2004	USA	PRJ	Quantitative Experimental design (2 experimental conditions; 1 control)	Burnout	Not given	93	100 (SA counsellors)	63% women	ACT training vs Multicultural training

ID	Study	Country	Publication Status	Study Design	EBS Construct Targeted	Setting	N	% of MHWs (specific role)	Gender	Intervention
47	Anderson, 1982	USA	UT/D	Quantitative Experimental design	Burnout	MH centre	40	100% (counsellor-therapists)	60% women	Facilitator-led peer groups
48	Mehr et al., 1994	USA	PRJ	Quantitative Repeated measures	Burnout	Community	27	100% (‘MH workers’)	100% women	Stress-reduction/positive imagery conference + follow up meetings
49	Ballew 2020	USA	UT/D	Quantitative Quasi-experimental control time series design	CF	Not specified	43 (19 at FU)	68% (‘MH professionals’)	76.9% women (2 people identified with multiple genders)	Professional Resilience and Optimization workshop
50	Chochol et al., 2021	USA	PRJ	Mixed methods Repeated measures	Burnout	Child and adolescent psychiatry	6 (pilot study)	100% (Child and adolescent psychiatry fellows)	Not specified	Balint-like group incorporating brief emotional awareness modules

ID	Study	Country	Publication Status	Study Design	EBS Construct Targeted	Setting	N	% of MHWs (specific role)	Gender	Intervention
51	Bartels-Velthuis et al., 2020	Netherlands	PRJ	Quantitative Experimental design	CF	Outpatient MH clinic	47	100% (doctors; psychols; nurses; mindfulness teacher; SW; MH counsellor; physiotherapist; drama therapist)	91.5% women	Interpersonal Mindfulness Program + 45–60 min daily home practice

306 *Note:* ACT = acceptance and commitment therapy; BPD = borderline personality disorder; CF = compassion fatigue; CM = case managers; CNS =
307 clinical nurse specialist; CS = compassion satisfaction; CSA = child sexual abuse; CYP= children and young people; DBT = dialectical behaviour
308 therapy; FU = follow up; HCA= health care assistants; PD= personality disorder; PRJ= peer reviewed journal; MBCT = mindfulness-based cognitive
309 therapy; MBSR = mindfulness-based stress reduction; MH = mental health; MSc= masters NB= non-binary; OT= occupational therapist; PD =
310 personality disorder; psychr = psychiatrist; psychol = psychologist; RCT = randomised controlled trial; SA = substance abuse; ST= secondary
311 trauma; STS = secondary traumatic stress; SW = social worker; UM/T= unpublished masters/thesis **UT/D = University Thesis/Dissertation?**

312

313

314

315

316

317

318

319

320

Study	Ethnicity of Participants
Razzaque & Wood, 2015	White: 18(69 %) Black: 2(8 %) Asian: 3(12 %) 'Other' ethnic backgrounds: 2(8 %).
Rollins et al., 2016	White: 111(77%)
Duckworth McNeil, 2012	African American: 8 Caucasian: 7 'Other' ethnic backgrounds: 1
Boone, 2006	White; 44 (83%) Black: 4 (7.5%) Hispanic: 5(9.4%)
Suyi, Meredith & Khan, 2017	Chinese: 29 (78.4%) Malay: 2 (5.4%) Indian: 4 (10.8%)
Wymer 2009	African American/Black: 1(33.3%) Hispanic: 1(33.3%) White: 1(33.3%)
Chilton et al., 2020	100% white
Askey-Jones, 2018	95% White British (no information on other 5%)
Kiley et al., 2018	African American: 18(26.1%) White: 43 (62.3%) Multiracial: 3 (4.4%) 'Other' ethnic backgrounds: 1(1.4%) Missing data: 4 (5.8%)
Riley et al., 2017	94.7% White (no information on other 5.3%)
Rosada et al., 2015	White: 30 (66.7%) 'Non-white': 15 (33.3%)
Ifrach & Miller, 2016	Caucasian/White: 21(70%) Hispanic/Latina: 3(10%) African American/Black: 1(3.3%) Mixed race: 1(3.3%) 'Other' ethnic backgrounds: 4(13.3%)

Study	Ethnicity of Participants
Luoma & Vilardaga, 2013	White: 19(95%) Hispanic/Latino: 1(5%)
Jones, 2009	Black African: 55 % White British: 16 % African Caribbean: 9% Black British: 8% Mauritian: 7% Indian: 3% White African: 2%
Salyers et al., 2011	White: 62(79%) Black:15(19%) 'Other' ethnic backgrounds: 2(3%). Hispanic: 3(4%)
Weingardt et al., 2009	African American: 29 (19.7%) Asian/Pacific Islander: 1 (0.7%) White: 98 (65.3%) Hispanic/Latino: 12 (8.2%) 'Other' ethnic backgrounds: 9 (6.1%)
Leykin et al., 2011	Not given; however, sample reportedly the same as Weingardt et al. (2009), plus two further participants
Hayes et al., 2004	White: 84%
Mehr et al., 1994	Asian American: 14.7% Latina: 18.5% White: 66.7%
Ballew, 2020	Caucasian: 35 (87.5%) Multiracial: 3 (7.5%) American Indian: 1 (2.5%) Other: 1 (2.5%)

323

324 **Interventions**

325 Studies were included if they evaluated an intervention for reducing/preventing EBS that was externally
 326 facilitated (i.e. the organisation/researchers provide and arranged it). Studies measuring internally facilitated or
 327 'self-care' behaviours (or other individual attributes) were excluded on the basis that it is primarily an

328 organisational responsibility, not an individual one, to tackle EBS. Two included studies challenged these
329 criteria. Van Kirk's (2020) employee wellness program was organised and coordinated by the researchers,
330 however relied on employees to deliver the wellness interventions (a different one for every week of the
331 program) and took place during employee's lunch hour. Kiley et al.'s (2018) guided imagery study provided
332 participants with guided imagery tracks on MP3 players and instructed them to listen to them three times a week
333 over the intervention period, during work breaks. These studies were included because of the external
334 facilitation by researchers in guiding participants on what, when and how to perform the interventions. Similar
335 studies that provided a potential moderator of EBS without this structure were excluded, e.g., case study by
336 Callander et al. (2019) who gave the participant a mindfulness app but no guidance on when and how to use
337 this. However, it is important to draw a distinction between interventions that are externally facilitated (but still
338 focus on change at the individual level) and those that are truly organisational (i.e., involving true system
339 change).

340 Given the diverse range of interventions described, for clarity, they have been grouped according to the main
341 intervention components as described in the studies themselves (Table 6). It is acknowledged some of the more
342 integrative interventions contained various elements and could potentially be categorised multiple ways.

343 The length of the interventions varied considerably. The majority ($n = 36$) were delivered over weeks/months,
344 with timeframes ranging from two weeks to 12 months. Some studies were delivered in workshops/retreats of
345 one ($n = 3$), two ($n = 6$) and three ($n = 1$) days in duration, with one described as two to three days long. Three
346 interventions spanned less than a day. For studies where the total hours of intervention received could be
347 calculated, this ranged from two to 104 hours. Six studies required 'homework' activities in-between/in addition
348 to attending the interventions, including meditation practice and unfacilitated peer groups.

349

350

351

352 **Table 6**

Intervention	Description
Training in a therapeutic modality (N = 11)	The most common training was Dialectical Behaviour Therapy (DBT; $n = 5$), followed by Psychosocial Intervention Training (PIT; $n = 3$); Cognitive Behavioural Therapy (CBT; $n = 2$) and specific training on understanding and working with psychosis ($n = 1$) and the knowledge and understanding framework (KUF) for working with borderline personality disorder (BPD; $n = 1$; this was compared with an ACT intervention- see below).
EBS training and prevention/ Resiliency training (N = 7)	Participants were educated on causes, signs and impacts of EBS and encouraged to develop coping/prevention strategies. Three studies included opportunities for participants to practice some of these strategies e.g. breathing exercises; progressive muscle relaxation; developing secondary trauma narratives. The study delivering resiliency training incorporated EBS education and development of coping strategies, as well as teaching resiliency skills (e.g. self-regulation, self-care) described as ‘antibodies’ that could bolster participants’ ability to fight off compassion fatigue (Ballew, 2020). One study involved training MHWs to deliver the certified compassion fatigue specialist training (CCFST; Gentry & Baranowsky, 2004) to other MHWs experiencing CF (given the experiential elements of the training, it was hypothesised there would be a ‘training-as-treatment’ impact on participants’ own CF).
Mindfulness (N = 7)	Studies using mindfulness-based stress reduction (MBSR; $n = 4$); group mindfulness-based cognitive therapy (MBCT; $n = 1$) and a mindfulness-based professional development (MBPD; $n = 1$) retreat taught participants mindfulness theory and practice and encouraged workers to develop their own practices.
Wellness/stress reduction (N = 6)	Two studies involved training in stress management techniques, including identifying stressors and practicing stress-reduction skills, including guided imagery and progressive muscle relaxation. Additional wellness interventions included guided imagery ($n = 1$); reiki ($n = 1$) and a wellness program ($n = 1$) incorporating multiple facets of wellness (e.g. mindfulness, nutrition and movement). One study compared yoga-based stress management (YBSM; combining yoga and aspects of cognitive stress management) with cognitive-behavioural stress management (CBSM; Antoni et al., 2006).

Intervention	Description
Peer support/ relationships (<i>N</i> = 4)	Two studies involved the facilitation of regular peer groups: one focused on sharing thoughts and emotions regarding burnout issues; the other primarily on case consultation. One study used a functional psychotherapeutic approach to foster emotional closeness between MHWs and one taught participants techniques for handling professional interpersonal relationships.
Expressive arts (<i>N</i> = 4)	Interventions included art therapy (<i>n</i> = 1); art therapy supervision (<i>n</i> = 1); poetry therapy (<i>n</i> = 1) and drumming (<i>n</i> = 1). Art therapy supervision was included here (rather than in supervision, below) because creating art was the central component of the intervention.
Supervision (<i>N</i> = 3)	Studies included psychodynamic group supervision (<i>n</i> = 1); relational family supervision (<i>n</i> = 1) and affective check in supervision (<i>n</i> = 1). It should be noted that whilst two studies introduced supervision to MHWs who had not previously been receiving it, the affective check in study recruited participants already receiving regular supervision and trained their supervisors to use the affective check in technique.
Acceptance and Commitment Therapy (ACT; <i>N</i> = 3)	These studies taught MHWs ACT skills for personal use. Two studies compared ACT with other interventions including BPD training (see above) and multicultural training.
BREATHE (<i>N</i> = 2)	The BREATHE (Burnout Reduction: Enhanced Awareness, Tools, Hand-outs, and Education) program (Salyers et al., 2011) incorporates burnout prevention information with practices including mindfulness, social (e.g. support structures and boundaries), physical (e.g. body scan), cognitive (e.g. thought challenging; identifying values), imagery, and other self-care activities.
Primary nursing (<i>N</i> = 1)	A nursing intervention whereby both psychiatric and practical nurses were assigned to patients as primary nurse caregivers.
Balint-like group (<i>N</i> =1)	One group member volunteered to share a challenging case, after which group attendees asked clarifying questions. For approximately 10 min, the other group members and the group facilitators processed the case and their emotional responses. Afterwards, the presenter returned to the group and had the opportunity to respond to the group's commentary, stimulating further discussion until the end of the hour.

Intervention	Description
Compassion enhancement program (N = 1)	The program involves standard mindfulness exercises such as breathing anchor and body scans, and compassion-focused exercises such as loving-kindness, and exercises of compassion with self and others.



354 *Note:* ACT = acceptance and commitment therapy; BPD = borderline personality disorder; MHWs = mental
 355 health workers.

356
 357
 358
 359

360 **Theoretical Underpinnings and Assumptions**

361 There are at least two levels on which to consider the theoretical rationale for the interventions used by included
 362 studies to address EBS in MHWs. Firstly, there is the route via which studies have attempted to intervene and
 363 what this reveals about their underlying assumptions as to the cause and nature of EBS. Secondly, there are the
 364 theoretical frameworks/rationales underpinning some of the specific interventions used.

365 ***Route of Intervention***

366 We have described the routes of intervention taken by the included studies as as: 1. individual (personal
 367 practices; awareness-raising and role training); 2. team and 3. organisational.

368 **Individual.** Individual interventions intervened at the level of the individual to help them manage the
 369 impact of EBS and/or other demands associated with work. There were three subtypes of individual
 370 intervention: Personal resilience; awareness-raising and role-training.

371 **Personal resilience.** These interventions sought to provide MHWs with treatment and/or train them in
 372 using personal practices to manage their own negative experiences. These included: Wellness/stress reduction;
 373 mindfulness; ACT; compassion enhancement; expressive arts and aspects of the BREATHE intervention and
 374 resiliency workshops.

375 **Awareness-raising.** Awareness-raising interventions were those that attempted to alleviate EBS by
376 making MHWs aware of and prepared to tackle the negative effects of EBS. These included: EBS prevention
377 and aspects of the BREATHE intervention and resiliency workshops.

378 **Role-training.** Role-training interventions were those that provided MHWs with training in therapeutic
379 modalities or approaches to use with their clients. These included: training in a therapeutic modality.

380 **Team.** Team interventions were those focussed on team relationships and supportiveness. These
381 included the interventions: supervision; peer groups/relationships and Balint-like groups.

382 **Organisational.** Organisational interventions are those in which changes to the structure/running of the
383 organisations employing MHWs were made. These included: primary nursing.

384 Figure 3 displays how these levels of intervention intersect with the three types of organisational intervention for
385 improving staff wellbeing and mental health described by Tetrick and Quick (2011): i) Primary interventions aim
386 to decrease or remove stressors at the organisational level i.e. at the source. They require changes to workplace
387 practice and usually involve employees in the processing of developing interventions ; ii) secondary interventions
388 attempt to alter the individual's perception of, or responses to a stressor. They enable work-related stressors to be
389 swiftly identified and attempt to mitigate these by increasing employees' coping skills, awareness and knowledge
390 e.g. through additional training; iii) tertiary interventions are reactive and focus on rehabilitation of those already
391 experiencing significant strain resulting from stressors e.g. psychological therapy or occupational health services
392 (Tetrick and Quick, 2011). This demonstrates that all but one of the interventions were of the secondary type
393 although they were operating at various levels of individual and team intervention.

394 ***Specific Theories and Rationale***

395 Reference to theory was inconsistent, with some studies referring to an evidence base rather than theoretical
396 concepts. Where theoretical underpinnings for interventions were made explicit, these are highlighted below
397 (grouped by type of intervention). Evidence base/rational given for interventions not explicitly linked to theory
398 are summarised in Table 7.

399 **Psychosocial Intervention Training (PIT).** Clients with serious mental health difficulties may be
400 perceived as troublesome, hard to understand and difficult to help and these perceptions may make MHWs feel
401 demotivated (Jones & Lowe, 2001) and burnt out (Savicki & Cooley, 1987; Moore et al., 1992). PIT gives
402 MHWs skills to empathically understand and intervene more effectively with these clients, which may increase
403 their sense of reward and efficacy in their role and decrease burnout.

404 **EBS education & prevention/resiliency training.** One study (Duckworth McNeil, 2012) highlighted that
405 secondary traumatic stress disorder is thought to occur when second-hand trauma is not integrated (Harris,
406 1995). Thus, teaching MHWs about this may enable them to better respond to their experiences. Salyers et al.
407 (2011) combined burnout-reduction principles such as boundary setting with mindfulness techniques. For the
408 study teaching MHWs to deliver Certified Compassion Fatigue Specialist Training (CCFST); it was
409 hypothesised that that this may reduce their own CF due to exposure to theories of EBS they learnt in the
410 process (Gentry et al., 2004). The study utilising resiliency training delivered a workshop- the Professional
411 Resilience and Optimization workshop (Gentry & Monson, 2017)- which is based in the same recovery program
412 as the CCFST- the Accelerated Recovery Program (Gentry & Baranowsky, 1998). The Accelerated Recovery
413 Program is a manualised treatment for people experiencing compassion fatigue. The proposed effectiveness of
414 the Professional Resilience and Optimization workshop was linked to various separate mechanisms that did not
415 constitute a unified theory and are summarised in Table 7. However the study utilising this program made
416 reference to emotional contagion theory (Figley, 1995; Rothschild & Rand, 2006) and proposed that the
417 resiliency skills being taught would act as ‘antibodies’ of CF that would allow participants to resist its effects
418 (Ballew, 2020).

419 **Mindfulness/compassion.** Mindfulness and self-compassion encourage non-judgmental attitudes
420 towards experiences and may therefore decrease unhelpful coping strategies and increase willingness to accept
421 and experience negative emotions that might arise (Raab, 2014).

422 **ACT.** Studies referenced how key components of ACT e.g. psychological flexibility, cognitive de-
423 fusion and acceptance may reduce impact and believability of negative thoughts and feelings that arise from
424 difficulties/stress in working with clients (Hayes, 2004; Hayes et al., 2006).

425 **Supervision.** Studies theorised that supervision may decrease burnout as support systems positively
426 influence people's ability to cope with work stress (Pines & Aronsson, 1988; Wessells et al., 1989). It has also
427 been found to help counsellors manage stress and prevent burnout (e.g. Moyer, 2011; Thompson et al., 2011).
428 The study specifically using affective check-in supervision drew on literature suggesting that secondary trauma
429 responses in trauma counsellors could be mitigated by supervision practices that validate their experiences of
430 being personally impacted both cognitively and emotionally by their work as well as giving them skills to
431 manage this (e.g. Canfield, 2005; Knight, 2013; Whitfield & Kanter, 2014).

432 **Arts therapy.** Expressive arts therapies can alleviate compassion fatigue (Pearlman & Saakvitne,
433 1995) and their inclusion in MHWs' supervision can increase self-awareness and reduce stress (e.g. Deaver &
434 Shiflett, 2011; Fish, 2017). The study using drumming highlighted that this has been used in cultural healing
435 practices since the beginning of ancient history (Bittman et al., 2001) and found to have positive physiological,
436 psychological, and social effects and has been trialled on patients with mental health difficulties (e.g. Bensimon
437 et al., 2008). The study using poetry referenced how writing may help in the integration of traumatic memories
438 (Metcalf & Jacobs, 1996).

439 **Wellness/stress reduction.** This group of studies provided a diverse range of rationales for their
440 interventions which largely were not linked to explicit theories and are captured in Table 7. However, one stress
441 management study highlighted work by Holroyd (1979) suggesting that people can be trained to observe how
442 environmental stressors impact their behaviour and cognition and thus choose more adaptive responses to stress.

Intervention Type	Evidence Base
CBT (plus supervision)	The two studies training MHWs in CBT skills emphasised the format of training delivery and supervision above the content, i.e. highly structured, inflexible methods vs flexible and responsive training and supervision. They drew on research findings that indicate that supportive organizations that promote flexibility and autonomy may reduce job burnout (Garner et al., 2007; Knudsen et al., 2003); whereas highly centralised management practices are associated with higher levels of emotional exhaustion among counselling staff (Knudsen et al., 2006).
Primary nursing	No research explicitly investigated the impact of primary nursing (PN) on burnout. However, it was hypothesised that it might alleviate it given that low autonomy is associated with higher burnout in nurses (Reed, 1988), and PN increases autonomy (MacGuire & Botting, 1990).
Peer/relationships	Studies noted that social support is negatively correlated with burnout (Pines & Kafry, 1978) and that sharing one's emotions with others who are also experiencing difficulties can be therapeutic (Festinger, 1954; Schachter, 1968; Shubin, 1987). In addition, lack of reciprocity amongst colleagues and unhelpful comparisons with peers seem to have a part in the development of burnout (Schaufeli, 2003; Buunk et al., 2001; Bakker et al., 2000). The study utilising a social connectedness intervention based this on the Interpersonal Process Model (IPM; Reis & Shaver, 1988), wherein a turn-by-turn relational process is thought to establish psychological intimacy. They highlight evidence suggesting that when members of a dyad engage in reciprocal self-disclosure and respond to one another with care and validation, this creates feelings of closeness and intimacy between them (e.g., Laurenceau et al., 1998; Laurenceau et al., 2005).
DBT	Within the DBT model, the behaviours of clients with BPD that increase risk of burnout in MHWs (e.g. excessive contact with MHWs, demands for more regular therapy or threats towards MHWs) are considered therapy-interfering behaviours (Linehan, 1993). The model therefore incorporates a consultation group which aims to support and encourage MHWs, whilst maintaining their adherence to the model and identifying when clients are pushing boundaries so that this can be swiftly addressed. This is thought to reduce risk of burnout. In addition, emphasis on DBT clinicians practicing the core DBT skills of distress tolerance, mindfulness, emotion regulation, and interpersonal effectiveness is thought to be protective against burnout (Dimeff & Linehan, 2001; Linehan, 1993).

Intervention Type	Evidence Base	
Balint-like groups	The study utilising Balint-like groups cited research demonstrating that these groups target various outcomes including burnout and wellbeing (Van Roy et al. 2015). In addition it was noted that physicians have reported the groups decrease feelings of isolation and support their processing of emotional interactions (Schwartz et al., 2020) as well as increasing perceptions of social support for physicians working in palliative care (Popa-Velea et al., 2019).	
Wellness/stress reduction	Stress Reduction workshop + follow up meetings	The study employing stress-reduction/positive imagery workshops highlighted that stress-reduction workshops can alter the course of occupational burnout via integrative methods including practical guidance and emotional support (Pines & Aronson, 1988). However, they point out that workshops alone are insufficient and frequent follow up is needed to maintain the changes in coping styles that can be introduced (Edelwich & Brodsky, 1980).
	Wellness programme	This study suggested that the energy resulting from compassion fatigue can positively impact compassion satisfaction if handled differently Radley and Figley (2007). They cite findings that reduction in CF and gains in compassion satisfaction have been demonstrated for: Expressive writing (Henry, 2014), guided imagery (Kiley et al., 2018), yoga and mindfulness (Gregory, 2015), and music therapy (Hilliard, 2006).
	Yoga-Based Stress Management	The study authors had devised this novel method through combining yoga with aspects of cognitive stress management. They cite research that yoga and cognitive stress management programs are associated with improved in wellbeing and stress reduction (Granath et al., 2006). The authors also suggested the intervention was similar to MBSR and created opportunities to practice mindfulness (unreferenced).
	Guided Imagery	The study using guided imagery cited findings that imagining an activity produces similar physiological reactions to actually carrying it out (Morewedge et al., 2010; Pascual-Leone et al., 1995), and therefore suggested that calming and peaceful imagery may produce similar responses to real situations of tranquility.
	Reiki	The study using Reiki described how this energy-based treatment promotes energetic balance in recipients that holistically addresses emotional, physical and spiritual aspects of their self to balance and heal their energy (Mitchell, 1994).

448 *Note.* CBT= cognitive behavioural therapy; CF= compassion fatigue BPD = borderline personality disorder; DBT = dialectical behaviour therapy;
449 MBSR = mindfulness based stress reduction; MHWs – mental health workers; PN = primary nursing

450 **Measures**

451 An overview of the quantitative measures of EBS used in included studies is provided in
452 Appendix C. Variations of the Maslach Burnout Inventory were most commonly used (n =
453 28) consistent with the fact that burnout was the most common outcome targeted in the
454 included studies. Of the studies utilising qualitative/ mixed methods (n = 8), qualitative data
455 collection was done via interviews (n = 4); free text/semi-structured questionnaires (n = 3);
456 art created by participants (n = 1); focus groups (n = 1) and narrative feedback (n = 1).

457 The majority of studies (n = 49) collected EBS measures both pre and post intervention. Most
458 of these (n = 30) only collected post data at one time point. Of these 30 studies, 25 collected
459 post measures within two weeks of the intervention end. However, some (n = 6) collected
460 post measures between five weeks and 12 months post intervention. Four studies comprising
461 long-term interventions (supervision; DBT training) took multiple measures throughout
462 intervention phases. Twelve studies collected additional follow up data after the initial post
463 data, either once (n = 11) twice (n = 2) or three times (n = 1). Timeframes for additional post
464 data ranged from two to 18 months. Finally, one study (Rollins et al., 2016) collected their
465 first post-intervention data at six weeks following intervention, and then a further follow up
466 at six months.

467 **Discussion**

468 This scoping review is, to our knowledge, the first to describe the available evidence on all
469 available interventions for EBS in MHWs as a distinct professional group. In addition, it
470 aimed to consider the theoretical underpinnings for available interventions; determine how
471 EBS has been measured within included studies; and to make recommendations for future
472 research. The findings relating to each of these aims will be discussed below, with reference
473 to the wider literature. Overall, the group of studies identified in this review were
474 heterogeneous and differed substantially with regards to: the interventions used; the length

475 and delivery of interventions and the rationale for applying them. We discuss how the lack of
476 consistency across these areas, coupled with the conceptual difficulties in the field, create
477 issues for EBS intervention literature. In response, we put forth a research agenda with
478 recommendations for how future researchers might navigate and address these difficulties.

479 **Available interventions for EBS**

480 A total of 51 studies were identified, most of which (n=41) targeted burnout. Studies also
481 targeted STS and CF, though none targeted VT. The overwhelming focus on burnout is
482 notable; according to the EBS model proposed by Rauvola et al (2019), STS, CF and VT are
483 all forms of empathy-based strain that drive the process of EBS, whilst burnout is one of the
484 resulting adverse outcomes. Through the EBS lens, the majority of interventions described in
485 this review are therefore targeting one of the end results of EBS, rather than addressing the
486 three active constructs. Similarly, all studies aimed to address EBS already present in MHWs
487 (rather than solely aiming to prevent it). This suggests a recognition of the scale of the
488 problem in the mental health work force (NHS England, 2018; NHS, 2023), however is
489 further suggestive of an emphasis on cure over prevention

490 **Participants**

491 There was considerable variation in the specific roles/professions of MHWs included studies,
492 with nurses working in a mental health setting/capacity being the profession most
493 represented. Other professions included psychiatrists, psychologists, social workers,
494 expressive arts therapists, occupational therapists and health care assistants; thus the
495 collective sample of the review differed considerably with regards to the nature and amount
496 of training they would have had to complete their role.

497 Of further importance was participant ethnicity, which was reported by only 20 studies. Of
498 these, eight did not provide any information on participants who were not White, or referred



499 to them as ‘other’/ ‘non-white’. The omission and/or inadequate description of participant
500 ethnicity is significant in light of research demonstrating that NHS staff belonging to ethnic
501 minorities (in the context of the population of the UK) may face additional stress in the
502 workplace. The NHS staff survey (NHS, 2023a) showed that one in five staff from minority
503 ethnic groups (other than white minority groups) experienced discrimination in the 12 months
504 preceding the survey. The latest version of the Workforce Race Equality Standard (WRES)
505 report showed that only 44.4% of Black and minority ethnic (BME²) NHS staff felt that their
506 organisation offered equal opportunities for career progression; compared with 58.7% of
507 White staff (NHS England, 2023b). Ethnic minority NHS staff have reported in interview
508 data the racism and racial microaggressions they face at work and the lack of equal
509 opportunities in their professional roles (Ross et al., 2020). Thus, there is substantial evidence
510 that MHWs, along with other NHS staff, may face additional workplace stress and potentially
511 even racial trauma as a result of belonging to an ethnic minority. If organisations and
512 researchers are not monitoring ethnic diversity when designing and implementing EBS
513 interventions, then these are unlikely to be sufficiently inclusive or sensitive to the additional
514 stressors faced by MHWs in minority ethnic groups.

515 **Theoretical underpinnings and assumptions**

516 There was significant variation in methodology, suggesting little consensus about how
517 research exploring EBS should be approached. To better summarise the available elements,
518 studies were grouped into three categories based on the level at which they sought to
519 intervene, namely individual (including personal resilience; awareness-raising and role
520 training); team and organisational approaches to addressing EBS. To contextualise these

² It is acknowledged that there is a helpful ongoing debate regarding the use of the term BME (see Aspinall, 2020 for a critical review), which is not widely accepted by the groups it represents yet continues to be used. This review uses the above term only in order to refer to data from existing reports which have used it.

521 findings, these levels were considered alongside the three types of organisational
522 interventions identified by Tetrick and Quick (2011).

523 This comparison demonstrates that all but one of the interventions were of the secondary type
524 (Tetrick & Quick, 2011) although they were operating at various levels of individual and
525 team intervention. The only intervention possibly acting at the organisational level and of
526 primary type (Tetrick & Quick, 2011) was the primary nursing study by Melchior et al.
527 (1996). Primary nursing is a model of nursing care delivery where each patient's care is the
528 responsibility of one nurse, with care focusing on the needs of the patient rather than the ward
529 (Gardner, 1991). Therefore, whilst this may be applicable to inpatient psychiatric settings, it
530 is not widely applicable to MHWs more generally.

531 In terms of the rationale for specific interventions, several studies provide an evidence base
532 rather than an explicit theoretical rationale. These studies generally noted EBS is negatively
533 correlated with: flexibility and autonomy (flexible CBT format; primary nursing); social
534 support and emotional reciprocity with colleagues (peer support/relationships) and
535 therapeutic boundaries (DBT training). Aside from Van Kirk (2020), studies utilising
536 wellness/stress reduction interventions did not cite research specifically on EBS; however,
537 they highlighted evidence of the positive effects of these interventions on stress and
538 wellbeing in general.

539 Where theoretical rationales were provided, a common thread appeared to be teaching
540 MHWs to respond differently to the EBS inducing aspects of their jobs. For ACT,
541 mindfulness and compassion-based interventions, this took the form of taking accepting and
542 non-judgemental attitudes towards negative thoughts and feelings that might arise from work.
543 For EBS prevention and education, the emphasis was on teaching MHWs the causes and
544 signs of EBS so they could better respond to negative experiences. For art therapy, the
545 emphasis was on resolving mental health issues and integrating traumatic experiences

546 through various therapeutic art forms. PIT, supervision and some aspects of EBS-awareness
547 (e.g. boundary setting) were the only interventions that suggest a preventative element.
548 However, the onus was still on the MHW to change an aspect of their practice or complete an
549 additional task. This seems consistent with the perspective of Montgomery (2014), who
550 suggested healthcare systems which are still dominated by the medical model take a
551 pathogenic perspective on burnout as a problem to be ‘treated’ at the individual level, rather
552 than questioning why the organisations consistently produce this issue.

553 Such a focus on the individual echoes findings from Cocker and Joss’ (2016) review of
554 interventions to reduce CF in healthcare, emergency and community services workers which
555 found all 13 included studies had an individual focus, and the majority employed stress
556 reduction and/or holistic interventions (e.g. yoga, meditation). Several reviews of EBS
557 interventions for healthcare staff have focused only on mindfulness-based interventions (e.g.
558 Conversano et al., 2020; Gilmartin et al., 2017; Klein et al., 2020; Mensah & Anderson,
559 2015). Thus, the wider literature seems to share the emphasis on individual interventions to
560 EBS that were found in this review.

561 However, studies of factors influencing EBS suggest organisational elements are key. For
562 example, Singh et al. (2020) demonstrated job-related factors including work environment,
563 workload and workplace trauma were associated with CF in MHWs. One of the most
564 consistently highlighted predictors of EBS across healthcare providers is high caseload/client
565 contact (Lasalvia et al., 2009; Maslach et al., 2001; Turgoose and Maddox, 2017; Singh et al.,
566 2020). In a systematic review of burnout determinants, O’Connor et al. (2018) concluded that
567 reasonable caseloads, clinician autonomy, good team functioning, and proper supervision
568 should be the focus of organisational attempts to prevent and reduce burnout in MHWs.
569 Sutton et al (2022) meanwhile, found in their narrative synthesis of organisational factors that
570 ameliorate EBS in MHWs that regular supervision, balanced and diverse caseloads, strong

571 peer support networks and an organisational culture that acknowledges secondary trauma
572 were key.

573 Furthermore, authors in this field have emphasised that organisations employing MHWs have
574 a responsibility to prevent and reduce EBS (Abendroth & Figley, 2013) and warn that failure
575 to address the systemic nature of this problem is resulting in significant clinician distress
576 (Dean et al., 2019). As Killian (2008) noted upon finding no significant relationship between
577 individual self-care strategies and reported levels of EBS, perhaps we should “stop expecting
578 helping professionals to “pull themselves up by their bootstraps” by reducing their stress with
579 standard individual coping strategies of leisure and continuing education” (pp. 42). They
580 instead called on organisations to protect the wellbeing of MHWs by altering their workloads
581 and giving them greater autonomy.

582 Also of note is the theory-practice gap between the theorised causes of EBS and the majority
583 of interventions used to address them. STS and CF are thought to arise through traumatic
584 exposure to the distress of others (Figley, 1995) and burnout is said to develop due to
585 repeated interpersonal stress (Maslach, 1993). However, very few of the studies reviewed
586 here addressed these issues directly. Of the seven studies intending to target STS, only three
587 (Boone, 2006; Landis, 2010; Wymer, 2009) mentioned working with trauma responses.
588 Despite the interpersonal mechanisms indicated in all descriptions of EBS; most studies
589 utilised individual interventions, and the interventions that had relational factors centred on
590 relationships with colleagues and supervisors, not clients (the supposed source of the EBS). It
591 is therefore unclear how these interventions are proposing to address the root of the problem.

592 As described above, it is challenging to cohesively summarise the varied interventions
593 outlined in this review. One lens through which to view these is the Job Demands- Resources
594 (JD- R; Bakker & Demerouti, 2007) model. The JD-R assumes that all work inherently
595 consists of job demands (JDs) and job resources (JRs). JDs are effortful, psychologically or

596 physically costly aspects of a job that are negatively valued; whilst JRs are rewarding,
597 positively valued aspects of a job that offer personal development or mitigate the impact of
598 JDs (Schaufeli & Taris, 2014). High JDs and insufficient JRs predict burnout; however, JDs
599 may not increase burnout if workers also have access to resources such as autonomy and
600 high- quality relationships with supervisors (Bakker et al., 2005). Using this framework,
601 most of the included studies seem to be aiming to increase JRs, either through increasing the
602 EBS-awareness, therapeutic skill or personal resources of individual MHWs, or through
603 team-based efforts i.e. supervision. However a recent study found that both lowering JDs and
604 increasing JRs is necessary to enable MHWs to provide high quality, person-centred care and
605 recommends mental health organisations address both (Fukui et al., 2021).

606 **Measurement of EBS**

607 Whilst repeated measures designs were the most common, there was notable inconsistency in
608 methodologies, as well as in the duration of interventions and the timing/frequency of follow
609 ups. The Maslach Burnout Inventory was the most common measure used which is consistent
610 with the majority of studies targeting burnout as a construct. Qualitative/mixed methods were
611 rare, and there appears to have been minimal attempts to discover what factors were
612 associated with EBS for the study populations before intervening. This may be significant
613 given the diverse roles of MHWs samples and multiple settings featured in included studies.
614 Bakker and Demerouti (2007) describe using a two-stage process in their research into the
615 unique burnout risks in different jobs/settings. This begins with qualitative interviews with
616 workers within the organisation, exploring the unique demands and resources inherent in
617 their role. The second phase involves operationalising the specific jobs demands and
618 resources into items in a custom-made questionnaire which is then completed by all workers.
619 This allows for organisation-specific interventions to be implemented when seeking to
620 address burnout. Employing similar methods with MHWs may be useful , and indeed there
621 are examples to be found in the literature e.g. Wilkie et al (2022) report on a two year process

622 of developing and implementing wellness initiatives in a Canadian hospital wherein survey
623 and consultation processes with staff led to individual, team and organisational changes (this
624 paper is not reviewed here as the initiatives have yet to be evaluated).

625 It is beyond the remit of a scoping review to assess and compare effectiveness of
626 interventions. However, it is worth noting that even superficial observations about the relative
627 outcomes of different approaches were difficult to make due to: the amount of different
628 interventions used; the multi-component nature of many of the interventions and the tendency
629 for studies to report changes across multiple subscales of EBS measures.

630 **Limitations**

631 The literature regarding how CF, burnout, VT and STS are conceptualised is inconsistent,
632 which may hamper the usefulness of the findings to some extent. Ravoula et al.'s (2019)
633 concept of EBS was used by this review in an attempt to overcome this conceptual
634 inconsistency. Whilst this enabled an inclusive search strategy; it is possible that
635 amalgamating concepts that have related however not identical mechanisms constituted an
636 over-simplification. Nonetheless, over-specificity may also be unhelpful. In a review that
637 solely targeted CF, Cocker and Joss (2016) found their earliest study published in 2011 and
638 concluded that the evidence base for these interventions is relatively recent. By including all
639 EBS concepts, the current review spans 40 years of research and arguably captures a more
640 representative view of attempts to help MHWs manage the impact of empathically
641 demanding work, independent of changing conventions in terminology.

642 A further limitation was excluding studies in languages other than English, thus this review
643 has an inherent bias towards evidence produced in Western, English-speaking countries and
644 cultures. However, the search strategy took a broad approach to publication status,
645 methodology, and type of intervention whilst retaining a specific focus on MHWs.

646

Research Agenda

647 Drawing together the findings of this review with the wider literature, we have produced a
648 research agenda to address the current issues in this field of study and drive it forward in a
649 manner that focuses predominantly on the urgent need for effective EBS interventions in
650 mental health services.

651 Current Issues

652 We highlight several key issues with existing research, including:

- 653 1) Inconsistent terminology/ lack of conceptual clarity, which creates unhelpful
654 partitions between similar areas of research
- 655 2) A disconnect between the proposed causal mechanisms of EBS and the interventions
656 being chosen to address it
- 657 3) An emphasis on individually focused interventions, which is at odds with literature
658 showing organisational factors are the leading cause of EBS
- 659 4) An emphasis on treating the outcomes of EBS rather than addressing or preventing
660 the causes
- 661 5) A failure to identify from MHWs themselves what the sources of EBS are in their
662 specific organisations before implementing preventatives or treatments
- 663 6) Potentially relevant clinical audits and initiatives being carried out in services and not
664 widely disseminated beyond the organisation, resulting in a disconnect between
665 knowledge held in clinical and academic arenas
- 666 7) Little regard given to diversity factors which may influence or interact with EBS, and
667 poor reporting on diversity data such as ethnicity
- 668 8) Methodological issues including inconsistency in study design/rigor, inconsistent
669 measurement of EBS and lack of control for profession of MHW

670 **Suggested Solutions**

671 1) **Inconsistent terminology/ lack of conceptual clarity**

672 **Adopting and Updating the EBS Model.** Greater conceptual agreement and consistency in
673 the literature regarding EBS would be helpful to avoid duplication of effort and bring
674 together inter-related fields of knowledge and research. We suggest that adopting Rauvola et
675 al.'s (2019) EBS model offers a practical solution to the problem of inconsistent
676 nomenclature in the field by distilling the core elements of the various concepts available in a
677 dynamic process model.

678 We note that the EBS model was born out of a conceptual review and hope that by
679 considering it in an applied context (i.e. how to helpfully intervene in EBS) we can support
680 its translation into the clinical research environment. Based on the current review we
681 therefore suggest one minor amendment. The model depicted in Rauvola et al. (2019) shows
682 that both individual and contextual factors influence the onset of EBS. Examples of what
683 constitutes contextual factors are given as 'emotional display norms & expectations,
684 form/frequency of trauma exposure, support' (pp. 298). Whilst the authors' intended meaning
685 may have been to clearly separate the two, we suggest the wording is potentially ambiguous
686 and may not be clear enough in emphasising organisational contextual factors related to the
687 workplace, rather than personal contextual factors. For example, 'support' and 'trauma
688 exposure' could refer to these elements both in and outside of the workplace. We therefore
689 suggest a clearer distinction between organisational/job role context (e.g. workload, role
690 autonomy, degree of exposure to EBS via work role) and individual factors (e.g. trauma
691 history, coping style, personality, support network outside work).

692 **Testing the EBS Model.** As Rauvola et al. (2019) note, empirical testing of this model is
693 warranted e.g. identifying underlying mechanisms, determining what factors may trigger
694 EBS, mapping how it arises over time. Additionally, determining the relative contribution of

695 contextual and individual factors is pertinent in light of the split described in this review
696 between organisation-related causes of EBS and individually-focused interventions. E.g. how
697 does the use of self-rostering systems (organisational context) to address EBS interact with
698 caring status (individual context)?

699 **Differentiating concepts and updating terminology.** Following the testing processes
700 described above, further clarification and differentiation of terms should be considered. It
701 may be helpful to further disentangle the three empathy-based strain constructs (STS, VT,
702 CF) from the concept of burnout (a proposed outcome of empathy-based strain- however
703 acknowledged by Rauvola et al to share common features). Other longstanding terminology
704 incorporated in the EBS model would benefit from review considering more recent empirical
705 work. For example, a neuroscientific study by Hofmeyer et al (2020) has contested the term
706 compassion fatigue after finding that compassion does not cause fatigue- the authors suggest
707 that ‘empathic distress fatigue’ should be used instead.

708 **The Need for Practicality and Action.** Whilst consistent conceptualisations and labels are
709 undeniably important when designing research and testing hypotheses, we note that the field
710 has been plagued by conceptual disagreements for decades and acknowledge these are
711 unlikely to be solved to the satisfaction of all. We therefore advocate a practical approach.
712 What seems evident from this and other reviews is that there can be negative consequences
713 arising from repeated empathic engagement with others’ distress and that this, whatever name
714 is given to it, translates into harmful outcomes for MHWs and clients alike. We encourage
715 applied, organisational and structured attempts to arrive at solutions to this chronic problem,
716 which has been made ever more pressing by the COVID-19 pandemic.

717 2) **Disconnect between the causes of EBS and the interventions used to address it**

718 **Clearly Linking Interventions to Causes.** We call for interventions that are theoretically
719 and conceptually linked to the theoretical understanding of EBS. If studies are using

720 individual concepts such as burnout, compassion fatigue etc then there should be clear links
721 between the assumed mechanisms of these conditions (see Table 1) and the steps being taken
722 to reduce or prevent them., and which take account of the dynamic interplay of the various
723 components of the model.

724 3) **Organisational Causes vs Individual Solutions**

725 **More Organisational Interventions.** Existing literature suggests organisational factors are
726 the key drivers of EBS, therefore there is a need for studies which employ
727 organisational/primary interventions for EBS in MHWs e.g. reduced/altered caseloads,
728 greater autonomy for staff. In calling for research of this nature, we are following the
729 example of West, Bailey and Willias (2020) who gave primary interventions the main focus
730 in their independent review of how to support nursing and midwives in delivering high
731 quality care. They describe several examples of primary interventions that have been used in
732 NHS mental health trusts. For example, Langley Green acute mental health hospital (Sussex,
733 UK) implemented a distributed leadership model under which staff could raise issues and
734 implement changes to the service. In addition, supervision commitments were reliably
735 increased to over 90% fulfilment per week and monthly ‘transparency boards’ monitored
736 levels of staff training and supervision. This model led to a decrease in staff sickness over the
737 course of implementation and medication omissions fell from 36% to 0%, illustrating the dual
738 benefits to staff and patients alike. East London NHS Foundation Trust (mental health and
739 community services) reviewed all their clinical audit processes with the involvement of
740 service users and staff and ultimately ceased 85% of all audit activity. This triggered two
741 broader initiatives to: i) identify any activities that did not add value to services (e.g.
742 duplication of clinical recording) which the trust then acted to address; ii) commence a ‘break
743 the rules’ campaign, encouraging staff to highlight pointless procedural rules that could be
744 eradicated. Whilst these were not research studies, they illustrate that organisational
745 interventions do exist and would benefit from robust evaluation.

746 A further reason for increasing organisational interventions is suggested by the Job Demands-
747 Resources model (JDR; Bakker & Demerouti, 2007), which tells us that job stress and
748 burnout arise out of an imbalance between both job demands and job resources. As noted
749 above, the majority of existing studies are addressing EBS by providing resources rather than
750 focusing on addressing the demands faced by MHWs. Therefore, research that addresses
751 these inherently organisational demands is warranted.

752 4) **An emphasis on treating the outcomes of EBS rather than addressing or**
753 **preventing the causes**

754 **Focus On Prevention.** It is easier to prevent EBS than to cure it once it has been established.
755 As noted by NHS Employers (2022), organisations employing MHWs should aim to reduce
756 the drivers of EBS and equip themselves to recognise and intervene early when signs of EBS
757 emerge. This focus on prevention should be reflected in the research being carried out in this
758 field.

759 5) **Failure to Ask MHWs About Sources of EBS**

760 **Ask First, Intervene Second.** Existing research suggests there are key recurring
761 organisational factors that can lead to EBS, with high caseloads being the most consistent
762 (e.g. Singh et al., 2020). Whilst this suggests there may be reliable trends across services, it is
763 expected that the relevant EBS-causing factors will vary between different organisations
764 which differ in terms of e.g. structure, remit and funding. When developing interventions to
765 address EBS in MHWs, a suggested preliminary step is for researchers to consult with
766 MHWs to gain an understanding of e.g. the sources/causes of EBS in their particular
767 organisation/setting (as did the East London trust leaders in the example given in West et al.,
768 2020), and design interventions accordingly. The two-step process described by Bakker and
769 Demerouti (2007; see discussion section) is a further example of using specific staff concerns
770 to generate outcome measures for establishing an interventions success.

771 6) **Disconnect between Clinical and Academic Domains**

772 **Practice-informed Research.** Whilst the in-house clinical initiatives and audits described in
773 West et al. (2020) may lack specific outcome measures relevant to EBS, they highlight that
774 organisational and primary interventions are happening. However, such data is not typically
775 widely disseminated beyond the trust or organisation in such a way that it could be identified
776 by this and other similar reviews or stimulate further research. We therefore recommend
777 greater connection and collaboration between clinical and academic organisations seeking to
778 tackle the issue of EBS. It is suggested that such collaboration could provide greater benefits
779 for the clinical staff and services participating in studies (e.g. greater chance of having
780 context specific and relevant issues addressed); the research teams themselves (e.g. help to
781 identify meaningful targets for outcome monitoring ; ideas for intervention) and the quality
782 and usefulness of the resultant findings (e.g. greater validity and relevance of findings). At
783 the least, we advocate that further research in this area be informed by e.g., outstanding case
784 examples, to promote the development of robust real-world research.

785 7) **Insufficient Attention Given to Diversity Factors**

786 **Record and Control for Diversity.** Researchers and services must attend to and address the
787 additional stressors that may be faced by MHWs who belong to ethnic minorities (e.g. NHS,
788 2023a) when designing interventions for EBS e.g. by using additional measures that capture
789 experiences of discrimination. Furthermore, they must record and control for a wide range of
790 diversity factors when implementing and evaluating EBS interventions, to ensure these are
791 applicable across staff who represent a diverse workforce in terms of factors including
792 ethnicity, race, gender, gender identity, sexuality, ability.

793 8) **Methodological issues**

794 **Higher Quality, Consistent Methodology.** Most of the studies included in this review
795 themselves made recommendations for future research (see Appendix D for a summary). The
796 most consistent of these was for greater methodological rigor i.e., controlled, experimental
797 designs and larger samples. We also note that in order for future reviews to draw conclusions
798 about the relative effectiveness of different EBS interventions, there is a need for greater
799 methodological consistency across studies.

800 **Improving Outcome Measurement.** The second most commonly suggested research target
801 from included papers was for studies to have a longer duration of intervention and/or follow
802 up prior to measurement. Several studies in the review also questioned whether changes in
803 EBS for MHWs would lead to improved outcomes for clients and recommended this be
804 measured. We highlight the wide variation in specific outcome measures used and suggest
805 greater consistency in measurement tools to allow direct comparison of studies.

806 **Controlling for Specific Role.** Several studies in the review acknowledged the diverse
807 professions and settings that MHWs may represent and suggested this be controlled for in
808 future studies- a recommendation we echo.

809 **Summary Recommendations**

810 For ease of reference, we have summarised the above into bullet points intended to guide
811 researchers undertaking further studies into interventions for EBS in MHWs. Future studies
812 should:

- 813 • Prioritise applied intervention research over conceptual debate.
- 814 • Consider adopting Rauvola et al.'s (2019) EBS model to overcome conceptual
815 discrepancies, including our suggestion to ensure that the 'contextual factors' aspect
816 of the model refers specifically to organisational context.

- 817 • Carry out empirical testing of the EBS framework (if using this) and clarify
818 terminology accordingly
- 819 • Regardless of EBS construct being targeted, ensure that the chosen intervention is
820 theoretically and conceptually linked with the proposed cause of the problem.
- 821 • Prioritise organisational or primary interventions (e.g. caseloads, service structure,
822 worker autonomy) over those that target individual MHW qualities or skills.
- 823 • Design studies that reduce job demands rather than solely attempting to provide job
824 resources.
- 825 • Prioritise studies that focus on preventing EBS rather than seeking to cure it once it
826 has been established.
- 827 • Find out from MHWs what the sources of EBS are in their particular setting before
828 intervening.
- 829 • Collaborate with services and organisations employing MHWs to share knowledge
830 and data between the clinical and academic realms. Be informed by outstanding case
831 examples and examples of positive deviance in clinical practice.
- 832 • Measure and control for a range of diversity factors in participants, to ensure
833 interventions are applicable to diverse MHWs. Consider how diversity issues such as
834 institutional racism may interact with EBS and introduce additional measures to
835 evaluate this.
- 836 • Design larger, more methodologically robust experimental studies that control for the
837 setting and specific role of MHWs.
- 838 • Consider longer periods of intervention and follow up prior to/over the course of
839 outcome measurement.
- 840 • Consistency in the EBS measures used to allow better comparison between studies.
- 841 • Consider measuring the outcomes of EBS interventions for MHWs in both MHWs
842 and their clients in parallel.

843

Conclusion

844 This scoping review outlined the current state of the literature regarding interventions for
845 EBS in MHWs. Overall, the findings showed that there is little consistency across available
846 interventions; aside from the general trend towards intervening at the level of the individual.
847 Given the additional pressures placed on mental health services by the COVID-19 pandemic,
848 we have questioned whether this individual focus is helpful or sufficient in the context of
849 ever-increasing need for the services MHWs provide. Finally, we have presented a research
850 agenda detailing how these and other issues can be addressed by future researchers studying
851 interventions for EBS in MHWs. Overall, this emphasises the need for practicality and
852 encourages applied interventions focusing on organisational causes of EBS.

853

References

854 *indicates studies included in the review

855 Abendroth, M., Figley, C.R. (2013). Vicarious Trauma and the Therapeutic Relationship. In
856 Murphy, D., Joseph, S., Harris, B. (Eds.) Trauma and the Therapeutic Relationship:
857 Approaches to Process and Practice (pp. 111-125). London: Palgrave Macmillan.

858 Ahmedani, B.K. (2011). Mental Health Stigma: Society, Individuals, and the Profession.
859 Journal of Social Work and Values Ethics. 8(2), 4-1–4-16.

860 *Alenezi, A., McAndrew, S., Fallon, P. (2019). Burning out physical and emotional fatigue:
861 evaluating the effects of a programme aimed at reducing burnout among mental health
862 nurses. International Journal of Mental Health Nursing. 28(5): 1042- 1052. [https://doi-](https://doi-org.ezproxy1.bath.ac.uk/10.1111/inm.12608)
863 [org.ezproxy1.bath.ac.uk/10.1111/inm.12608](https://doi-org.ezproxy1.bath.ac.uk/10.1111/inm.12608)

864 Al-Turki, H. A., Al-Turki, R. A., Al-Dardas, H. A., Al-Gazal, M. R., Al-Maghrabi, G. H., Al-
865 Enizi, N. H., & Ghareeb, B. A. (2010). Burnout syndrome among multinational nurses

866 working in Saudi Arabia. *Annals of African medicine*, 9(4), 226–229. [https://doi-](https://doi-org.ezproxy1.bath.ac.uk/10.4103/1596-3519.70960)
867 [org.ezproxy1.bath.ac.uk/10.4103/1596-3519.70960](https://doi-org.ezproxy1.bath.ac.uk/10.4103/1596-3519.70960)

868 *Anderson, C. M. (1982). The effects of peer support groups on levels of burnout among
869 mental health professionals. *Dissertation Abstracts International Section A: Humanities and*
870 *Social Sciences*, 43(6-A), 1880–1881.

871 Arksey, H. & O'Malley, L. (2005). Scoping studies: towards a
872 methodological framework, *International Journal of Social Research Methodology*, 8(1), 19-
873 32. <https://doi.org/10.1080/1364557032000119616>

874 *Askey-Jones R. (2018). Mindfulness-based cognitive therapy: An efficacy study for mental
875 health care staff. *Journal of psychiatric and mental health nursing*, 25(7), 380–389.
876 <https://doi-org.ezproxy1.bath.ac.uk/10.1111/jpm.12472>

877 Aspinall, P.J. (2020) Ethnic/Racial Terminology as a Form of Representation: A Critical
878 Review of the Lexicon of Collective and Specific Terms in Use in Britain. *Genealogy*, 4, 87.
879 <https://doi.org/10.3390/genealogy4030087>

880 Atkins, P.W.B., & Parker, S.K. (2012). Understanding individual compassion in
881 organizations: The role of appraisals and psychological flexibility. *Academy of Management*
882 *Review*, 37(4), 524–46. <https://doi.org/10.5465/amr.2010.0490>

883 Bakker, A. B., & Demerouti, E. (2007). The Job Demands-Resources model: State of the art.
884 *Journal of Managerial Psychology*, 22(3), 309–328.
885 <https://doi.org/10.1108/02683940710733115>

886 Bakker, A. B., Demerouti, E., & Euwema, M. (2005). Job resources buffer the impact of job
887 demands on burnout. *Journal of occupational health psychology*, 10(2), 170–180. [https://doi-](https://doi-org.ezproxy1.bath.ac.uk/10.1037/1076-8998.10.2.170)
888 [org.ezproxy1.bath.ac.uk/10.1037/1076-8998.10.2.170](https://doi-org.ezproxy1.bath.ac.uk/10.1037/1076-8998.10.2.170)

889 * Ballew, J.K. (2020) The effects of resiliency training on self-reported compassion fatigue
890 and compassion satisfaction in mental health professionals and counselors-in-training
891 [Unpublished thesis] University of Montana

892 Baranowsky, A. (2002). The silencing response in clinical practice: on the road to dialogue.
893 In C.R. Figley (Ed.), *Treating Compassion Fatigue*, Brunner/Mazel, New York.

894 *Bartels-Velthuis, A.A., van den Brink, E., Koster, F., Rogier Hoenders, H. J. (2020). The
895 Interpersonal Mindfulness Program for Health Care Professionals: a Feasibility Study.
896 *Mindfulness* 11, 2629–2638. <https://doi.org/10.1007/s12671-020-01477-5>

897 Beaton, R. D., & Murphy, S. A. (1995). Working with people in crisis: Research
898 implications. In C. R. Figley (Ed.), *Brunner/Mazel psychological stress series*, No. 23.
899 *Compassion fatigue: Coping with secondary traumatic stress disorder in those who treat the*
900 *traumatized* (p. 51–81). Brunner/Mazel.

901 Bensimon, M., Amir, D., & Wolf, Y. (2008). Drumming through trauma: Music therapy with
902 post-traumatic soldiers. *The Arts in Psychotherapy*, 35, 34–48.
903 <https://doi.org/10.1016/j.aip.2007.09.002>

904 Bittman, B. B., Berk, L. S., Felten, D. L., Westergard, J., Simonton, O. C., Pappas, J., &
905 Ninehouser, M. (2001). Composite effects of group drumming music therapy on modulation
906 of neuroendocrine-immune parameters in normal subjects. *Alternative Therapies*, 7, 38–47.

907 *Boone, B.C. (2012). The impact of poetry therapy on symptoms of secondary posttraumatic
908 stress disorder in domestic violence counsellors [Unpublished thesis] Office of Graduate
909 Studies, Texas A&M University.

910 Borritz, M., Rugulies, R., Bjorner, J. B., Villadsen, E., Mikkelsen, O. A., & Kristensen, T. S.
911 (2006). Burnout among employees in human service work: design and baseline findings of

912 the PUMA study. *Scandinavian journal of public health*, 34(1), 49–58.
913 <https://doi.org/10.1080/14034940510032275>

914 Bride, B., Radey, M., Figley, C. (2007). Measuring Compassion Fatigue. *Clinical Social*
915 *Work Journal*. 35, 155–163.

916 *Brady, S., O'Connor, N., Burgermeister, D., & Hanson, P. (2012). The impact of
917 mindfulness meditation in promoting a culture of safety on an acute psychiatric unit.
918 *Perspectives in psychiatric care*, 48(3), 129–137. [https://doi-](https://doi-org.ezproxy1.bath.ac.uk/10.1111/j.1744-6163.2011.00315.x)
919 [org.ezproxy1.bath.ac.uk/10.1111/j.1744-6163.2011.00315.x](https://doi-org.ezproxy1.bath.ac.uk/10.1111/j.1744-6163.2011.00315.x)

920 Callender, K. A., Trustey, C. E., Alton, L., & Hao, Y. (2019). Single case evaluation of a
921 mindfulness-based mobile application with a substance abuse counselor. *Counseling*
922 *Outcome Research and Evaluation*. Advance online publication.
923 <https://doi.org/10.1080/21501378.2019.1686353>

924 Canfield, J. (2005). Secondary Traumatization, Burnout, and Vicarious Traumatization: A
925 Review of the Literature as It Relates to Therapists Who Treat Trauma. *Smith College*
926 *Studies in Social Work*, 75, 81–101. https://doi.org/10.1300/J497v75n02_06

927 Care Quality Commission (2022) State of Care 2021/22. Available from:
928 <https://www.cqc.org.uk/publication/state-care-202122> [Accessed: 10.04.2023]

929 *Carmel, A., Fruzzetti, A. E., & Rose, M. L. (2014). Dialectical behavior therapy training to
930 reduce clinical burnout in a public behavioral health system. *Community mental health*
931 *journal*, 50(1), 25–30. <https://doi-org.ezproxy1.bath.ac.uk/10.1007/s10597-013-9679-2>

932 Cavanagh, N., Cockett, G., Heinrich, C., Doig, L., Fiest, K., Guichon, J. R., Page, S.,
933 Mitchell, I., & Doig, C. J. (2020). Compassion fatigue in healthcare providers: A systematic
934 review and meta-analysis. *Nursing Ethics*, 27(3), 639–665.
935 <https://doi.org/10.1177/0969733019889400>

- 936 Cocker, F., & Joss, N. (2016). Compassion Fatigue among Healthcare, Emergency and
937 Community Service Workers: A Systematic Review. *International journal of environmental
938 research and public health*, 13(6), 618. https://doi.org/10.1300/J497v75n02_06
- 939 *Chilton, G., Lynskey, K., Ohnstad, E., Manders, E. (2020). A Case of El Duende: Art-Based
940 Supervision in Addiction Treatment, *Art Therapy*.
941 <https://doi.org/10.1080/07421656.2020.1771138>
- 942 *Chochol, M. D., Pease, E., Swintak, C., & Anand, U. (2021). Addressing Well-being,
943 Burnout, and Professional Fulfillment in Child and Adolescent Psychiatry Trainees During
944 the COVID-19 Pandemic Through Implementation of a Virtual Balint-Like Group: a Pilot
945 Study. *Academic psychiatry : the journal of the American Association of Directors of
946 Psychiatric Residency Training and the Association for Academic Psychiatry*, 45(5), 581–
947 586. <https://doi.org/10.1007/s40596-021-01488-x>
- 948 Conversano C., Ciacchini R., Orrù G., Di Giuseppe M., Gemignani A., Poli A. (2020).
949 Mindfulness, Compassion, and Self-Compassion Among Health Care Professionals: What's
950 New? A Systematic Review. *Frontiers in Psychology*. 11.1683.
951 <https://doi.org/10.1080/07421656.2020.1771138>
- 952 Corcoran, K. & Fischer, J. (1994). *Measures for clinical practice: A Sourcebook* (3rd Ed.,
953 Vol. 2) Adults. New York: The Free Press.
- 954 *Clarke, S., Taylor, G., Lancaster, J., & Remington, B. (2015). Acceptance and commitment
955 therapy-based self-management versus psychoeducation training for staff caring for clients
956 with a personality disorder: a randomized controlled trial. *Journal of personality disorders*,
957 29(2), 163–176. https://doi-org.ezproxy1.bath.ac.uk/10.1521/pedi_2014_28_149
- 958 Cuneo, C. L., Curtis Cooper, M. R., Drew, C. S., Naoum-Heffernan, C., Sherman, T., Walz,
959 K., & Weinberg, J. (2011). The effect of Reiki on work-related stress of the registered nurse.

960 Journal of holistic nursing : official journal of the American Holistic Nurses' Association,
961 29(1), 33–43. <https://doi-org.ezproxy1.bath.ac.uk/10.1177/0898010110377294>

962 Dean, W., Talbot, S., & Dean, A. (2019). Reframing Clinician Distress: Moral Injury Not
963 Burnout. *Federal practitioner: for the health care professionals of the VA, DoD, and PHS*,
964 36(9), 400–402.

965 Deaver, S. P., & Shiflett, C. (2011). Art-based supervision techniques. *The Clinical*
966 *Supervisor*, 30(2), 257–276. <https://doi.org/10.1080/07325223.2011.619456>

967 Demerouti, E. & Bakker, A.B. (2008). The Oldenburg Burnout Inventory: A good alternative
968 to measure burnout and engagement, in: J. Halbesleben (Ed.), *Handbook of stress and*
969 *burnout in health care*, Nova Science Publishers, Hauppauge, NY, 65-78

970 Demerouti, E., Bakker, A. B., Vardakou, I., & Kantas, A. (2003). The convergent validity of
971 two burnout instruments: A multitrait-multimethod analysis. *European Journal of*
972 *Psychological Assessment*, 19(1), 12–23. <https://doi.org/10.1080/07421656.2020.1771138>

973 Demerouti, E., Bakker, A. B., Nachreiner, F., & Schaufeli, W. B. (2000). A model of burnout
974 and life satisfaction amongst nurses. *Journal of advanced nursing*, 32(2), 454–464.
975 <https://doi.org/10.1046/j.1365-2648.2000.01496.x>

976 Demerouti, E. & Nachreiner, F. (1998). Zur Spezifität von Burnout für Dienstleistungsberufe:
977 Fakt oder Artefakt [The specificity of burnout for human services: Fact or artefact] *Zeitschrift*
978 *für Arbeitswissenschaft*, 52, 82-89.

979 Department of Health & Social Care (2012). *The NHS Constitution for England*. Retrieved
980 from: <https://www.gov.uk/government/publications/the-nhs-constitution-for-england#history>
981 [Accessed: 14/12/20]

- 982 Dimeff, L., & Linehan, M.M. (2001). Dialectical behavior therapy in a nutshell. The
983 California Psychologist, 34, 10-13.
- 984 *Doyle, M., Kelly, D., Clarke, S., & Braynion, P. (2007). Burnout: The impact of
985 psychosocial interventions training. *Mental Health Practice*, 10, 16–19.
986 <http://dx.doi.org/10.7748/mhp2007.04.10.7.16.c4298>
- 987 *Duckworth McNeil, A. (2012). Identification and Prevention of Secondary Traumatic Stress
988 in Mental Health Professionals Who Work With Child Sexual Abuse Victim [Unpublished
989 thesis] The University of Southern Mississippi
- 990 Dyrbye, L. N., Satele, D., Sloan, J., & Shanafelt, T. D. (2013). Utility of a brief screening
991 tool to identify physicians in distress. *Journal of general internal medicine*, 28(3), 421–427.
992 <https://doi.org/10.1007/s11606-012-2252-9>
- 993 Dyrbye, L. N., Satele, D., Sloan, J., & Shanafelt, T. D. (2014). Ability of the physician well-
994 being index to identify residents in distress. *Journal of graduate medical education*, 6(1), 78–
995 84. <https://doi.org/10.4300/JGME-D-13-00117.1>
- 996 Edelwich, J. & Brodsky, A. (1980). *Burn-out: Stages of Disillusionment in the Helping*
997 *Professions*. Human Sciences Press: New York.
- 998 Elliott, R., Bohart, A. C., Watson, J. C., & Murphy, D. (2018). Therapist empathy and client
999 outcome: An updated meta-analysis. *Psychotherapy (Chicago, Ill.)*, 55(4), 399–410.
1000 <https://doi.org/10.1037/pst0000175>
- 1001 *Eriksson, T., Germundsjö, L., Åström, E., & Rönnlund, M. (2018). Mindful Self-
1002 Compassion Training Reduces Stress and Burnout Symptoms Among Practicing
1003 Psychologists: A Randomized Controlled Trial of a Brief Web-Based Intervention. *Frontiers*
1004 *in psychology*, 9, 2340. <https://doi-org.ezproxy1.bath.ac.uk/10.3389/fpsyg.2018.02340>

1005 *Ewers, P., Bradshaw, T., McGovern, J., & Ewers, B. (2002). Does training in psychosocial
1006 interventions reduce burnout rates in forensic nurses?. *Journal of advanced nursing*, 37(5),
1007 470–476. <https://doi-org.ezproxy1.bath.ac.uk/10.1046/j.1365-2648.2002.02115.x>

1008 Festinger, L. (1954). A theory of social comparison processes. *Human Relations*, 7, 117-140.

1009 Figley, C. R. (1995). Compassion fatigue: Toward a new understanding of the costs of caring.
1010 In B. H. Stamm (Ed.), *Secondary traumatic stress: Self-care issues for clinicians, researchers,*
1011 *and educators* (pp. 3–28). Baltimore: The Sidran Press.

1012 Figley, C. R. (2002). Compassion fatigue: Psychotherapists’ chronic lack of self-care. *Journal*
1013 *of Clinical Psychology*, 58(11), 1433–1441. <https://doi.org/10.1002/jclp.10090>

1014 Figley, C. R., & Stamm, B. H. (1996). Psychometric review of Compassion Fatigue Self-
1015 Test. In B. H. Stamm (Ed.), *Measurement of stress, trauma, and adaptation* (pp. 127–130).
1016 Brooklandville, MD: Sidran Press.

1017 *Finnamore, C., Rocca, F., Parker, J., Blazdell, J., Dale, O. (2020). The impact of a co-
1018 produced personality disorder training on staff burnout, knowledge and attitudes. *Mental*
1019 *Health Review Journal*, 25(3):269-280. <https://doi.org/10.1108/MHRJ-01-2020-0009>

1020 Fish, B. (2017). *Arts-based Supervision*. Routledge.

1021 *Flarity, K., Nash, K., Jones, W., & Steinbruner, D. (2016). Intervening to Improve
1022 Compassion Fatigue Resiliency in Forensic Nurses. *Advanced emergency nursing journal*,
1023 38(2), 147–156. <https://doi-org.ezproxy1.bath.ac.uk/10.1097/TME.000000000000101>

1024 Fukui, S., Salyers, M. P., Morse, G., & Rollins, A. L. (2021). Factors that affect quality of
1025 care among mental health providers: Focusing on job stress and resources. *Psychiatric*
1026 *rehabilitation journal*, 44(3), 204–211. <https://doi.org/10.1037/prj0000469>

1027 Garner, B. R., Knight, K., & Simpson, D. D. (2007). Burnout among corrections-based drug
1028 treatment staff: Impact of individual and organizational factors. *International Journal of*
1029 *Offender Therapy and Comparative Criminology*, 51(5), 510–522. [https://doi-](https://doi-org.ezproxy1.bath.ac.uk/10.1177/0306624X06298708)
1030 [org.ezproxy1.bath.ac.uk/10.1177/0306624X06298708](https://doi-org.ezproxy1.bath.ac.uk/10.1177/0306624X06298708)

1031 Gardner K. (1991). A summary of findings of a five-year comparison study of primary and
1032 team nursing. *Nursing research*, 40(2), 113–117.

1033 Gentry, J. E. (1996). Solution-focused trauma recovery scale (TRS). Unpublished. Florida
1034 State University: Tallahassee

1035 Gentry, J. E., Baggerly, J., & Baranowsky, A. (2004). Training-as-treatment: effectiveness of
1036 the Certified Compassion Fatigue Specialist Training. *International journal of emergency*
1037 *mental health*, 6(3), 147–155.

1038 Gentry, J. E.; Baranowsky, A. B. (1998). Treatment manual for the accelerated recovery
1039 program: Set II. Toronto, Canada: PsychInk.

1040 Gentry, J. E.; Monson, M. (2017). Professional resilience: Helping doesn't have to hurt.
1041 Phoenix, AZ: Compassion Unlimited.

1042 Gilbert, P. (Ed). (2005). *Compassion: Conceptualizations, research, and use in*
1043 *psychotherapy*. New York, NY: Routledge

1044 Gilbert P. (2014). The origins and nature of compassion focused therapy. *The British journal*
1045 *of clinical psychology*, 53(1), 6–41. <https://doi.org/10.1111/bjc.12043>

1046 Gilmartin, H., Goyal, A., Hamati, M. C., Mann, J., Saint, S., & Chopra, V. (2017). Brief
1047 Mindfulness Practices for Healthcare Providers - A Systematic Literature Review. *The*
1048 *American journal of medicine*, 130(10), 1219.e1–1219.e17.
1049 <https://doi.org/10.1016/j.amjmed.2017.05.041>

1050 Granath, J., Ingvarsson, S., von Thiele, U., & Lundberg, U. (2006). Stress management: a
1051 randomized study of cognitive behavioural therapy and yoga. *Cognitive behaviour therapy*,
1052 35(1), 3–10. <https://doi-org.ezproxy1.bath.ac.uk/10.1080/16506070500401292>

1053 Gregory, A. (2015). Yoga and mindfulness program: The effects on compassion fatigue and
1054 compassion satisfaction in social workers. *Journal of Religion & Spirituality in Social Work:*
1055 *Social Thought*, 34(4), 372–393. <https://doi.org/10.1080/15426432.2015.1080604>

1056 Grossi, G., Perski, A., Evengård, B., Blomkvist, V., & Orth-Gomér, K. (2003). Physiological
1057 correlates of burnout among women. *Journal of psychosomatic research*, 55(4), 309–316.
1058 [https://doi.org/10.1016/s0022-3999\(02\)00633-5](https://doi.org/10.1016/s0022-3999(02)00633-5)

1059 Halbesleben, J. R. B. (2010). The role of exhaustion and workarounds in predicting
1060 occupational injuries: A cross-lagged panel study of health care professionals. *Journal of*
1061 *Occupational Health Psychology*, 15(1), 1–16. <https://doi.org/10.1037/a0017634>

1062 Halbesleben, J. R. B., & Demerouti, E. (2005). The construct validity of an alternative
1063 measure of burnout: Investigating the English translation of the Oldenburg Burnout
1064 Inventory. *Work & Stress*, 19(3), 208–220. [https://doi-](https://doi-org.ezproxy1.bath.ac.uk/10.1080/02678370500340728)
1065 [org.ezproxy1.bath.ac.uk/10.1080/02678370500340728](https://doi-org.ezproxy1.bath.ac.uk/10.1080/02678370500340728)

1066 *Hallberg I. R. (1994). Systematic clinical supervision in a child psychiatric ward:
1067 satisfaction with nursing care, tedium, burnout, and the nurses' own report on the effects of it.
1068 *Archives of psychiatric nursing*, 8(1), 44–52. [https://doi-](https://doi-org.ezproxy1.bath.ac.uk/10.1016/0883-9417(94)90020-5)
1069 [org.ezproxy1.bath.ac.uk/10.1016/0883-9417\(94\)90020-5](https://doi-org.ezproxy1.bath.ac.uk/10.1016/0883-9417(94)90020-5)

1070 Hamaideh S. H. (2011). Burnout, social support, and job satisfaction among Jordanian mental
1071 health nurses. *Issues in mental health nursing*, 32(4), 234–242. [https://doi-](https://doi-org.ezproxy1.bath.ac.uk/10.3109/01612840.2010.546494)
1072 [org.ezproxy1.bath.ac.uk/10.3109/01612840.2010.546494](https://doi-org.ezproxy1.bath.ac.uk/10.3109/01612840.2010.546494)

1073 Harris, C.J. (1995). Sensory-based Therapy for Crisis Counselors. In: C.R. Figley (Ed.),
1074 Compassion fatigue: coping with secondary traumatic stress disorder, 110-114. New York,
1075 NY: Brunner/Mazel.

1076 Hayes, S. C. (2004). Acceptance and Commitment Therapy, Relational Frame Theory, and
1077 the third wave of behavior therapy. *Behavior Therapy*, 35, 639-665.
1078 <https://doi.org/10.1016/j.beth.2016.11.006>

1079 *Hayes, S. C., Bissett, R., Roget, N., Padilla, M., Kohlenberg, B. S., Fisher, G., Masuda, A.,
1080 Pistorello, J., Rye, A. K., Berry, K., & Niccolls, R. (2004). The Impact of Acceptance and
1081 Commitment Training and Multicultural Training on the Stigmatizing Attitudes and
1082 Professional Burnout of Substance Abuse Counselors. *Behavior Therapy*, 35(4), 821–835.
1083 [https://doi-org.ezproxy1.bath.ac.uk/10.1016/S0005-7894\(04\)80022-4](https://doi-org.ezproxy1.bath.ac.uk/10.1016/S0005-7894(04)80022-4)

1084 Hayes, S. C., Luoma, J. B., Bond, F. W., Ma-suda, A., & Lillis, J. (2006). Acceptance and
1085 Commitment Therapy: Model, processes and outcomes. *Behaviour Research and Therapy*,
1086 44(1), 1–25. <https://doi.org/10.1016/j.brat.2005.06.006>

1087 *Haynos, A. F., Fruzzetti, A. E., Anderson, C., Briggs, D., & Walenta, J. (2016). Effects of
1088 dialectical behavior therapy skills training on outcomes for mental health staff in a child and
1089 adolescent residential setting. *Journal of hospital administration*, 5(2), 55–61. [https://doi-](https://doi-org.ezproxy1.bath.ac.uk/10.5430/jha.v5n2p55)
1090 [org.ezproxy1.bath.ac.uk/10.5430/jha.v5n2p55](https://doi-org.ezproxy1.bath.ac.uk/10.5430/jha.v5n2p55)

1091 Henderson, C., Noblett, J., Parke, H., Clement, S., Caffrey, A., Gale-Grant, O. et al. (2014)
1092 Mental health-related stigma in healthcare and mental health–care settings. *Lancet*
1093 *Psychiatry*. 1(6), 467–482. [https://doi.org/10.1016/S2215-0366\(14\)00023-6](https://doi.org/10.1016/S2215-0366(14)00023-6)

1094 Henry, B. J. (2014). Nursing burnout interventions: what is being done?. *Clinical journal of*
1095 *oncology nursing*, 18(2), 211–214. [https://doi-](https://doi-org.ezproxy1.bath.ac.uk/10.1188/14.CJON.211-21)
1096 [org.ezproxy1.bath.ac.uk/10.1188/14.CJON.211-21](https://doi-org.ezproxy1.bath.ac.uk/10.1188/14.CJON.211-21)

1097 Hilliard, R. (2006). The effect of music therapy sessions on compassion fatigue and team
1098 building of professional hospice caregivers. *The Arts in Psychotherapy*, 33, 395–401.
1099 <https://doi.org/10.1016/j.aip.2006.06.002>

1100 Hofmeyer, A., Kennedy, K., & Taylor, R. (2020). Contesting the term ‘compassion
1101 fatigue’: Integrating findings from Social Neuroscience and self-care research. *Collegian*,
1102 27(2), 232–237. <https://doi.org/10.1016/j.colegn.2019.07.001>

1103 Holroyd, K. (1979). Stress, coping, and the treatment of stress related illness. In J. McNamara
1104 (Ed.), *Behavioral approaches to medicine*. New York: Plenum Press.

1105 Horowitz, M., Wilner, N., & Alvarez, W. (1979). Impact of Event Scale: a measure of
1106 subjective stress. *Psychosomatic medicine*, 41(3), 209–218. [https://doi-](https://doi-org.ezproxy1.bath.ac.uk/10.1097/00006842-197905000-00004)
1107 [org.ezproxy1.bath.ac.uk/10.1097/00006842-197905000-00004](https://doi-org.ezproxy1.bath.ac.uk/10.1097/00006842-197905000-00004)

1108 *Hunnicut, A.W., MacMillan, T.F. (1983). Beating Burnout: Findings from a three-year
1109 study. *Journal of Mental Health Administration* 10, 7–9. <https://doi.org/10.1007/BF02833043>

1110 *Ifrach, E.R., Miller, A. (2016). Social action art therapy as an intervention for compassion
1111 fatigue, *The Arts in Psychotherapy*, 50, 34-39 <https://doi.org/10.1016/j.aip.2016.05.009>

1112 Jachens, L., Houdmont, J., Thomas, R. (2018). Work-related stress in a humanitarian context:
1113 A qualitative investigation. *Disasters*, 42, 619–634. <https://doi.org/10.1111/disa.12278>

1114 *Jones M. (2009). The side effects of evidence-based training. *Journal of psychiatric and*
1115 *mental health nursing*, 16(7), 593–598. [https://doi-org.ezproxy1.bath.ac.uk/10.1111/j.1365-](https://doi-org.ezproxy1.bath.ac.uk/10.1111/j.1365-2850.2009.01401.x)
1116 [2850.2009.01401.x](https://doi-org.ezproxy1.bath.ac.uk/10.1111/j.1365-2850.2009.01401.x)

1117 Jones, J., & Lowe, T. (2001). The post-registration education and training of mental health
1118 nurses working in acute inpatient psychiatric settings. *Mental Health Practice*. 5 4), 8-11.
1119 [https://doi.org/10.1016/S0260-6917\(03\)00101-1](https://doi.org/10.1016/S0260-6917(03)00101-1)

1120 *Kiley, K.A., Sehgal, A.R., Neth, S., Dolata, J., Pike, E., Spilsbury, J.C., Albert, J.M.
1121 (2018).The Effectiveness of Guided Imagery in Treating Compassion Fatigue and Anxiety of
1122 Mental Health Workers, *Social Work Research*, 42(1) 33–43,
1123 <https://doi.org/10.1093/swr/svx026>

1124 Killian, K. D. (2008). Helping till it hurts? A multimethod study of compassion fatigue,
1125 burnout, and self-care in clinicians working with trauma survivors. *Traumatology*, 14(2), 32–
1126 44. <https://doi.org/10.1177/1534765608319083>

1127 Kirby, J.N., Tellegen, C.L. and Steindl, S.R. (2017). A meta-analysis of compassion-based
1128 interventions: Current state of knowledge and future directions. *Behavior Therapy*, 48(6),
1129 778-792. <https://doi.org/10.1016/j.beth.2017.06.003>

1130 Klein, A., Taieb, O., Xavier, S., Baubet, T., Reyre, A. (2020). The benefits of mindfulness-
1131 based interventions on burnout among health professionals: A systematic review, *EXPLORE*,
1132 16(1), 35-43. <https://doi.org/10.1016/j.explore.2019.09.002>.

1133 Klimecki, O.M., Leiberg, S., Lamm, C. and Singer, T. (2013). Functional neural plasticity
1134 and associated changes in positive affect after compassion training. *Cerebral cortex*, 23(7),
1135 1552-1561. <https://doi.org/10.1093/cercor/bhs142>

1136 Klimecki, O.M., Leiberg, S., Ricard, M. and Singer, T. (2014). Differential pattern of
1137 functional brain plasticity after compassion and empathy training. *Social cognitive and*
1138 *affective neuroscience*, 9(6), 873-879. <https://doi.org/10.1093/scan/nst060>

1139 Knight, C. (2013). Indirect trauma: Implications for self-care, supervision, the organization,
1140 and the academic institution. *The Clinical Supervisor*, 32(2), 224-243.
1141 <https://doi.org/10.1080/07325223.2013.850139>

1142 Knudsen, H. K., Ducharme, L. J., & Roman, P. M. (2006). Counselor emotional exhaustion
1143 and turnover intention in therapeutic communities. *Journal of substance abuse treatment*,
1144 31(2), 173–180. <https://doi.org/10.1016/j.jsat.2006.04.003>

1145 Knudsen, H. K., Johnson, J. A., & Roman, P. M. (2003). Retaining counseling staff at
1146 substance abuse treatment centers: Effects of management practices. *Journal of Substance*
1147 *Abuse Treatment*, 24(2), 129–135. [https://doi.org/10.1016/S0740-5472\(02\)00357-4](https://doi.org/10.1016/S0740-5472(02)00357-4)

1148 *Kovač, J., Krečič, M.J., Čagran, B., Mulej, M (2017). Effect of Supervision on Stress and
1149 Burnout in School Counsellors: A Case of Action Research. *Systemic Practice and Action*
1150 *Research* 30, 395–406. <https://doi.org/10.1007/s11213-016-9400-9>

1151 Kristensen, T. S., Borritz, M., Villadsen, E., & Christensen, K. B. (2005). The Copenhagen
1152 Burnout Inventory: A new tool for the assessment of burnout. *Work & Stress*, 19(3), 192–
1153 207. <https://doi.org/10.1080/02678370500297720>

1154 Lamothe, M., Rondeau, É., Malboeuf-Hurtubise, C., Duval, M., Sultan, S. (2016). Outcomes
1155 of MBSR or MBSR-based interventions in health care providers: A systematic review with a
1156 focus on empathy and emotional competencies, *Complementary Therapies in Medicine*, 24,
1157 19-28. <https://doi.org/10.1016/j.ctim.2015.11.001>.

1158 *Landis, E.M. (2010). Sharevision: A Collarative-Reflective, Expressive Arts Intervention To
1159 Address Trauma [Unpublished thesis] Graduate School of Arts and Social Sciences, Lesley
1160 University

1161 Lasalvia, A., Bonetto, C., Bertani, M., Bissoli, S., Cristofalo, D., Marrella, G., . . . Tansella,
1162 M. (2009). Influence of perceived organisational factors on job burnout: Survey of
1163 community mental health staff. *British Journal of Psychiatry*, 195(6), 537-544.
1164 <https://doi.org/10.1192/bjp.bp.108.060871>

- 1165 Laurenceau, J., Barrett, L. F., & Pietromonaco, P. R. (1998). Intimacy as an interpersonal
1166 process: The importance of self-disclosure, partner disclosure, and perceived partner
1167 responsiveness in interpersonal exchanges. *Journal of Personality & Social Psychology*,
1168 74(5),1238–1251. <https://doi.org/10.1037/0022-3514.74.5.1238>.
- 1169 Laurenceau, J., Barrett, L. F., & Rovine, M. J. (2005). The interpersonal process model of
1170 intimacy in marriage: A daily-diary and multi-level modelling approach. *Journal of Family*
1171 *Psychology*, 19(2), 314–323. <https://doi.org/10.1037/0893-3200.19.2.314>.
- 1172 Leiter, M. P., & Schaufeli, W. B. (1996). Consistency of the burnout construct across
1173 occupations. *Anxiety, Stress & Coping: An International Journal*, 9(3), 229–243.
1174 <https://doi.org/10.1080/10615809608249404>
- 1175 *Leykin, Y., Cucciare, M. A., & Weingardt, K. R. (2011). Differential effects of online
1176 training on job-related burnout among substance abuse counsellors. *Journal of Substance*
1177 *Use*, 16(2), 127–135. <https://doi-org.ezproxy1.bath.ac.uk/10.3109/14659891.2010.526168>
- 1178 Linehan, M. M. (1993). *Diagnosis and treatment of mental disorders. Cognitive-behavioral*
1179 *treatment of borderline personality disorder*. Guilford Press.
- 1180 *Little, L. B. (2000). *Training in dialectical behavior therapy as a means of reducing therapist*
1181 *burnout*. *Dissertation Abstracts International: Section B: The Sciences and Engineering*, 61(5-
1182 B), 2769.
- 1183 Luken, M., Sammons, A. (2016). Systematic review of mindfulness practice for reducing job
1184 burnout. *American Journal of Occupational Therapy*, 70.
1185 <https://doi.org/10.5014/ajot.2016.016956>
- 1186 Lundgren-Nilsson, Å., Jonsdottir, I.H., Pallant, J. & Ahlborg Junior, G. (2012). Internal
1187 construct validity of the Shirom-Melamed Burnout Questionnaire (SMBQ). *BMC Public*
1188 *Health*, 12(1) <https://doi.org/10.1186/1471-2458-12-1>

1189 *Luoma, J. B., & Vilardaga, J. P. (2013). Improving therapist psychological flexibility while
1190 training acceptance and commitment therapy: a pilot study. *Cognitive behaviour therapy*,
1191 42(1), 1–8. <https://doi-org.ezproxy1.bath.ac.uk/10.1080/16506073.2012.701662>

1192 MacGuire, J. M., & Botting, D. A. (1990). The use of the Ethnograph program to identify the
1193 perceptions of nursing staff following the introduction of primary nursing in an acute medical
1194 ward for elderly people. *Journal of advanced nursing*, 15(10), 1120–1127. [https://doi-](https://doi-org.ezproxy1.bath.ac.uk/10.1111/j.1365-2648.1990.tb01702.x)
1195 [org.ezproxy1.bath.ac.uk/10.1111/j.1365-2648.1990.tb01702.x](https://doi-org.ezproxy1.bath.ac.uk/10.1111/j.1365-2648.1990.tb01702.x)

1196 Maslach, C. (1982). *Burnout: The cost of caring*. Englewood Cliffs, NJ: Prentice Hall

1197 Maslach, C. (1993). Burnout: A multidimensional perspective. In W. B. Schaufeli, C.
1198 Maslach, & T. Marek (Eds.). *Series in applied psychology: Social issues and questions*.
1199 *Professional burnout: Recent developments in theory and research* (p. 19–32). Taylor &
1200 Francis.

1201 Maslach, C. (2006). Understanding Job Burnout, In: Rossi, A.M., Pamela L. Perrewé, P. L.,
1202 & Sauter, S. L. *Stress and Quality of Working Life: Current Perspectives in Occupational*
1203 *Health*. Information Age Publishing

1204 Maslach, C. & Jackson, S.E. (1981). *Maslach Burnout Inventory*. 2. Palo Alto: Consulting
1205 Psychologists Press

1206 Maslach, C., Schaufeli, W.B. & Leiter, M.P. (2001). Job Burnout. *Annual Review of*
1207 *Psychology*. 52, 397-422. <https://doi.org/10.1146/annurev.psych.52.1.397>

1208 Maslow, A. (1968). *Toward a psychology of being*. New York: Van Nostrand,

1209 Mathieu, F. (2007). Running on empty: Compassion fatigue in health professionals. *Rehab*
1210 *Community Care Medicine*. 4, 1–7.

1211 Mauritz, M., Goossens, P., Draijer, N., & van Achterberg, T. (2013). Prevalence of
1212 interpersonal trauma exposure and trauma-related disorders in severe mental illness.
1213 European Journal Of Psychotraumatology, 4(1), 19985.
1214 <https://doi.org/10.3402/ejpt.v4i0.19985>

1215 *Mehr, L. M., Senteney, A., & Creadie, T. M. (1995). Daydreams, stress and burn-out in
1216 women mental health workers: A preliminary clinical report. Imagination, Cognition and
1217 Personality, 14(2), 105–115. [https://doi-org.ezproxy1.bath.ac.uk/10.2190/103X-40HD-](https://doi-org.ezproxy1.bath.ac.uk/10.2190/103X-40HD-A1QN-7KC5)
1218 [A1QN-7KC5](https://doi-org.ezproxy1.bath.ac.uk/10.2190/103X-40HD-A1QN-7KC5)

1219 Melamed, S., Ugarten, U., Shirom, A., Kahana, L., Lerman, Y., & Froom, P. (1999). Chronic
1220 burnout, somatic arousal and elevated salivary cortisol levels. Journal of psychosomatic
1221 research, 46(6), 591–598. [https://doi-org.ezproxy1.bath.ac.uk/10.1016/s0022-3999\(99\)00007-](https://doi-org.ezproxy1.bath.ac.uk/10.1016/s0022-3999(99)00007-0)
1222 [0](https://doi-org.ezproxy1.bath.ac.uk/10.1016/s0022-3999(99)00007-0)

1223 *Melchior, M. E., Philipsen, H., Abu-Saad, H. H., Halfens, R. J., van de Berg, A. A., &
1224 Gassman, P. (1996). The effectiveness of primary nursing on burnout among psychiatric
1225 nurses in long-stay settings. Journal of advanced nursing, 24(4), 694–702.
1226 <https://doi.org/10.1046/j.1365-2648.1996.02457.x>

1227 Mensah, S. B., & Anderson, J. G. (2015). Barriers and facilitators of the use of mind-body
1228 therapies by healthcare providers and clinicians to care for themselves. Complementary
1229 therapies in clinical practice, 21(2), 124–130. [https://doi-](https://doi-org.ezproxy1.bath.ac.uk/10.1016/j.ctcp.2015.01.004)
1230 [org.ezproxy1.bath.ac.uk/10.1016/j.ctcp.2015.01.004](https://doi-org.ezproxy1.bath.ac.uk/10.1016/j.ctcp.2015.01.004)

1231 Metcalf, J. & Jacobs, W. J. (1996). A “hot-system/cool-system” view of memory under
1232 stress. The National Center for Post-Traumatic Stress Disorder Research Quarterly, 7(2), 1-3.
1233 <https://doi.org/10.1016/B978-0-12-800951-2.00020-0>

1234 Mitchell K. (1994). Reiki: A Torch in Daylight. St. Charles, IL: Mind River Publications.

1235 Montgomery, A. (2014). The inevitability of physician burnout: Implications for
1236 interventions. *Burnout Research*, 1, 50–6. <https://doi.org/10.1016/j.burn.2014.04.002>

1237 Moore E., Ball R.A. & Kuipers L. (1992). Expressed emotion in staff working with the long
1238 term adult mentally ill. *British Journal of Psychiatry*, 161, 802-808.
1239 <https://doi.org/10.1192/bjp.161.6.802>

1240 Morewedge, C. K., Huh, Y. E., & Vosgerau, J. (2010). Thought for food: imagined
1241 consumption reduces actual consumption. *Science (New York, N.Y.)*, 330(6010), 1530–1533.
1242 <https://doi-org.ezproxy1.bath.ac.uk/10.1126/science.1195701>

1243 Moyer, M. (2011). Effects of non-guidance activities, supervision, and student-to counsellor
1244 ratios on school counsellor burnout. *Journal of School Counselling*, 9(5):1–31

1245 Munn, Z., Peters, M. D. J., Stern, C., Tufanaru, C., McArthur, A., & Aromataris, E. (2018).
1246 Systematic review or scoping review? Guidance for authors when choosing between a
1247 systematic or scoping review approach. *BMC medical research methodology*, 18(1).
1248 <https://doi.org/10.1186/s12874-018-0611-x>

1249 Newell, J.M. & MacNeil, Gordon. (2010). Professional burnout, vicarious trauma, secondary
1250 traumatic stress, and compassion fatigue: A review of theoretical terms, risk factors, and
1251 preventive methods for clinicians and researchers. *Best Practice in Mental Health*. 6. 57-68.

1252 *Newman, G.F., Maggott, C., Alexander, D.G. (2015). Group drumming as a burnout
1253 prevention initiative among staff members at a child and adolescent mental health care
1254 facility. *South African Journal of Psychology*, 45(4), 439-451.
1255 <https://doi.org/10.1177/0081246315581346>

1256 NHS (2020). NHS Staff Survey 2019: National results briefing. Available from:
1257 [https://www.nhsstaffsurveys.com/Caches/Files/ST19_National%20briefing_FINAL%20V2.p](https://www.nhsstaffsurveys.com/Caches/Files/ST19_National%20briefing_FINAL%20V2.pdf)
1258 [df](https://www.nhsstaffsurveys.com/Caches/Files/ST19_National%20briefing_FINAL%20V2.pdf) [Accessed: 10/01/21]

- 1259 NHS (2023a) NHS Staff Survey 2022: National results briefing. Available from:
- 1260 [https://www.nhsstaffsurveys.com/static/8c6442c8d92624a830e6656baf633c3f/NHS-](https://www.nhsstaffsurveys.com/static/8c6442c8d92624a830e6656baf633c3f/NHS-Staff-Survey-2022-National-briefing.pdf)
- 1261 [Staff-Survey-2022-National-briefing.pdf](https://www.nhsstaffsurveys.com/static/8c6442c8d92624a830e6656baf633c3f/NHS-Staff-Survey-2022-National-briefing.pdf) [Accessed 10.04.23]
- 1262 NHS Employers (2022, April 9) Beating burnout in the NHS.
- 1263 <https://www.nhsemployers.org/articles/beating-burnout-nhs>
- 1264 NHS England (2018). Towards commissioning for workplace compassion: a support guide.
- 1265 Retrieved from: [https://www.england.nhs.uk/wp-content/uploads/2018/10/towards-](https://www.england.nhs.uk/wp-content/uploads/2018/10/towards-commissioning-for-workplace-compassion-a-support-guide-v2.pdf)
- 1266 [commissioning-for-workplace-compassion-a-support-guide-v2.pdf](https://www.england.nhs.uk/wp-content/uploads/2018/10/towards-commissioning-for-workplace-compassion-a-support-guide-v2.pdf) [Accessed: 10/01/21]
- 1267 NHS England (2023b) Workforce Race Equality Standard Report. Retrieved from:
- 1268 [https://www.england.nhs.uk/publication/nhs-workforce-race-equality-standard-](https://www.england.nhs.uk/publication/nhs-workforce-race-equality-standard-2022/)
- 1269 [2022/](https://www.england.nhs.uk/publication/nhs-workforce-race-equality-standard-2022/)[Accessed: 10/04/2023]
- 1270 O'Connor, K., Muller Neff, D., & Pitman, S. (2018). Burnout in mental health professionals:
- 1271 A systematic review and meta-analysis of prevalence and determinants. *European Psychiatry*,
- 1272 53, 74-99. <https://doi.org/10.1016/j.eurpsy.2018.06.003>
- 1273 Ortlepp, K., & Friedman, M. (2001). The relationship between sense of coherence and
- 1274 indicators of secondary traumatic stress in non-professional trauma counsellors. *South*
- 1275 *African Journal of Psychology*, 31, 38-45. <https://doi.org/10.1177/008124630103100205>
- 1276 Pascual-Leone, A., Nguyet, D., Cohen, L. G., Brasil-Neto, J. P., Cammarota, A., & Hallett,
- 1277 M. (1995). Modulation of muscle responses evoked by transcranial magnetic stimulation
- 1278 during the acquisition of new fine motor skills. *Journal of neurophysiology*, 74(3), 1037–
- 1279 1045. <https://doi-org.ezproxy1.bath.ac.uk/10.1152/jn.1995.74.3.1037>

1280 *Paulson, L. R., Casile, W. J., & Jones, D. (2015). Tech it out: Implementing an online peer
1281 consultation network for rural mental health professionals. *Journal of Rural Mental Health*,
1282 39(3-4), 125–136. <https://doi-org.ezproxy1.bath.ac.uk/10.1037/rmh0000034>

1283 Pearlman, L. A., & Saakvitne, K. (1995). *Trauma and the therapist: Countertransference and*
1284 *vicarious traumatization in psychotherapy with incest survivors*. New York: W. W. Norton.

1285 Penko, T. (1994). Izgorelost pri delu. In: Lamovec T (ed) *Psihodiagnostika I. Znanstveni*
1286 *institut Filozofske fakultete, Ljubljana*, 324–343

1287 *Perseius, K. I., Kåver, A., Ekdahl, S., Asberg, M., & Samuelsson, M. (2007). Stress and
1288 burnout in psychiatric professionals when starting to use dialectical behavioural therapy in
1289 the work with young self-harming women showing borderline personality symptoms. *Journal*
1290 *of psychiatric and mental health nursing*, 14(7), 635–643. [https://doi-](https://doi-org.ezproxy1.bath.ac.uk/10.1111/j.1365-2850.2007.01146.x)
1291 [org.ezproxy1.bath.ac.uk/10.1111/j.1365-2850.2007.01146.x](https://doi-org.ezproxy1.bath.ac.uk/10.1111/j.1365-2850.2007.01146.x)

1292 Peters, M., Godfrey, C., Khalil, H., McInerney, P., Soares, C. & Parker, D. (2017). 2017
1293 Guidance for the Conduct of JBI Scoping Reviews. In: Aromataris, E. & Munn, J. (eds.)
1294 (2017) *Joanna Briggs Institute Reviewer's Manual*. The Joanna Briggs Institute.

1295 Pines, A., & Aronson, E. (1988). *Career Burnout. Causes & Cures*. New York, NY: The Free
1296 Press.

1297 Pines, A. and Kafry, D. (1978). Occupational tedium in the social services. *Social work*,
1298 23(6), 499-507.

1299 Poghosyan, L., Aiken, L. H., & Sloane, D. M. (2009). Factor structure of the Maslach
1300 burnout inventory: an analysis of data from large scale cross-sectional surveys of nurses from
1301 eight countries. *International journal of nursing studies*, 46(7), 894–902. [https://doi-](https://doi-org.ezproxy1.bath.ac.uk/10.1016/j.ijnurstu.2009.03.004)
1302 [org.ezproxy1.bath.ac.uk/10.1016/j.ijnurstu.2009.03.004](https://doi-org.ezproxy1.bath.ac.uk/10.1016/j.ijnurstu.2009.03.004)

- 1303 Popa-Velea, O., Trutescu, C. I., & Diaconescu, L. V. (2019). The impact of Balint work on
1304 alexithymia, perceived stress, perceived social support and burnout among physicians
1305 working in palliative care: a longitudinal study. *International journal of occupational*
1306 *medicine and environmental health*, 32(1), 53–63.
1307 <https://doi.org/10.13075/ijomeh.1896.01302>
- 1308 Raab, K. (2014). Mindfulness, self-compassion, and empathy among health care
1309 professionals: A review of the literature. *Journal of Health Care Chaplaincy*, 20, 95–108.
1310 <https://doi.org/10.1080/08854726.2014.913876>
- 1311 *Raab, K., Sogge, K., Parker, N., & Flament, M.F. (2015). Mindfulness-based stress
1312 reduction and self-compassion among mental healthcare professionals: a pilot study. *Mental*
1313 *Health, Religion & Culture*, 18(6), 503-512. <https://doi.org/10.1080/13674676.2015.1081588>
- 1314 Rauvola, R.S., Vega, D.M. & Lavigne, K.N. (2019). Compassion Fatigue, Secondary
1315 Traumatic Stress, and Vicarious Traumatization: a Qualitative Review and Research Agenda.
1316 *Occupational Health Science*, 3, 297–336 <https://doi.org/10.1007/s41542-019-00045-1>
- 1317 *Ray, C.A. (1981). Holistic stress management training: a burnout strategy for mental health
1318 workers [Unpublished dissertation] North Texas State University
- 1319 *Razzaque, R., Wood, L. (2016). Exploration of the Effectiveness and Acceptability of a
1320 Professional Mindfulness Retreat for Psychiatrists. *Mindfulness* 7, 340–348.
1321 <https://doi.org/10.1007/s12671-015-0443-2>
- 1322 Read, J., Hammersley, P., & Rudegeair, T. (2007). Why, when and how to ask about
1323 childhood abuse. *Advances in Psychiatric Treatment*, 13(2), 101-110.
1324 <https://doi.org/10.1192/apt.bp.106.002840>
- 1325 *Redhead, K., Bradshaw, T., Braynion, P., & Doyle, M. (2011). An evaluation of the
1326 outcomes of psychosocial intervention training for qualified and unqualified nursing staff

- 1327 working in a low-secure mental health unit. *Journal of psychiatric and mental health nursing*,
1328 18(1), 59–66. <https://doi-org.ezproxy1.bath.ac.uk/10.1111/j.1365-2850.2010.01629.x>
- 1329 Reed S. E. (1988). A comparison of nurse-related behaviour, philosophy of care and job
1330 satisfaction in team and primary nursing. *Journal of advanced nursing*, 13(3), 383–395.
1331 <https://doi-org.ezproxy1.bath.ac.uk/10.1111/j.1365-2648.1988.tb01433.x>
- 1332 Reis, H. T., & Shaver, P. (1988). Intimacy as an interpersonal process. In Duck, S. (Ed.),
1333 *Handbook of personal relationships: Theory, research & interventions* (pp. 367–389).
1334 Chichester: Wiley.
- 1335 *Reyes Ortega, M.A., Kuczynski, A.M., Kanter, J.W., de Montis, I. A., Santos, M.M. (2019).
1336 A Preliminary Test of a Social Connectedness Burnout Intervention for Mexican Mental
1337 Health Professionals. *The Psychological Record*, 69, 267–276.
1338 <https://doi.org/10.1007/s40732-019-00338-5>
- 1339 *Riley, K. E., Park, C. L., Wilson, A., Sabo, A., Antoni, M. H., Braun, T. D., Harrington, J.,
1340 Reiss, J., Pasalis, E., Harris, A., & Cope, S. (2017). Improving physical and mental health in
1341 frontline mental health care providers: Yoga-based stress management versus cognitive
1342 behavioral stress management. *Journal of workplace behavioral health*, 32(1), 26–48.
1343 <https://doi-org.ezproxy1.bath.ac.uk/10.1080/15555240.2016.1261254>
- 1344 *Roberts, C. (2018). *The Effectiveness of Dialectical Behaviour Therapy Skills Training on*
1345 *Outcomes for Mental Health & Learning Disability Staff* [Unpublished doctoral dissertation]
1346 Bangor University
- 1347 Rogers, C.R. (1967). *On Becoming a Person: A Therapist's View of Psychotherapy*;
1348 Constable: London, UK,
- 1349 *Rollins, A. L., Kukla, M., Morse, G., Davis, L., Leiter, M., Monroe-DeVita, M., Flanagan,
1350 M. E., Russ, A., Wasmuth, S., Eliacin, J., Collins, L., & Salyers, M. P. (2016). Comparative

1351 Effectiveness of a Burnout Reduction Intervention for Behavioral Health Providers.
1352 Psychiatric services (Washington, D.C.), 67(8), 920–923. [https://doi-](https://doi-org.ezproxy1.bath.ac.uk/10.1176/appi.ps.201500220)
1353 [org.ezproxy1.bath.ac.uk/10.1176/appi.ps.201500220](https://doi-org.ezproxy1.bath.ac.uk/10.1176/appi.ps.201500220)

1354 *Rosada, R. M., Rubik, B., Mainguy, B., Plummer, J., & Mehl-Madrona, L. (2015). Reiki
1355 Reduces Burnout Among Community Mental Health Clinicians. *Journal of alternative and*
1356 *complementary medicine (New York, N.Y.)*, 21(8), 489–495. [https://doi-](https://doi-org.ezproxy1.bath.ac.uk/10.1089/acm.2014.0403)
1357 [org.ezproxy1.bath.ac.uk/10.1089/acm.2014.0403](https://doi-org.ezproxy1.bath.ac.uk/10.1089/acm.2014.0403)

1358 Ross, S., Randhawa, M., Jabbal, J., Dahir, S., West, R., Murphy, S., Thorley, J., Joubert, H.
1359 (2020) 'A long way to go': ethnic minority NHS staff share their stories. The King's Fund.
1360 Retrieved from: [https://features.kingsfund.org.uk/2020/07/ethnic-minority-nhs-staff-racism-](https://features.kingsfund.org.uk/2020/07/ethnic-minority-nhs-staff-racism-discrimination/index.html)
1361 [discrimination/index.html](https://features.kingsfund.org.uk/2020/07/ethnic-minority-nhs-staff-racism-discrimination/index.html)

1362 Rothschild, B., Rand, .L. (2006). *Help for the helper: Self-care strategies for managing*
1363 *burnout and stress*. New York, NY: W.W. Norton & Company. [Accessed 15/07/2021]

1364 *Scarnera, P., Bosco, A., Soleti, E., & Lancioni, G. E. (2009). Preventing burnout in mental
1365 health workers at interpersonal level: an Italian pilot study. *Community mental health journal*,
1366 45(3), 222–227. <https://doi-org.ezproxy1.bath.ac.uk/10.1007/s10597-008-9178-z>

1367 Schacter, S. (1967). The interaction of cognitive and physiological determinants of emotional
1368 states. In Berkowitz (Ed), *Advances in Experimental Social Psychology*, Vol. 1, New York :
1369 Academic Press.

1370 Schaufeli, W. B. (1990). *Opgebrd: Over de achtergronden van werkstress bij contactuele*
1371 *beroepen - her burnout-syndroom (Burnout: About the job stress in the human services*
1372 *professions - The burnout syndrome)*, Ad. Donker, Rotterdam.

1373 Schaufeli, W.B., Leiter, M.P. and Maslach, C. (2009). Burnout: 35 years of research and
1374 practice. *Career Development International*, 14(3), 204-220.
1375 <https://doi.org/10.1108/13620430910966406>

1376 Schaufeli, W.B., Leiter, M.P., Maslach, C., & Jackson, S.E. (1996). The MBI-General
1377 Survey. In C. Maslach, S.E. Jackson, & M.P. Leiter (Eds.), *Maslach Burnout Inventory*.
1378 *Manual* (3rd ed., pp. 19-26). Palo Alto, CA: Consulting Psychologists Press

1379 Schaufeli, W. B., & Taris, T. W. (2014). A critical review of the job demands-resources
1380 model: Implications for improving work and health. In G. F. Bauer & O. Hämmig (Eds.),
1381 *Bridging occupational, organizational and public health: A transdisciplinary approach* (p. 43–
1382 68). Springer Science + Business Media. https://doi.org/10.1007/978-94-007-5640-3_4

1383 Schaufeli, W. B., & Van Dierendonck, D. (1993). The construct validity of two burnout
1384 measures. *Journal of Organizational Behavior*, 14(7), 631–647.
1385 <https://doi.org/10.1002/job.4030140703>

1386 Schwartz, R., Shanafelt, T. D., Gimmler, C., & Osterberg, L. (2020). Developing institutional
1387 infrastructure for physician wellness: qualitative Insights from VA physicians. *BMC health*
1388 *services research*, 20(1), 7. <https://doi.org/10.1186/s12913-019-4783-9>

1389 Shay, J. (2014). Moral injury. *Psychoanalytic Psychology*, 31(2), 182–191.
1390 <https://doi.org/10.1037/a0036090>

1391 Shubin, S. (1978). Burnout, the professional hazard you face in nursing. *Nursing*, 8(7), 22-27.

1392 Singh, J., Karanika-Murray, M., Baguley, T., & Hudson, J. (2020). A Systematic Review of
1393 Job Demands and Resources Associated with Compassion Fatigue in Mental Health
1394 Professionals. *International Journal of Environmental Research and Public Health*, 17(19),
1395 6987. <https://doi.org/10.3390/ijerph17196987>

- 1396 Sirigatti S., Stefanile C. (1993). “Adattamento e taratura per l’Italia” in MBI Maslach burnout
1397 inventory. eds. Maslach C., Jackson S. (Firenze: Organizzazioni Speciali;), 33–42.
- 1398 Spandler, H., & Stickley, T. (2011). No Hope without Compassion: The Importance of
1399 Compassion in Recovery-focused Mental Health Services. *Journal of Mental Health* 20(6):
1400 555–566. <https://doi.org/10.3109/09638237.2011.583949>
- 1401 Stamm, B.H. (2002). Measuring compassion satisfaction as well as fatigue: developmental
1402 history of the compassion fatigue and satisfaction test. In: Figley CR, editor. *Treating*
1403 *compassion fatigue*. New York: Brunner-Routledge. p. 107–19.
- 1404 Stamm, B. H. (2009). Professional Quality of Life: Compassion Satisfaction and Fatigue
1405 Version 5 (ProQOL). Retrieved from: [/www.isu.edu/~bhstamm](http://www.isu.edu/~bhstamm) [Accessed 22.04.21]
- 1406 Strauss, C., Taylor, B.L., Gu, J., Kuyken, W., Baer, R., Jones, F., Cavanagh, K. (2016). What
1407 is compassion and how can we measure it? A review of definitions and measures. *Clinical*
1408 *Psychology Review*, 47, 15-27. <https://doi.org/10.1016/j.cpr.2016.05.004>
- 1409 Sutton, L., Rowe, S., Hammerton, G., & Billings, J. (2022). The contribution of
1410 organisational factors to vicarious trauma in mental health professionals: a systematic review
1411 and narrative synthesis. *European journal of psychotraumatology*, 13(1), 2022278.
1412 <https://doi.org/10.1080/20008198.2021.2022278>
- 1413 *Suyi, Y., Meredith, P., & Khan, A. (2017). Effectiveness of Mindfulness Intervention in
1414 Reducing Stress and Burnout for Mental Health Professionals in Singapore. *Explore* (New
1415 York, N.Y.), 13(5), 319–326. [https://doi-](https://doi-org.ezproxy1.bath.ac.uk/10.1016/j.explore.2017.06.001)
1416 [org.ezproxy1.bath.ac.uk/10.1016/j.explore.2017.06.001](https://doi-org.ezproxy1.bath.ac.uk/10.1016/j.explore.2017.06.001)
- 1417 Tetrack L. E., Quick J. C. (2011). Overview of occupational health psychology: public health
1418 in occupational settings, in *Handbook of Occupational Health Psychology*, 2nd Edn, eds
1419 Quick J. C., Tetrack L. E. (Washington, DC: American Psychological Association;), 3–20.

1420 Thompson, E.H., Frick, M.H., Trice-Black, S. (2011). Counselor-in-training perceptions of
1421 supervision practices related to self-care and burnout. *The Professional Counsellor*, 1(3),
1422 152–162, <https://doi.org/10.15241/eht.1.3.152>

1423 Thompson, J. (1980). Burnout in group home houseparents. *American Journal of Psychiatry*,
1424 137(6), 710-714. <https://doi.org/10.1176/ajp.137.6.710>

1425 Timms, C., Brough, P., & Graham, D. (2012). Burnt-out but engaged: the co-existence of
1426 psychological burnout and engagement. *Journal of Educational Administration*, 50(3), 327-
1427 345. <https://doi.org/10.1108/09578231211223338>

1428 Ting, L., Jacobson, J.M., Sanders, S. Bride, B.E. & Harrington, D. (2005). The Secondary
1429 Traumatic Stress Scale (STSS), *Journal of Human Behavior in the Social Environment*, 11
1430 (3-4), 177-194. https://doi.org/0.1300/J137v11n03_09

1431 Tricco, A, C., Lillie, E., Zarin, W., O'Brien, K.K., Colquhoun, H., Levac, D., Moher, D.,
1432 Peters, M,D., Horsley, T., Weeks, L., Hempel, S., et al. (2018). PRISMA extension for
1433 scoping reviews (PRISMA-ScR): checklist and explanation. *Annals of Internal Medicine*,
1434 169(7), 467-473. <https://doi.org/10.7326/M18-0850>

1435 Trockel, M., Bohman, B., Lesure, E., Hamidi, M. S., Welle, D., Roberts, L., & Shanafelt, T.
1436 (2018). A Brief Instrument to Assess Both Burnout and Professional Fulfillment in
1437 Physicians: Reliability and Validity, Including Correlation with Self-Reported Medical
1438 Errors, in a Sample of Resident and Practicing Physicians. *Academic psychiatry : the journal*
1439 *of the American Association of Directors of Psychiatric Residency Training and the*
1440 *Association for Academic Psychiatry*, 42(1), 11–24. [https://doi.org/10.1007/s40596-017-](https://doi.org/10.1007/s40596-017-0849-3)
1441 [0849-3](https://doi.org/10.1007/s40596-017-0849-3)

- 1442 Turgoose, D., & Maddox, L. (2017). Predictors of compassion fatigue in mental health
1443 professionals: A narrative review. *Traumatology*, 23(2), 172–185.
1444 <https://doi.org/10.1037/trm0000116>
- 1445 *Van Kirk M. L. (2021). Employee Wellness Pilot Program. *Workplace health & safety*,
1446 69(5), 192–197. <https://doi-org.ezproxy1.bath.ac.uk/10.1177/2165079920976522>
- 1447 Van Roy, K., Vanheule, S., & Inslegers, R. (2015). Research on Balint groups: A literature
1448 review. *Patient education and counseling*, 98(6), 685–694.
1449 <https://doi.org/10.1016/j.pec.2015.01.014>
- 1450 *Walker, D. (2018). Examining the Effects of a Brief Mindfulness-Based Stress Reduction
1451 Program on Burnout in Mental Health Professionals [Unpublished thesis] Grand Canyon
1452 University
- 1453 *Weingardt, K. R., Cucciare, M. A., Bellotti, C., & Lai, W. P. (2009). A randomized trial
1454 comparing two models of web-based training in cognitive-behavioral therapy for substance
1455 abuse counselors. *Journal of substance abuse treatment*, 37(3), 219–227. [https://doi-](https://doi-org.ezproxy1.bath.ac.uk/10.1016/j.jsat.2009.01.002)
1456 [org.ezproxy1.bath.ac.uk/10.1016/j.jsat.2009.01.002](https://doi-org.ezproxy1.bath.ac.uk/10.1016/j.jsat.2009.01.002)
- 1457 Wessells, D.T., Jr., Kutscher, A.H., Seeland, LB., Selder, F.E., Cherico, D.J., & Clark, E.J.
1458 (1989). *Professional Burnout in Medicine and the Helping Professions*. New York, NY: The
1459 Haworth Press.
- 1460 West, M. A. & Chowla, R. (2017). Compassionate leadership for compassionate health care.
1461 In Gilbert, P. (ed.). *Compassion: Concepts, Research and Applications*. (pp. 237-257).
1462 London: Routledge.
- 1463 Whitfield, N., & Kanter, D. (2014). Helpers in distress: Preventing secondary trauma.
1464 *Reclaiming Children and Youth*, 22(4), 59-61.

1465 Wilkie, T., Tajirian, T., & Stergiopoulos, V. (2022). Advancing physician wellness,
 1466 engagement and excellence in a mental health setting: a Canadian perspective. *Health*
 1467 *promotion international*, 37(1), daab061. <https://doi.org/10.1093/heapro/daab061>

1468 *Wymer, B. (2019). An Investigation of the Impact of a Supervision Intervention on
 1469 Secondary Traumatic Stress Responses Among Counselors Treating Child Survivors of
 1470 Sexual Abuse [Unpublished thesis] University of South Carolina.

1471

1472

1473

1474

Appendices

1475 **Appendix A: Preferred Reporting Items for Systematic reviews and Meta-Analyses** 1476 **extension for Scoping Reviews (PRISMA-ScR) Checklist**

SECTION	ITEM	PRISMA-ScR CHECKLIST ITEM	REPORTED ON PAGE #
TITLE			
Title	1	Identify the report as a scoping review.	1
ABSTRACT			
Structured summary	2	Provide a structured summary that includes (as applicable): background, objectives, eligibility criteria, sources of evidence, charting methods, results, and conclusions that relate to the review questions and objectives.	2
INTRODUCTION			
Rationale	3	Describe the rationale for the review in the context of what is already known. Explain why the review questions/objectives lend themselves to a scoping review approach.	2-5
Objectives	4	Provide an explicit statement of the questions and objectives being addressed with reference to their key elements (e.g., population or participants, concepts, and context) or other relevant key elements used to conceptualize the review questions and/or objectives.	5-6

SECTION	ITEM	PRISMA-ScR CHECKLIST ITEM	REPORTED ON PAGE #
METHODS			
Protocol and registration	5	Indicate whether a review protocol exists; state if and where it can be accessed (e.g., a Web address); and if available, provide registration information, including the registration number.	8
Eligibility criteria	6	Specify characteristics of the sources of evidence used as eligibility criteria (e.g., years considered, language, and publication status), and provide a rationale.	9-10
Information sources*	7	Describe all information sources in the search (e.g., databases with dates of coverage and contact with authors to identify additional sources), as well as the date the most recent search was executed.	8, 83-35
Search	8	Present the full electronic search strategy for at least 1 database, including any limits used, such that it could be repeated.	83-85
Selection of sources of evidence†	9	State the process for selecting sources of evidence (i.e., screening and eligibility) included in the scoping review.	9-10
Data charting process‡	10	Describe the methods of charting data from the included sources of evidence (e.g., calibrated forms or forms that have been tested by the team before their use, and whether data charting was done independently or in duplicate) and any processes for obtaining and confirming data from investigators.	11
Data items	11	List and define all variables for which data were sought and any assumptions and simplifications made.	9-10
Critical appraisal of individual sources of evidence§	12	If done, provide a rationale for conducting a critical appraisal of included sources of evidence; describe the methods used and how this information was used in any data synthesis (if appropriate).	N/A
Synthesis of results	13	Describe the methods of handling and summarizing the data that were charted.	12-13
RESULTS			
Selection of sources of evidence	14	Give numbers of sources of evidence screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally using a flow diagram.	11-12

SECTION	ITEM	PRISMA-ScR CHECKLIST ITEM	REPORTED ON PAGE #
Characteristics of sources of evidence	15	For each source of evidence, present characteristics for which data were charted and provide the citations.	13-25

1477

1478

1479 **Appendix B: Full Search Strategies**

- 1480 • Initial searches run on 15.01.2021, 1970-2021 date filters used
- 1481 Updated searches run on 12.09.2022, 2021-2022 date filters used

1482

1483 ***PsychINFO (all databases):***

1484

1485 (((**Any Field:** ("compassion fatigue")) OR (**Any Field:** ("secondary traumatic
1486 stress")) OR (**Any Field:** (burnout)) OR (**Any Field:** (burn-out)) OR (**Any Field:** ("vicarious
1487 trauma")) OR (**Any Field:** ("occupational stress")) OR (**IndexTermsFilt:** ("Occupational
1488 Stress")) OR (**IndexTermsFilt:** ("Compassion Fatigue")) OR (**Any Field:** ("compassion
1489 stress")) OR (**Any Field:** "secondary trauma" OR (**Any Field:** "secondary
1490 traumatisation" OR (**Any Field:** "secondary traumatization") AND ((**Year:** [1970 TO
1491 2021]))) AND (((**AnyField:**(psychotherapist*) OR (**AnyField:**("forensic nurse")) OR
1492 (**AnyField:**("forensic psychologist")) OR (**AnyField:**("forensic psychiatrist")) OR
1493 (**AnyField:**(therapist*)) OR (**AnyField:**("support time and recovery worker")) OR
1494 (**AnyField:**("mental health employee")) OR (**AnyField:**(counsellor*)) OR
1495 (**AnyField:**("counselling psychologist")) OR (**AnyField:**("family therapist")) OR
1496 (**AnyField:**(counselor)) OR (**AnyField:**("mental health support worker")) OR
1497 (**AnyField:**("mental health social worker")) OR (**IndexTermsFilt:**("Mental Health
1498 Personnel")) OR (**IndexTermsFilt:**("Counselors")) OR OR (**AnyField:**("mental health care
1499 provider")) OR (**AnyField:**("mental health personnel")) OR (**AnyField:**("mental health
1500 nurse")) OR (**AnyField:**("mental health service")) OR (**AnyField:**("mental health service
1501 provider")) OR (**AnyField:**("mental health professional")) OR (**AnyField:**(psychologist*)) OR
1502 (**AnyField:**("mental health worker")))) AND ((**PublicationYear:**[1970 TO 2021] OR
1503 **TestYear:**[1970 TO 2021]))))

1504

1505 **Pubmed**

1506

1507 **Mesh search:**

1508

1509 ("social work, psychiatric"[MeSH Major Topic] OR "social work, psychiatric"[MeSH Major
1510 Topic] OR "emergency services, psychiatric"[MeSH Major Topic] OR "psychiatric
1511 nursing"[MeSH Major Topic] OR "health personnel/psychology"[MeSH Major Topic]) AND
1512 "burnout, professional/prevention and control"[MeSH Major Topic] OR "compassion
1513 fatigue/prevention and control"[MeSH Major Topic])

1514

1515 **Keyword search:**

1516

1517 ("psychotherapist*" [Title/Abstract] OR "forensic nurs*" [Title/Abstract] OR "forensic
1518 psychologist*" [Title/Abstract] OR "forensic psychiatrist*" [Title/Abstract] OR
1519 "therapist*" [Title/Abstract] OR "mental health employee*" [Title/Abstract] OR

1520 "counsellor*"[Title/Abstract] OR "counselling psychologist*"[Title/Abstract] OR "family
1521 therapist*"[Title/Abstract] OR "counselor*"[Title/Abstract] OR "mental health support
1522 worker*"[Title/Abstract] OR "mental health social worker*"[Title/Abstract] OR "mental
1523 health care provider*"[Title/Abstract] OR "mental health personnel"[Title/Abstract] OR
1524 "mental health nurs*"[Title/Abstract] OR "mental health service*"[Title/Abstract] OR
1525 "mental health service provider*"[Title/Abstract] OR "mental health
1526 professional*"[Title/Abstract] OR "psychologist*"[Title/Abstract] OR "mental health
1527 worker"[Title/Abstract])) AND ("compassion fatigue"[Title/Abstract] OR "vicarious
1528 trauma"[Title/Abstract] OR "secondary traumatic stress"[Title/Abstract] OR
1529 "burnout"[Title/Abstract] OR "burn out"[Title/Abstract] OR "compassion
1530 stress"[Title/Abstract] OR "secondary trauma"[Title/Abstract] OR "secondary
1531 traumatisation"[Title/Abstract] OR "secondary traumatization"[Title/Abstract])

1532

1533 *Embase*

1534

1535 • #1 ('compassion fatigue'/exp OR 'compassion fatigue' OR 'compassion stress' OR
1536 'vicarious trauma'/exp OR 'vicarious trauma' OR 'burnout'/exp OR burnout OR 'burn
1537 out'/exp OR 'burn out' OR 'secondary traumatic stress'/exp OR 'secondary traumatic
1538 stress' OR 'secondary trauma' OR 'secondary traumatisation' OR 'secondary
1539 traumatization') AND ([embase]/lim OR [embase classic]/lim) AND [1970-
1540 2021]/pyEBS & MHWs 08.01.212021-01-082021-01-0814187

1541

1542 • #2 ('vicarious trauma'/exp OR 'burnout'/exp OR 'professional burnout'/exp OR
1543 'compassion fatigue'/exp OR 'secondary traumatic stress'/exp) AND ([embase]/lim
1544 OR [embase classic]/lim) AND [1970-2021]/pyEBS & MHWs 08.01.212021-
1545 01-082021-01-0811764

1546 • #3 (psychotherapist* OR 'forensic nurse' OR 'forensic psychologist' OR
1547 'forensic psychiatrist'/exp OR 'forensic psychiatrist' OR therapist* OR
1548 'support time and recovery worker' OR 'mental health employee' OR
1549 counsellor* OR 'counselling psychologist' OR 'family therapist' OR
1550 'counselor'/exp OR counselor OR 'mental health support worker' OR
1551 'mental health social worker' OR 'mental health care provider' OR 'mental
1552 health personnel'/exp OR 'mental health personnel' OR 'mental health
1553 nurse' OR 'mental health service'/exp OR 'mental health service' OR
1554 'mental health service provider' OR 'mental health professional'/exp OR
1555 'mental health professional' OR psychologist* OR 'mental health
1556 worker'/exp OR 'mental health worker') AND ([embase]/lim OR [embase
1557 classic]/lim) AND [1970-2021]/pyEBS & MHWs 08.01.212021-01-
1558 082021-01-08121632

1559

1560

1561 • #4 ('mental health care personnel'/exp OR 'psychotherapist'/exp OR
1562 'counselor'/exp OR 'mental health service'/exp OR 'psychologist'/exp OR
1563 'psychiatric nurse'/exp OR 'forensic psychology'/exp) AND ([embase]/lim
1564 OR [embase classic]/lim) AND [1970-2021]/pyEBS & MHWs
1565 08.01.212021-01-082021-01-0856216

1566

1567 • #5 #1 OR #2

1567 • #6 #3 OR #4

1568 • #7 #5 AND #6

1569

1570 ***PTSDPubs:***

1571

1572 MAINSUBJECT.EXACT("Vicarious Traumatization") OR

1573 MAINSUBJECT.EXACT("Burnout") OR MAINSUBJECT.EXACT("Trauma Contagion")

1574 OR "compassion fatigue" OR "compassion stress" OR "secondary traumatic stress" OR

1575 "vicarious trauma" OR burnout OR burn-out

1576 AND

1577 MAINSUBJECT.EXACT("Mental Health Personnel") OR (psychotherapist* OR "forensic

1578 nurse" OR "forensic psychologist" OR "forensic psychiatrist" OR therapist* OR "support

1579 time and recovery worker" OR "mental health employee" OR counsellor* OR "counselling

1580 psychologist" OR "family therapist" OR counselor OR "mental health support worker" OR

1581 "mental health social worker" OR "mental health care provider" OR "mental health

1582 personnel" OR "mental health nurse" OR "mental health service" OR "mental health service

1583 provider" OR "mental health professional" OR psychologist*)

1584

1585 ***Central:***

1586

1587 "compassion fatigue" OR "compassion stress" OR "secondary traumatic stress" OR

1588 "vicarious trauma" OR burnout OR burn-out OR "Compassion Fatigue" OR "burnout,

1589 professional" OR "burnout, psychological" OR "burnout, professional/prevention and

1590 control" OR "compassion fatigue/prevention and control" in Title Abstract Keyword AND

1591 'social work, psychiatric' OR 'emergency services, psychiatric' OR 'psychiatric nursing' OR

1592 'health personnel, psychology' OR "psychotherapist*" OR "forensic nurs*" OR "forensic

1593 psychologist*" OR "forensic psychiatrist*" OR "therapist*" OR "mental health employee*"

1594 OR "counsellor*" OR "counselling psychologist*" OR "family therapist*" OR "counselor*"

1595 OR "mental health support worker*" OR "mental health social worker*" OR "mental health

1596 care provider*" OR "mental health personnel" OR "mental health nurs*" OR "mental health

1597 service*" OR "mental health service provider*" OR "mental health professional*" OR

1598 "psychologist*" OR "mental health worker" in Title Abstract Keyword - with Cochrane

1599 Library publication date Between Jan 1970 and Jan 2021

1600

1601

1602

1603

1604

1605

1606

1607

1608 **Appendix C**1609 *Table showing quantitative measures of EBS used*

1610

Measure	Description	No. of Studies Used In
The Maslach Burnout Inventory (MBI) (Maslach & Jackson, 1981).	The MBI is a 22-item self-report measure of burnout in human services professionals. It produces 3 dimensions of burnout: emotional exhaustion, depersonalization, and personal accomplishment. It is the most commonly used measure of burnout with high reliability ($\alpha = .71$ or higher) across 7 countries (Poghosyan et al., 2009).	24
The Dutch version of the MBI (DMBI) (Schaufeli, 1990).	The questionnaire consists of 22 items on a 7-point Likert scale. The Dutch MBI has been shown to be valid and reliable in the assessment of burnout in nurses (Schaufeli & van Dierendonck, 1993).	1
The Arabic version of the MBI (Al-Turki et al., 2010; Hamaideh, 2011).	The Arabic MBI has been used in a number of Arabic-speaking countries, in studies assessing burnout levels among nurses (Al-Turki et al., 2010; Hamaideh, 2011). It has been shown to have good internal consistency across subscales: $\alpha = .80$ for emotional exhaustion; $\alpha = .73$ for depersonalisation; $\alpha = .77$ for personal accomplishment (Sabbah et al., 2012).	1
The Italian version of the MBI (Sirigatti & Stefanile, 1993).	This Italian version of the MBI has been shown to have good internal consistency across subscales: $\alpha = .88$ for emotional exhaustion; $\alpha = .70$ for depersonalisation; $\alpha = .83$ for personal realisation (Maslach & Jackson, 1981; Sirigatti & Stefanile, 1993).	1
MBI-General Survey (MBI-GS) (Schaufeli et al., 1996).	The MBI-GS is a 16-item self-report measure of burnout in any occupation (not just health/care). The generic items measure burnout on 3 subscales that parallel the MBI: exhaustion, cynicism, and	1

Measure	Description	No. of Studies Used In
	professional efficacy. The MBI-GS has been shown to have satisfactory internal consistency ($\alpha = .84$ to $.90$ for exhaustion, $.74$ to $.84$ for cynicism, and from $.70$ to $.78$ for professional efficacy) (Leiter & Schaufeli, 1996).	
The Copenhagen Burnout Inventory (CBI) (Kristensen et al., 2005).	The CBI is a 19-item self-report tool that produces 3 scales measuring personal burnout, work-related burnout, and client-related burnout. It is designed for use with employees in any sector. A prospective study of burnout in human service workers sector found that all 3 scales had high internal reliability $\alpha = .87$ for both personal and work-related burnout; $\alpha = 0.85$ for client-related burnout) (Borritz et al., 2006).	3
The Oldenburg Burnout Inventory (OLBI) (Demerouti & Nachreiner, 1998).	The OLBI is a self-report measure consisting of 16 positively and negatively formulated items that are used to evaluate 2 dimensions of burnout: the identification continuum (ranging from disengagement to dedication) and the energy continuum (ranging from exhaustion to vigour). Studies have demonstrated the convergent validity of the OLBI and the MBI-General Survey (Demerouti et al., 2003; Halbesleben & Demerouti, 2005). The reliability of the exhaustion subscale ranges from $\alpha = .74$ to $\alpha = .85$, and the reliability of the disengagement subscale from $.73$ to $.85$ (Demerouti & Bakker, 2008; Demerouti et al., 2003; Halbesleben & Demerouti, 2005, Halbesleben, 2010; Sonnentag, Binnewies & Mojza, 2010; Timms et al., 2012).	2
Silencing Response Scale (SRS) (Baranowsky, 2002).	The SRS is a 15-item scale that helps care-givers identify specific communication difficulties in their trauma work with clients i.e. selective listening and active avoidance. It has been shown to have good reliability, $\alpha = .85$ (Ortlepp & Friedman, 2001).	1
The Trauma Recovery Scale (Gentry, 1996).	The TRS is a 10-item self-report scale. Parts I & II provide a traumatic experiences inventory. Part III measures relative recovery from traumatic experiences. It is found to have good internal reliability ($\alpha = .86$) and good convergent validity with the impact of events scale ($.71$). (Gentry, 1996).	1
Secondary Traumatic Stress Scale (STSS) (Bride et al., 2004).	The STSS is a 17-item self-report measure of secondary trauma in practitioners who have experienced traumatic stress through their work with clients. It comprises 3 subscales; Intrusion, Avoidance, and Arousal. It was found to have good internal consistency, ranging from $\alpha = .93$ - $.94$ for the total STSS	1

Measure	Description	No. of Studies Used In
The Impact of Events Scale (IES) (Horowitz et al., 1979).	<p>scale; $\alpha = .79-80$ for the Intrusion subscale, $\alpha = .85-87$ for the Avoidance subscale, and $\alpha = .83-87$ for the Arousal subscale (Bride et al., 2004; Ting et al., 2005).</p> <p>The IES is a 15-item self-report measure of current subjective distress for any specific life event. It measures distress along two subscales: Intrusive and Avoidance symptoms. The IES has shown very good internal consistency ($r = .79$ to $.92$, with an average of $.86$ for the intrusive subscale and $.90$ for the avoidance subscale) (Corcoran and Fischer, 1994). The split-half reliability of the IES scale was high ($r = .86$) and internal consistency of the subscales was also high: $\alpha = .78$ for intrusion and $\alpha = .82$ for avoidance. Test-retest reliability was high: $r = .87$ for the total stress scores; $.89$ for the intrusion subscale; and $.79$ for the avoidance subscale (Horowitz et al., 1979).</p>	1
The questionnaire on the perception of occupational burnout (Penko, 1994).	According to the information in Kovač et al (2016), this questionnaire consists of 3 dimensions of occupational burnout: less fulfilment, emotional exhaustion and depersonalization. No further information was accessible in English regarding this scale.	6
Professional Quality of Life Scale version 5 (ProQOL-5) (Stamm, 2009).	The ProQOL-5 is a 30 item self-report measure of the positive and negative effects of working with people who have experienced extremely stressful events. It comprises of 3 subscales measuring Compassion Fatigue, Burnout and Compassion Satisfaction. CS alpha reliabilities = $.87$ BO alpha reliabilities = $.72$ CF alpha reliabilities = $.80$. Early information on test-retest data show good reliability.	1
Swedish version of the Shirom Melamed Burnout Questionnaire (SMBQ) (Grossi et al., 2003)	The SMBQ (originally by Melamed et al., 1999) is a 22-item self-report burnout measure consisting of 4 subscales: Physical Fatigue; Cognitive weariness; Tension and Listlessness. The Swedish version was shown to have consistencies ($\alpha > 0.70$) across the subscales, but a Rasch-analysis suggested omission of the Tension factor (Lundgren-Nilsson et al., 2012).	1

Measure	Description	No. of Studies Used In
Compassion Satisfaction/Fatigue Self-Test (CSFST) (Figley & Stamm, 1996).	The CSFT is a 66-item self-report measure comprised of three subscales: compassion fatigue, compassion satisfaction, and burnout. According to Stamm (2002), all 3 subscales have demonstrated good alpha reliability: $\alpha = .87$ on the compassion fatigue subscale; $\alpha = .90$ on the burnout subscale; and $\alpha = .87$ for the compassion satisfaction subscale.	3
Well-being Index (WBI) (Dyrbye et al., 2014)	The WBI is a seven item self-report index, measuring domains of burnout, depression, stress, fatigue, and mental and physical quality of life. It consists of yes/no items and respondents receive a score from 0 to 7 based on responses. At a threshold score of ≥ 4 , the PWBI's sensitivity for identifying physicians with low mental quality of life is 73.3 % with a specificity of 81.0 % (Dyrbye et al. 2013).	1
Stanford Professional Fulfilment Index (PFI) (Trockel et al., 2018)	The PFI is a 16 item self-report measure assessing the degree of intrinsic positive reward the individual derives from his or her work, including happiness, meaningfulness, contribution, self-worth, satisfaction, and feeling in control when dealing with difficult problems at work. It consists of 3 scales- professional fulfillment; work exhaustion; and interpersonal disengagement and an overall burnout scale. Response options are on a five-point Likert scale ("not at all true" to "completely true" for professional fulfillment items and "not at all" to "extremely" for work exhaustion and interpersonal disengagement items). Each PFI item is scored from 0 to 4, using the associated five-point Likert scale. All scales have demonstrated good alpha reliability: $\alpha = .86$ on the work exhaustion scale; $\alpha = .92$ on the interpersonal disengagement scale; $\alpha = .91$ on the professional fulfillment scale; $\alpha = .92$ on the overall burnout scale (Trockel et al., 2018).	1
Stress Rating Form & Burnout Adjectives (described in Ray, 1981).	The stress rating form was devised for the study to collect pre- and post-ratings on participants by one close friend/relative and co-workers or supervisor. The observer was asked to rate the amount of stress that he/she believed that the subject was currently experiencing in five areas (home, job interpersonal relationships, health, and school), and to rate the amount of burnout the subject was feeling, on a 1 to 10 scale, with 1 being little stress and 10 being highly stressed. Additionally, the observer rated the subject on 16 adjectives on a 1 to 10-point scale, in order to compute an overall burnout score (the method for this is reportedly described in Thompson (1980) but this could not be accessed by reviewers).	1

1611 **Appendix D**

1612 *Table showing recommendations for future research*


Intervention	Recommendations/ Questions for future research
DBT skills training	<ul style="list-style-type: none"> • Larger and more diverse samples; control groups • Direct patient-related outcomes • Compare effectiveness of person-directed interventions, versus organizational interventions only, versus a combination. • Would longer training and supervision groups increase effectiveness? • Include clinician burnout as a variable within DBT effectiveness and efficacy studies • Account/control for different settings differing initial levels of clinician burnout • Account for variations in learning and utilization of training • Administer questionnaires to assess subjects' knowledge of and utilization of DBT.
PSI training	<ul style="list-style-type: none"> • Focus on PSI competencies • Longer-term evaluation of implementation in practice • Outcome-based intervention trials • Larger samples, different settings, more rigorous methodologies
ACT training	<ul style="list-style-type: none"> • Compare the impact of brief interventions based on ACT, psychoeducation, or an amalgam of both. • Improved rigour
Supervision	<ul style="list-style-type: none"> • Larger samples, inclusion of different settings/specific roles of participants. • What structure of systematic clinical supervision is most effective? • Longer duration of study/supervision may be required to see changes in burnout • Incorporate qualitative and quantitative data to measure STS responses. • Can supervision serve as a preventative strategy for MHWs who do not yet have STS responses?


Intervention	Recommendations/ Questions for future research
EBS training/ prevention	<ul style="list-style-type: none"> • Larger samples, collect demographic data • Separate the various components of the intervention for comparison • Longer and more intensive STS education program • Follow up 2-3 months later, giving MHWs time to implement/ incorporate STS prevention plans • Exploring the effects of intermittent booster sessions of the intervention • More rigorous designs e.g. multiple baseline/ wait-list control groups, randomization • Measure primary traumatic stress and burnout • Participant interviews to distill the active elements of interventions
Resiliency training	<ul style="list-style-type: none"> • More CF exploratory longitudinal studies with treatment and control groups • Additional CF research exploring CF over the course of clinicians' careers • Comparing CF symptoms between CITs and practicing mental health professionals • Qualitative approaches exploring lived experiences with CF symptoms as well as during and after CF treatments to determine active ingredients for meaningful CF symptom reduction.
Mindfulness/ compassion	<ul style="list-style-type: none"> • Are MBIs alone sufficient to generate self-compassion? • Increased rigour: large, controlled studies; random sampling across settings; long term follow up • Explore possible applications to other disciplines/ health workers • Explore mechanisms through which the Interpersonal Mindfulness Program works • Explore the adverse reactions reported in one study (Bartels-Velthuis et al., 2020).
'BREATHE' burnout reduction workshop	<ul style="list-style-type: none"> • Separate BREATHE content into smaller modules delivered over time • Trial additional electronic resources and reminders outside the workshop setting. • Revise BREATHE materials to specifically address how to cope with job demands that promote burnout. • Target interventions to MHWs experiencing higher levels of burnout. • Randomised control groups and longer follow-up periods • Ensure acceptability and feasibility with racial and ethnic minority groups
Expressive arts	<ul style="list-style-type: none"> • Recruit participants from a variety of settings to examine individual differences


Intervention	Recommendations/ Questions for future research
Peer support/ relationships	<ul style="list-style-type: none"> • Use a wider variety of research methods to explore impact • Investigate the precise effects of group drumming on factors associated with burnout <ul style="list-style-type: none"> • Longitudinal designs with follow-up to explore longevity of effects. • Compare: online and face-to-face peer groups; hybrid models that include both online and face-to-face components. • Focus on both process and outcome measures • Control groups and larger samples. • Recruit participants with higher levels of emotional exhaustion to explore impact on this aspect of burnout more specifically
Balint-like group Training in other therapeutic interventions (CBT; psychosis/BPD interventions).	<ul style="list-style-type: none"> • Larger sample sizes with more statistical power • More rigorous experimental designs • Does training result in improved patient outcomes? • Will results generalise to other mental health providers? • Do reductions in burnout for MHWs attending training sustain over time?
Wellness/stress reduction	<ul style="list-style-type: none"> • Recruit homogeneous samples or else stratify more varied samples by job classification • Larger samples, randomization to groups • Explore differences between single and partnered MHWs, including experience and/or frequency of touch in their lives.
	•

1613 *Note.* ACT = acceptance and commitment therapy; BREATHE = Burnout Reduction: Enhanced Awareness, Tools, Hand-outs, and Education; CF =
1614 compassion fatigue; CITs = counsellors in training; DBT= dialectical behaviour therapy; MBIs= mindfulness-based interventions; MHWs= mental
1615 health workers; PSI = psychosocial interventions; STS = secondary traumatic stress

		Primary	Secondary	Tertiary
Individual	Personal practice		<ul style="list-style-type: none"> Wellness/stress reduction Mindfulness ACT Compassion enhancement Expressive arts Aspects of the BREATHE intervention and resiliency workshops 	
	Awareness raising		<ul style="list-style-type: none"> EBS prevention Aspects of the BREATHE intervention and resiliency workshops 	
	Role training		<ul style="list-style-type: none"> Training in a therapeutic modality (DBT, PSI, CBT) 	
Team			<ul style="list-style-type: none"> Supervision Peer groups/relationships Balint-like group 	
Organisational		<ul style="list-style-type: none"> Primary nursing 		







Click here to access/download
Supporting Information
PRISMA Chart.docx